



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 07:13 am GMT

PDB ID : 2CLZ
Title : Mhc Class I Natural Mutant H-2Kbm8 Heavy Chain Complexed With beta-2 Microglobulin and pBM1 peptide
Authors : Mazza, C.; Auphan-Anezin, N.; Guimezanes, A.; Barrett-Wilt, G.A.; Montero-Julian, F.; Roussel, A.; Hunt, D.F.; Schmitt-Verhulst, A.M.; Malissen, B.
Deposited on : 2006-05-03
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	trunk30967
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	5502 (1.90-1.90)
Clashscore	122126	6115 (1.90-1.90)
Ramachandran outliers	120053	6048 (1.90-1.90)
Sidechain outliers	120020	6048 (1.90-1.90)
RSRZ outliers	108989	5379 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	<div> <div>5%</div> <div> <div></div> <div>90%</div> <div>10%</div> </div> </div>
1	H	279	<div> <div>5%</div> <div> <div></div> <div>87%</div> <div>12%</div> </div> </div>
2	B	99	<div> <div>%</div> <div> <div></div> <div>91%</div> <div>8%</div> </div> </div>
2	P	99	<div> <div>%</div> <div> <div></div> <div>86%</div> <div>13%</div> </div> </div>
3	C	8	<div> <div></div> <div>100%</div> </div>
3	M	8	<div> <div></div> <div>75%</div> <div>25%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6939 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 CLASS I HISTOCOMPATIBILITY ANTIGEN, K-B ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	279	Total	C	N	O	S	0	0	1
			2258	1427	399	424	8			
1	H	279	Total	C	N	O	S	0	0	1
			2258	1427	399	424	8			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	PHE	TYR	engineered mutation	UNP P01901
A	23	ILE	MET	engineered mutation	UNP P01901
A	24	SER	GLU	engineered mutation	UNP P01901
A	30	ASN	ASP	engineered mutation	UNP P01901
H	22	PHE	TYR	engineered mutation	UNP P01901
H	23	ILE	MET	engineered mutation	UNP P01901
H	24	SER	GLU	engineered mutation	UNP P01901
H	30	ASN	ASP	engineered mutation	UNP P01901

- Molecule 2 is a protein called BETA-2 MICROGLOBULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	0	0
			821	524	138	152	7			
2	P	99	Total	C	N	O	S	0	0	0
			821	524	138	152	7			

- Molecule 3 is a protein called RBM5 PROTEIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	8	Total	C	N	O	0	0	0
			70	46	10	14			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	M	8	Total	C	N	O	0	0	0
			70	46	10	14			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	222	Total	O	0	0
			222	222		
4	B	92	Total	O	0	0
			92	92		
4	C	10	Total	O	0	0
			10	10		
4	H	208	Total	O	0	0
			208	208		
4	M	7	Total	O	0	0
			7	7		
4	P	102	Total	O	0	0
			102	102		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 CLASS I HISTOCOMPATIBILITY ANTIGEN, K-B ALPHA CHAIN




- Molecule 1: H-2 CLASS I HISTOCOMPATIBILITY ANTIGEN, K-B ALPHA CHAIN



- Molecule 2: BETA-2 MICROGLOBULIN



- Molecule 2: BETA-2 MICROGLOBULIN




- Molecule 3: RBM5 PROTEIN



There are no outlier residues recorded for this chain.

● Molecule 3: RBM5 PROTEIN

Chain M:  75% 25%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.59Å 90.76Å 89.25Å 90.00° 111.35° 90.00°	Depositor
Resolution (Å)	15.00 – 1.90 50.97 – 1.90	Depositor EDS
% Data completeness (in resolution range)	96.7 (15.00-1.90) 96.6 (50.97-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.206 , 0.247 0.218 , 0.259	Depositor DCC
R_{free} test set	3788 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.595	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.019 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6939	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.24 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.7960e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/2322	0.59	0/3158
1	H	0.45	0/2322	0.60	0/3158
2	B	0.45	0/847	0.58	0/1148
2	P	0.46	0/847	0.60	0/1148
3	C	0.57	0/71	0.56	0/94
3	M	0.50	0/71	0.50	0/94
All	All	0.46	0/6480	0.59	0/8800

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2258	0	2153	24	0
1	H	2258	0	2153	30	0
2	B	821	0	796	6	0
2	P	821	0	796	10	0
3	C	70	0	65	0	0
3	M	70	0	65	2	0
4	A	222	0	0	3	0
4	B	92	0	0	0	0
4	C	10	0	0	0	0
4	H	208	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	M	7	0	0	0	0
4	P	102	0	0	1	0
All	All	6939	0	6028	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:GLU:HB2	1:H:138:MET:HE1	1.10	1.09
1:H:108:ARG:HG3	1:H:108:ARG:HH11	1.18	1.01
1:A:61:GLU:HB2	1:H:138:MET:CE	1.93	0.96
1:A:111:ARG:HG2	1:A:111:ARG:HH11	1.28	0.95
2:P:3:LYS:HE3	4:P:2066:HOH:O	1.73	0.89
1:A:61:GLU:CB	1:H:138:MET:HE1	2.04	0.78
1:H:108:ARG:HH11	1:H:108:ARG:CG	1.97	0.76
1:A:111:ARG:CG	1:A:111:ARG:HH11	2.01	0.72
2:P:21:ASN:HD22	2:P:22:ILE:H	1.38	0.70
2:P:13:HIS:H	2:P:21:ASN:HD21	1.41	0.68
1:H:50:ARG:HH11	1:H:50:ARG:HG3	1.59	0.68
1:H:263:HIS:CD2	1:H:265:GLY:H	2.13	0.66
1:A:111:ARG:HG2	1:A:111:ARG:NH1	2.07	0.65
1:A:258:THR:HG22	1:A:273:ARG:HE	1.62	0.64
2:P:37:ILE:HD12	2:P:51:MET:HE1	1.80	0.63
1:A:82:LEU:HD21	1:A:89:LYS:HA	1.81	0.62
1:H:81:LEU:HD13	1:H:118:TYR:CD1	2.36	0.61
1:A:108:ARG:HD3	4:A:2102:HOH:O	2.01	0.60
1:H:81:LEU:HG	3:M:8:ILE:HD11	1.82	0.59
1:A:258:THR:OG1	1:A:260:HIS:HE1	1.89	0.56
1:H:108:ARG:HG3	1:H:108:ARG:NH1	1.99	0.55
1:H:230:LEU:HD22	1:H:243:LYS:HE3	1.89	0.55
1:A:263:HIS:CD2	1:A:265:GLY:H	2.25	0.54
1:H:4:SER:HB3	1:H:102:GLU:HG2	1.90	0.54
1:H:234:ARG:HE	1:H:242:GLN:HE21	1.55	0.52
1:A:14:ARG:HB3	1:A:17:LEU:HB2	1.94	0.49
1:H:258:THR:HG22	1:H:273:ARG:HD3	1.95	0.49
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.47	0.49
1:H:6:ARG:NH2	4:H:2005:HOH:O	2.44	0.49
1:A:234:ARG:HE	1:A:242:GLN:HE21	1.61	0.49
4:A:2060:HOH:O	1:H:138:MET:HE2	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:263:HIS:HD2	1:H:265:GLY:H	1.59	0.48
2:P:17:ASN:HD21	2:P:97:ARG:HH22	1.61	0.48
2:P:21:ASN:ND2	2:P:22:ILE:H	2.09	0.47
1:A:6:ARG:NH2	1:A:30:ASN:OD1	2.47	0.47
1:A:260:HIS:HA	1:A:270:LEU:O	2.15	0.47
1:A:258:THR:OG1	1:A:260:HIS:CE1	2.67	0.47
1:H:210:PRO:O	1:H:263:HIS:HE1	1.98	0.47
1:H:50:ARG:HH11	1:H:50:ARG:CG	2.26	0.46
1:H:117:ALA:HB2	2:P:60:TRP:CE2	2.50	0.46
1:H:108:ARG:NH1	1:H:108:ARG:CG	2.67	0.46
1:A:11:ALA:HA	1:A:21:ARG:O	2.17	0.45
1:A:94:THR:HG21	2:B:33:PRO:HD3	1.99	0.45
2:B:40:LEU:HD12	2:B:45:LYS:HA	1.98	0.44
2:P:24:ASN:HD22	2:P:67:HIS:HB3	1.83	0.44
1:A:210:PRO:O	1:A:263:HIS:HE1	2.01	0.44
1:A:234:ARG:HH11	2:B:8:GLN:NE2	2.16	0.44
1:A:6:ARG:NH1	4:A:2006:HOH:O	2.23	0.43
2:P:21:ASN:HD22	2:P:22:ILE:N	2.13	0.43
1:H:216:THR:OG1	1:H:260:HIS:HB2	2.19	0.42
1:A:81:LEU:HD13	1:A:118:TYR:CD1	2.54	0.42
1:H:275:GLU:HA	1:H:276:PRO:HD2	1.97	0.41
2:B:12:ARG:CZ	2:B:22:ILE:HD13	2.50	0.41
1:H:4:SER:CB	1:H:102:GLU:HG2	2.50	0.41
1:H:234:ARG:HH11	2:P:8:GLN:NE2	2.17	0.41
1:H:6:ARG:NH1	1:H:30:ASN:OD1	2.53	0.41
1:A:48:ARG:NH2	2:B:53:ASP:OD1	2.51	0.41
1:H:62:ARG:HE	1:H:62:ARG:HB3	1.56	0.41
1:H:81:LEU:HD22	1:H:85:TYR:HE2	1.85	0.41
1:H:258:THR:HG22	1:H:273:ARG:CD	2.50	0.41
1:H:97:VAL:HG11	3:M:5:PHE:CE2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	277/279 (99%)	268 (97%)	8 (3%)	1 (0%)	36	25
1	H	277/279 (99%)	270 (98%)	7 (2%)	0	100	100
2	B	97/99 (98%)	95 (98%)	2 (2%)	0	100	100
2	P	97/99 (98%)	95 (98%)	2 (2%)	0	100	100
3	C	6/8 (75%)	6 (100%)	0	0	100	100
3	M	6/8 (75%)	5 (83%)	1 (17%)	0	100	100
All	All	760/772 (98%)	739 (97%)	20 (3%)	1 (0%)	53	44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/237 (100%)	232 (98%)	4 (2%)	63	60
1	H	236/237 (100%)	222 (94%)	14 (6%)	21	11
2	B	94/94 (100%)	92 (98%)	2 (2%)	56	51
2	P	94/94 (100%)	91 (97%)	3 (3%)	42	33
3	C	8/8 (100%)	8 (100%)	0	100	100
3	M	8/8 (100%)	8 (100%)	0	100	100
All	All	676/678 (100%)	653 (97%)	23 (3%)	40	30

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	111	ARG
1	A	114	GLN
1	A	194	ARG
2	B	40	LEU
2	B	70	PHE
1	H	21	ARG
1	H	45	TYR
1	H	50	ARG
1	H	62	ARG
1	H	108	ARG
1	H	109	LEU
1	H	141	LEU
1	H	163	THR
1	H	166	GLU
1	H	180	LEU
1	H	181	ARG
1	H	186	LYS
1	H	201	LEU
1	H	224	LEU
2	P	21	ASN
2	P	48	LYS
2	P	70	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	GLN
1	A	174	ASN
1	A	218	GLN
1	A	242	GLN
1	A	260	HIS
1	A	263	HIS
2	B	2	GLN
2	B	8	GLN
1	H	3	HIS
1	H	127	ASN
1	H	242	GLN
1	H	263	HIS
2	P	8	GLN
2	P	17	ASN
2	P	21	ASN
2	P	24	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	P	38	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	279/279 (100%)	0.19	14 (5%) 29 32	14, 24, 38, 43	0
1	H	279/279 (100%)	0.19	14 (5%) 29 32	13, 25, 38, 45	0
2	B	99/99 (100%)	-0.06	1 (1%) 82 84	15, 23, 32, 43	0
2	P	99/99 (100%)	-0.04	1 (1%) 82 84	15, 23, 32, 40	0
3	C	8/8 (100%)	0.17	0 100 100	17, 22, 26, 26	0
3	M	8/8 (100%)	-0.02	0 100 100	21, 25, 31, 31	0
All	All	772/772 (100%)	0.13	30 (3%) 39 43	13, 24, 37, 45	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	225	ILE	5.4
1	H	278	PRO	5.3
1	A	225	ILE	4.3
2	B	1	ILE	4.3
1	A	42	ASN	3.9
2	P	1	ILE	3.8
1	A	278	PRO	3.7
1	A	41	GLU	3.5
1	H	41	GLU	3.4
1	H	279	SER	3.3
1	A	89	LYS	3.3
1	H	111	ARG	2.8
1	A	138	MET	2.7
1	A	267	PRO	2.6
1	A	195	PRO	2.6
1	A	16	GLY	2.5
1	H	220	ASN	2.5
1	H	16	GLY	2.4
1	H	277	PRO	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	220	ASN	2.4
1	A	150	ALA	2.4
1	H	145	HIS	2.3
1	H	195	PRO	2.3
1	A	226	GLN	2.2
1	A	149	GLN	2.2
1	H	149	GLN	2.2
1	H	226	GLN	2.2
1	H	42	ASN	2.1
1	H	40	ALA	2.1
1	A	275	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.