



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 10:35 pm GMT

PDB ID : 4D0Z
Title : GalNAc-T2 crystal soaked with UDP-5SGalNAc, mEA2 and manganese (Higher resolution dataset)
Authors : Lira-Navarrete, E.; Iglesias-Fernandez, J.; Zandberg, W.F.; Companon, I.; Kong, Y.; Corzana, F.; Pinto, B.M.; Clausen, H.; Peregrina, J.M.; Vocadlo, D.; Rovira, C.; Hurtado-Guerrero, R.
Deposited on : 2014-04-30
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4343 (2.20-2.20)
Clashscore	122126	5027 (2.20-2.20)
Ramachandran outliers	120053	4952 (2.20-2.20)
Sidechain outliers	120020	4953 (2.20-2.20)
RSRZ outliers	108989	4245 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	571	<div> <div>3%</div> <div> <div>71%</div> <div>13%</div> <div>•</div> <div>13%</div> </div> </div>
1	B	571	<div> <div>2%</div> <div> <div>73%</div> <div>12%</div> <div>•</div> <div>13%</div> </div> </div>
1	C	571	<div> <div>33%</div> <div> <div>64%</div> <div>15%</div> <div>• •</div> <div>17%</div> </div> </div>
1	D	571	<div> <div>4%</div> <div> <div>70%</div> <div>14%</div> <div>•</div> <div>13%</div> </div> </div>
1	E	571	<div> <div>2%</div> <div> <div>71%</div> <div>12%</div> <div>•</div> <div>13%</div> </div> </div>
1	F	571	<div> <div>15%</div> <div> <div>71%</div> <div>13%</div> <div>•</div> <div>14%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	X	6	
2	Y	6	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	A	1572	-	-	X	-
4	EDO	B	1573	-	-	X	-
4	EDO	D	1576	-	-	X	-
4	EDO	E	1572	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 24773 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called POLYPEPTIDE N-ACETYLGALACTOSAMINYLTRANSFERASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	495	Total	C	N	O	S	20	1	0
			3975	2502	722	727	24			
1	B	495	Total	C	N	O	S	20	0	0
			3967	2497	719	727	24			
1	C	476	Total	C	N	O	S	20	0	0
			3826	2407	693	702	24			
1	D	495	Total	C	N	O	S	20	0	0
			3967	2497	719	727	24			
1	E	495	Total	C	N	O	S	20	0	0
			3967	2497	719	727	24			
1	F	491	Total	C	N	O	S	20	0	0
			3926	2470	710	722	24			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	516	ASP	ASN	engineered mutation	UNP Q10471
B	516	ASP	ASN	engineered mutation	UNP Q10471
C	516	ASP	ASN	engineered mutation	UNP Q10471
D	516	ASP	ASN	engineered mutation	UNP Q10471
E	516	ASP	ASN	engineered mutation	UNP Q10471
F	516	ASP	ASN	engineered mutation	UNP Q10471


- Molecule 2 is a protein called PEPTIDE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	X	6	Total	C	N	O	S	0	0	1
			32	18	6	7	1			
2	Y	6	Total	C	N	O	S	0	0	1
			32	18	6	7	1			

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total Mn 1 1	0	0
3	E	1	Total Mn 1 1	0	0
3	B	1	Total Mn 1 1	0	0
3	C	1	Total Mn 1 1	0	0
3	A	1	Total Mn 1 1	0	0
3	F	1	Total Mn 1 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is 2-(acetylamino)-2-deoxy-5-thio-alpha-D-galactopyranose (three-letter code: BBK) (formula: C₈H₁₅NO₅S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			15	8	1	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	S	0	0
			15	8	1	5	1		
5	D	1	Total	C	N	O	S	0	0
			15	8	1	5	1		
5	E	1	Total	C	N	O	S	0	0
			15	8	1	5	1		

- Molecule 6 is (2R,3R,4R,5R,6R)-3-(acetylamino)-4,5-dihydroxy-6-(hydroxymethyl)tetrahydro-2H-thiopyran-2-yl [(2R,3S,4R,5R)-5-(2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)-3,4-dihydroxytetrahydrofuran-2-yl]methyl dihydrogen diphosphate (three-letter code: HWU) (formula: C₁₇H₂₇N₃O₁₆P₂S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
6	A	1	Total 39	C 17	N 3	O 16	P 2	S 1	0	0
6	B	1	Total 39	C 17	N 3	O 16	P 2	S 1	0	0
6	C	1	Total 39	C 17	N 3	O 16	P 2	S 1	0	0
6	D	1	Total 39	C 17	N 3	O 16	P 2	S 1	0	0
6	E	1	Total 39	C 17	N 3	O 16	P 2	S 1	0	0
6	F	1	Total 39	C 17	N 3	O 16	P 2	S 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	144	Total 144	O 144	0	0
7	B	125	Total 125	O 125	0	0
7	C	13	Total 13	O 13	0	0
7	D	122	Total 122	O 122	0	0
7	E	168	Total 168	O 168	0	0
7	F	61	Total 61	O 61	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: POLYPEPTIDE N-ACETYL GALACTOSAMINYLTRANSFERASE 2




• Molecule 1: POLYPEPTIDE N-ACETYL GALACTOSAMINYLTRANSFERASE 2




• Molecule 1: POLYPEPTIDE N-ACETYL GALACTOSAMINYLTRANSFERASE 2







● Molecule 1: POLYPEPTIDE N-ACETYL GALACTOSAMINYLTRANSFERASE 2



● Molecule 2: PEPTIDE



● Molecule 2: PEPTIDE





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	116.48Å 121.14Å 249.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	249.39 – 2.20 19.95 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (249.39-2.20) 99.8 (19.95-2.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 2.19Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.196 , 0.234 0.204 , 0.240	Depositor DCC
R_{free} test set	4936 reflections (2.77%)	wwPDB-VP
Wilson B-factor (Å ²)	38.8	Xtriage
Anisotropy	0.540	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.016 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	24773	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HWU, BBK, EDO, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	4/4070 (0.1%)	0.98	25/5503 (0.5%)
1	B	0.79	3/4059 (0.1%)	0.91	9/5489 (0.2%)
1	C	0.88	7/3912 (0.2%)	0.98	26/5287 (0.5%)
1	D	1.01	3/4059 (0.1%)	0.98	19/5489 (0.3%)
1	E	0.92	3/4059 (0.1%)	0.98	19/5489 (0.3%)
1	F	0.89	3/4015 (0.1%)	0.90	12/5427 (0.2%)
2	X	1.42	0/32	1.09	0/44
2	Y	1.82	1/32 (3.1%)	1.53	0/44
All	All	0.89	24/24238 (0.1%)	0.95	110/32772 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	F	0	1
All	All	0	4

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	75	LYS	CB-CG	-33.78	0.61	1.52
1	C	84	GLU	CB-CG	-26.41	1.01	1.52
1	D	486	LYS	CB-CG	-25.36	0.84	1.52
1	F	75	LYS	CB-CG	-25.00	0.85	1.52
1	E	84	GLU	CB-CG	-23.77	1.06	1.52

The worst 5 of 110 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	75	LYS	CA-CB-CG	21.27	160.19	113.40
1	D	194	ARG	NE-CZ-NH2	-17.68	111.46	120.30
1	E	75	LYS	CA-CB-CG	16.43	149.55	113.40
1	F	75	LYS	CA-CB-CG	15.46	147.41	113.40
1	D	75	LYS	CB-CG-CD	13.48	146.66	111.60

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	89	GLY	Peptide
1	B	89	GLY	Peptide
1	C	223	LEU	Mainchain
1	F	89	GLY	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3975	0	3903	54	0
1	B	3967	0	3890	30	0
1	C	3826	0	3748	81	0
1	D	3967	0	3890	55	0
1	E	3967	0	3890	49	0
1	F	3926	0	3849	31	0
2	X	32	0	28	3	0
2	Y	32	0	28	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
4	A	40	0	60	9	0
4	B	32	0	48	6	0
4	D	20	0	30	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	56	0	84	15	0
5	A	15	0	15	3	0
5	B	15	0	14	4	0
5	D	15	0	14	4	0
5	E	15	0	13	5	0
6	A	39	0	24	0	0
6	B	39	0	24	2	0
6	C	39	0	25	11	0
6	D	39	0	25	4	0
6	E	39	0	25	2	0
6	F	39	0	24	2	0
7	A	144	0	0	0	0
7	B	125	0	0	3	0
7	C	13	0	0	6	0
7	D	122	0	0	3	0
7	E	168	0	0	3	0
7	F	61	0	0	0	0
All	All	24773	0	23651	298	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 298 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:203:GLY:HA3	1:C:330:VAL:CG2	1.71	1.19
1:C:203:GLY:CA	1:C:330:VAL:CG2	2.26	1.13
1:A:291:ARG:CZ	1:D:435:PRO:HB2	1.79	1.11
1:C:203:GLY:HA3	1:C:330:VAL:HG22	1.20	1.10
1:C:330:VAL:HG12	1:C:331:TRP:HB3	1.32	1.08

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	494/571 (86%)	479 (97%)	14 (3%)	1 (0%)	49	57
1	B	493/571 (86%)	476 (97%)	16 (3%)	1 (0%)	49	57
1	C	470/571 (82%)	451 (96%)	16 (3%)	3 (1%)	27	28
1	D	493/571 (86%)	478 (97%)	14 (3%)	1 (0%)	49	57
1	E	493/571 (86%)	479 (97%)	13 (3%)	1 (0%)	49	57
1	F	487/571 (85%)	473 (97%)	13 (3%)	1 (0%)	49	57
2	X	4/6 (67%)	3 (75%)	1 (25%)	0	100	100
2	Y	4/6 (67%)	3 (75%)	1 (25%)	0	100	100
All	All	2938/3438 (86%)	2842 (97%)	88 (3%)	8 (0%)	43	48

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	330	VAL
1	C	440	PRO
1	B	330	VAL
1	C	332	GLY
1	C	516	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	429/485 (88%)	388 (90%)	41 (10%)	9	8
1	B	428/485 (88%)	390 (91%)	38 (9%)	11	10
1	C	414/485 (85%)	371 (90%)	43 (10%)	8	7
1	D	428/485 (88%)	385 (90%)	43 (10%)	8	7
1	E	428/485 (88%)	389 (91%)	39 (9%)	10	10
1	F	424/485 (87%)	382 (90%)	42 (10%)	8	8

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	X	4/4 (100%)	3 (75%)	1 (25%)	0	0
2	Y	4/4 (100%)	2 (50%)	2 (50%)	0	0
All	All	2559/2918 (88%)	2310 (90%)	249 (10%)	9	8

5 of 249 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	471	TYR
1	D	296	ASN
1	F	438	ARG
1	C	502	ARG
1	D	116	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 99 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	451	GLN
1	D	335	ASN
1	F	380	ASN
1	C	462	HIS
1	D	123	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 53 ligands modelled in this entry, 6 are monoatomic - leaving 47 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	A	1571	-	3,3,3	0.87	0	2,2,2	0.27	0
4	EDO	A	1572	-	3,3,3	0.51	0	2,2,2	0.56	0
4	EDO	A	1573	-	3,3,3	0.83	0	2,2,2	0.01	0
4	EDO	A	1574	-	3,3,3	0.54	0	2,2,2	0.13	0
4	EDO	A	1575	-	3,3,3	0.37	0	2,2,2	0.67	0
4	EDO	A	1576	-	3,3,3	0.31	0	2,2,2	0.77	0
4	EDO	A	1577	-	3,3,3	0.56	0	2,2,2	0.17	0
4	EDO	A	1578	-	3,3,3	0.71	0	2,2,2	0.66	0
4	EDO	A	1579	-	3,3,3	0.70	0	2,2,2	0.41	0
4	EDO	A	1580	-	3,3,3	0.97	0	2,2,2	1.19	0
5	BBK	A	1581	-	13,15,15	9.17	2 (15%)	15,21,21	2.16	5 (33%)
6	HWU	A	1582	3	31,41,41	3.84	5 (16%)	39,62,62	2.02	11 (28%)
6	HWU	B	1571	3	31,41,41	3.89	8 (25%)	39,62,62	2.34	13 (33%)
4	EDO	B	1572	-	3,3,3	0.53	0	2,2,2	0.64	0
4	EDO	B	1573	-	3,3,3	0.28	0	2,2,2	0.39	0
4	EDO	B	1574	-	3,3,3	0.62	0	2,2,2	0.33	0
4	EDO	B	1575	-	3,3,3	0.45	0	2,2,2	0.36	0
4	EDO	B	1576	-	3,3,3	0.65	0	2,2,2	0.13	0
4	EDO	B	1577	-	3,3,3	0.28	0	2,2,2	1.14	0
4	EDO	B	1578	-	3,3,3	0.55	0	2,2,2	0.39	0
4	EDO	B	1579	-	3,3,3	0.71	0	2,2,2	0.05	0
5	BBK	B	1580	-	13,15,15	9.03	2 (15%)	15,21,21	1.90	4 (26%)
6	HWU	C	1569	3	31,41,41	4.98	1 (3%)	39,62,62	2.62	16 (41%)
6	HWU	D	1571	3	31,41,41	4.42	5 (16%)	39,62,62	2.59	16 (41%)
5	BBK	D	1572	-	13,15,15	9.51	2 (15%)	15,21,21	3.39	5 (33%)
4	EDO	D	1573	-	3,3,3	0.51	0	2,2,2	0.77	0
4	EDO	D	1574	-	3,3,3	0.81	0	2,2,2	0.50	0
4	EDO	D	1575	-	3,3,3	0.91	0	2,2,2	0.41	0
4	EDO	D	1576	-	3,3,3	0.60	0	2,2,2	0.63	0
4	EDO	D	1577	-	3,3,3	0.46	0	2,2,2	0.70	0
4	EDO	E	1570	-	3,3,3	0.76	0	2,2,2	0.35	0
4	EDO	E	1571	-	3,3,3	0.69	0	2,2,2	0.29	0
4	EDO	E	1572	-	3,3,3	0.41	0	2,2,2	0.41	0
4	EDO	E	1573	-	3,3,3	0.31	0	2,2,2	1.00	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	E	1574	-	3,3,3	0.37	0	2,2,2	0.84	0
4	EDO	E	1575	-	3,3,3	0.47	0	2,2,2	0.35	0
4	EDO	E	1576	-	3,3,3	0.54	0	2,2,2	0.53	0
4	EDO	E	1577	-	3,3,3	0.36	0	2,2,2	0.77	0
4	EDO	E	1578	-	3,3,3	0.41	0	2,2,2	0.69	0
4	EDO	E	1579	-	3,3,3	0.47	0	2,2,2	0.69	0
4	EDO	E	1580	-	3,3,3	0.64	0	2,2,2	0.23	0
4	EDO	E	1581	-	3,3,3	0.36	0	2,2,2	0.88	0
4	EDO	E	1582	-	3,3,3	0.36	0	2,2,2	0.31	0
4	EDO	E	1583	-	3,3,3	0.59	0	2,2,2	0.15	0
5	BBK	E	1584	-	13,15,15	9.22	2 (15%)	15,21,21	2.14	5 (33%)
6	HWU	E	1585	3	31,41,41	3.94	3 (9%)	39,62,62	1.77	7 (17%)
6	HWU	F	1571	3	31,41,41	3.80	5 (16%)	39,62,62	1.66	6 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	1571	-	-	0/1/1/1	0/0/0/0
4	EDO	A	1572	-	-	0/1/1/1	0/0/0/0
4	EDO	A	1573	-	-	0/1/1/1	0/0/0/0
4	EDO	A	1574	-	-	0/1/1/1	0/0/0/0
4	EDO	A	1575	-	-	0/1/1/1	0/0/0/0
4	EDO	A	1576	-	-	0/1/1/1	0/0/0/0
4	EDO	A	1577	-	-	0/1/1/1	0/0/0/0
4	EDO	A	1578	-	-	0/1/1/1	0/0/0/0
4	EDO	A	1579	-	-	0/1/1/1	0/0/0/0
4	EDO	A	1580	-	-	0/1/1/1	0/0/0/0
5	BBK	A	1581	-	-	2/6/26/26	0/1/1/1
6	HWU	A	1582	3	-	0/21/63/63	0/3/3/3
6	HWU	B	1571	3	-	0/21/63/63	0/3/3/3
4	EDO	B	1572	-	-	0/1/1/1	0/0/0/0
4	EDO	B	1573	-	-	0/1/1/1	0/0/0/0
4	EDO	B	1574	-	-	0/1/1/1	0/0/0/0
4	EDO	B	1575	-	-	0/1/1/1	0/0/0/0
4	EDO	B	1576	-	-	0/1/1/1	0/0/0/0
4	EDO	B	1577	-	-	0/1/1/1	0/0/0/0
4	EDO	B	1578	-	-	0/1/1/1	0/0/0/0
4	EDO	B	1579	-	-	0/1/1/1	0/0/0/0
5	BBK	B	1580	-	-	0/6/26/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	HWU	C	1569	3	-	0/21/63/63	0/3/3/3
6	HWU	D	1571	3	-	0/21/63/63	0/3/3/3
5	BBK	D	1572	-	-	0/6/26/26	0/1/1/1
4	EDO	D	1573	-	-	0/1/1/1	0/0/0/0
4	EDO	D	1574	-	-	0/1/1/1	0/0/0/0
4	EDO	D	1575	-	-	0/1/1/1	0/0/0/0
4	EDO	D	1576	-	-	0/1/1/1	0/0/0/0
4	EDO	D	1577	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1570	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1571	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1572	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1573	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1574	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1575	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1576	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1577	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1578	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1579	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1580	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1581	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1582	-	-	0/1/1/1	0/0/0/0
4	EDO	E	1583	-	-	0/1/1/1	0/0/0/0
5	BBK	E	1584	-	-	0/6/26/26	0/1/1/1
6	HWU	E	1585	3	-	0/21/63/63	0/3/3/3
6	HWU	F	1571	3	-	0/21/63/63	0/3/3/3

The worst 5 of 35 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	C	1569	HWU	C5'-S5'	-27.24	1.41	1.82
5	D	1572	BBK	C5-S5	-27.03	1.42	1.82
5	E	1584	BBK	C5-S5	-26.64	1.42	1.82
5	B	1580	BBK	C5-S5	-25.96	1.43	1.82
5	A	1581	BBK	C5-S5	-25.79	1.43	1.82

The worst 5 of 88 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	1572	BBK	C1-C2-N2	-9.34	93.88	111.36
5	D	1572	BBK	O1-C1-C2	-7.78	93.37	109.25
6	C	1569	HWU	PB-O3A-PA	-4.47	117.62	132.63
6	D	1571	HWU	O1'-PB-O1B	-4.45	92.43	109.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	1569	HWU	O6'-C6'-C5'	-4.09	101.31	110.69

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1581	BBK	C8-C7-N2-C2
5	A	1581	BBK	O7-C7-N2-C2

There are no ring outliers.

25 monomers are involved in 72 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1572	EDO	4	0
4	A	1575	EDO	1	0
4	A	1578	EDO	1	0
4	A	1579	EDO	1	0
4	A	1580	EDO	2	0
5	A	1581	BBK	3	0
6	B	1571	HWU	2	0
4	B	1572	EDO	1	0
4	B	1573	EDO	4	0
4	B	1577	EDO	1	0
5	B	1580	BBK	4	0
6	C	1569	HWU	11	0
6	D	1571	HWU	4	0
5	D	1572	BBK	4	0
4	D	1573	EDO	1	0
4	D	1576	EDO	4	0
4	E	1570	EDO	1	0
4	E	1572	EDO	8	0
4	E	1573	EDO	1	0
4	E	1579	EDO	2	0
4	E	1580	EDO	1	0
4	E	1582	EDO	2	0
5	E	1584	BBK	5	0
6	E	1585	HWU	2	0
6	F	1571	HWU	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	C	3

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	222:PHE	C	223:LEU	N	1.63
1	C	225:SER	C	226:HIS	N	0.99
1	C	224:ASP	C	225:SER	N	0.87

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	495/571 (86%)	-0.03	16 (3%) 47 45	26, 42, 70, 130	5 (1%)
1	B	495/571 (86%)	-0.10	14 (2%) 53 50	26, 40, 66, 106	5 (1%)
1	C	476/571 (83%)	1.98	189 (39%) 0 0	42, 92, 135, 168	5 (1%)
1	D	495/571 (86%)	-0.04	20 (4%) 38 36	26, 41, 66, 142	5 (1%)
1	E	495/571 (86%)	-0.20	9 (1%) 68 66	23, 36, 58, 90	5 (1%)
1	F	491/571 (85%)	0.65	85 (17%) 1 1	34, 57, 112, 140	5 (1%)
2	X	6/6 (100%)	1.38	2 (33%) 0 0	40, 73, 84, 87	0
2	Y	6/6 (100%)	2.65	4 (66%) 0 0	42, 71, 77, 83	0
All	All	2959/3438 (86%)	0.37	339 (11%) 5 4	23, 45, 109, 168	30 (1%)

The worst 5 of 339 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	330	VAL	14.2
1	C	476	ALA	10.8
1	C	331	TRP	10.8
1	F	547	SER	10.4
1	F	556	GLY	8.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	BBK	B	1580	15/15	0.64	0.29	69,86,120,142	0
4	EDO	D	1575	4/4	0.64	0.22	60,63,68,70	0
5	BBK	D	1572	15/15	0.65	0.31	96,105,117,124	0
5	BBK	E	1584	15/15	0.70	0.32	96,105,117,124	0
4	EDO	B	1576	4/4	0.73	0.24	65,69,73,76	0
5	BBK	A	1581	15/15	0.77	0.23	69,86,120,142	0
4	EDO	E	1580	4/4	0.78	0.32	59,70,71,76	0
4	EDO	A	1580	4/4	0.80	0.22	45,52,57,59	0
4	EDO	E	1575	4/4	0.81	0.17	64,69,69,76	0
4	EDO	A	1572	4/4	0.81	0.24	57,66,67,69	0
4	EDO	E	1570	4/4	0.82	0.31	44,48,59,61	0
4	EDO	E	1574	4/4	0.82	0.19	62,68,68,75	0
4	EDO	A	1573	4/4	0.84	0.19	56,68,70,70	0
4	EDO	A	1577	4/4	0.85	0.48	62,68,71,76	0
4	EDO	A	1579	4/4	0.85	0.18	51,52,54,64	0
4	EDO	D	1577	4/4	0.85	0.20	67,68,69,69	0
6	HWU	C	1569	39/39	0.86	0.18	51,76,89,92	0
4	EDO	E	1576	4/4	0.87	0.19	68,68,69,69	0
4	EDO	A	1576	4/4	0.88	0.25	57,62,66,75	0
4	EDO	B	1573	4/4	0.88	0.28	40,41,42,48	0
4	EDO	E	1578	4/4	0.89	0.23	64,64,66,82	0
4	EDO	A	1574	4/4	0.89	0.15	51,52,54,57	0
4	EDO	B	1579	4/4	0.90	0.13	41,48,56,56	0
4	EDO	B	1578	4/4	0.90	0.28	55,58,62,69	0
4	EDO	E	1573	4/4	0.90	0.22	49,58,60,71	0
4	EDO	E	1572	4/4	0.91	0.25	38,45,48,51	0
4	EDO	D	1574	4/4	0.91	0.13	49,50,52,58	0
4	EDO	E	1583	4/4	0.91	0.15	58,61,62,68	0
6	HWU	F	1571	39/39	0.91	0.12	38,44,55,60	0
6	HWU	A	1582	39/39	0.92	0.12	26,34,45,48	0
4	EDO	D	1573	4/4	0.92	0.13	38,40,40,45	0
4	EDO	A	1578	4/4	0.92	0.19	38,45,48,63	0
4	EDO	E	1581	4/4	0.92	0.16	53,56,60,68	0
6	HWU	B	1571	39/39	0.92	0.13	30,36,44,49	0
6	HWU	D	1571	39/39	0.93	0.12	24,35,45,46	0
4	EDO	B	1575	4/4	0.93	0.25	66,68,69,69	0
4	EDO	B	1574	4/4	0.94	0.10	40,40,42,45	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	EDO	D	1576	4/4	0.94	0.15	39,43,45,49	0
4	EDO	B	1577	4/4	0.94	0.20	42,43,48,57	0
4	EDO	E	1577	4/4	0.94	0.21	61,61,64,68	0
4	EDO	A	1571	4/4	0.94	0.20	43,46,46,52	0
4	EDO	A	1575	4/4	0.95	0.11	53,55,57,59	0
3	MN	C	1570	1/1	0.95	0.04	59,59,59,59	0
6	HWU	E	1585	39/39	0.95	0.12	26,36,54,58	0
4	EDO	B	1572	4/4	0.95	0.10	34,35,35,42	0
4	EDO	E	1571	4/4	0.96	0.16	38,46,47,50	0
4	EDO	E	1579	4/4	0.96	0.11	43,44,52,53	0
4	EDO	E	1582	4/4	0.96	0.12	47,48,51,56	0
3	MN	E	1586	1/1	0.99	0.08	26,26,26,26	0
3	MN	F	1570	1/1	1.00	0.03	42,42,42,42	0
3	MN	B	1570	1/1	1.00	0.09	33,33,33,33	0
3	MN	A	1570	1/1	1.00	0.08	27,27,27,27	0
3	MN	D	1570	1/1	1.00	0.08	29,29,29,29	0

6.5 Other polymers

There are no such residues in this entry.