



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 02:30 am GMT

PDB ID : 2D24  
Title : Crystal structure of ES complex of catalytic-site mutant xylanase from *Streptomyces olivaceoviridis* E-86  
Authors : Suzuki, R.; Kuno, A.; Fujimoto, Z.; Ito, S.; Kawahara, S.I.; Kaneko, S.; Hasegawa, T.; Taira, K.  
Deposited on : 2005-09-02  
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2111 (1.86-1.86)
Clashscore	122126	2258 (1.86-1.86)
Ramachandran outliers	120053	2234 (1.86-1.86)
Sidechain outliers	120020	2234 (1.86-1.86)
RSRZ outliers	108989	2075 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	436	<div> <div>3%</div> <div> <div></div> <div>84%</div> <div>13%</div> <div>••</div> </div> </div>
1	B	436	<div> <div>7%</div> <div> <div></div> <div>82%</div> <div>15%</div> <div>•</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	XYS	A	451	X	-	-	-
2	XYS	A	452	X	-	-	-
2	XYS	A	453	X	-	-	-
2	XYS	A	454	X	-	-	-
2	XYS	A	455	X	-	-	-
2	XYS	A	456	X	-	-	X
2	XYS	A	457	X	-	-	-
2	XYS	A	458	X	-	-	-
2	XYS	A	459	X	-	-	-
2	XYS	A	956	X	-	-	X
2	XYS	A	957	X	-	-	-
2	XYS	B	951	X	-	-	-
2	XYS	B	952	X	-	-	-
2	XYS	B	953	X	-	-	-
2	XYS	B	954	X	-	-	-
2	XYS	B	955	X	-	-	-
2	XYS	B	958	X	-	-	-
2	XYS	B	959	X	-	-	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7333 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ENDO-1,4-BETA-D-XYLANASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	427	Total	C	N	O	S	0	0	0
			3233	1988	588	641	16			
1	B	427	Total	C	N	O	S	0	0	0
			3233	1988	588	641	16			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	SER	ASN	ENGINEERED MUTATION	UNP Q7SI98
A	128	HIS	GLU	ENGINEERED MUTATION	UNP Q7SI98
B	627	SER	ASN	ENGINEERED MUTATION	UNP Q7SI98
B	628	HIS	GLU	ENGINEERED MUTATION	UNP Q7SI98

- Molecule 2 is XYLOPYRANOSE (three-letter code: YYS) (formula: C<sub>5</sub>H<sub>10</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			9	5	4		
2	A	1	Total	C	O	0	0
			9	5	4		
2	A	1	Total	C	O	0	0
			9	5	4		
2	A	1	Total	C	O	0	0
			9	5	4		
2	A	1	Total	C	O	0	0
			10	5	5		
2	A	1	Total	C	O	0	0
			9	5	4		
2	A	1	Total	C	O	0	0
			9	5	4		
2	A	1	Total	C	O	0	0
			9	5	4		
2	A	1	Total	C	O	0	0
			10	5	5		
2	A	1	Total	C	O	0	0
			9	5	4		
2	B	1	Total	C	O	0	0
			9	5	4		
2	B	1	Total	C	O	0	0
			9	5	4		
2	B	1	Total	C	O	0	0
			9	5	4		
2	B	1	Total	C	O	0	0
			10	5	5		
2	B	1	Total	C	O	0	0
			9	5	4		
2	B	1	Total	C	O	0	0
			10	5	5		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	318	Total	O	0	0
			318	318		
5	B	354	Total	O	0	0
			354	354		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ENDO-1,4-BETA-D-XYLANASE



#### • Molecule 1: ENDO-1,4-BETA-D-XYLANASE





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.70Å 94.09Å 138.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.51 – 1.85 36.51 – 1.85	Depositor EDS
% Data completeness (in resolution range)	(Not available) (36.51-1.85) 99.9 (36.51-1.85)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.21 (at 1.85Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.177 , 0.199 0.177 , 0.198	Depositor DCC
$R_{free}$ test set	4311 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.1	Xtriage
Anisotropy	0.033	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 53.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7333	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.86% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, XYS, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.29	0/3298	0.61	0/4473
1	B	0.30	0/3298	0.61	0/4473
All	All	0.30	0/6596	0.61	0/8946

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3233	0	3043	43	0
1	B	3233	0	3040	50	0
2	A	102	0	86	7	0
2	B	65	0	55	2	0
3	A	10	0	0	0	0
4	B	18	0	24	0	0
5	A	318	0	0	3	0
5	B	354	0	0	3	0
All	All	7333	0	6248	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 95 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:935:ARG:HA	1:B:936:THR:HB	1.45	0.96
1:B:836:GLN:HB2	1:B:876:LYS:HE3	1.59	0.84
1:A:1:ALA:HB1	5:A:1194:HOH:O	1.85	0.77
1:B:926:SER:O	1:B:927:ASN:HB2	1.87	0.73
1:A:340:TYR:CZ	2:A:957:XYS:H52	2.24	0.71

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	423/436 (97%)	410 (97%)	13 (3%)	0	100	100
1	B	423/436 (97%)	406 (96%)	15 (4%)	2 (0%)	31	16
All	All	846/872 (97%)	816 (96%)	28 (3%)	2 (0%)	49	34

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	927	ASN
1	B	844	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	334/341 (98%)	329 (98%)	5 (2%)	67	56
1	B	334/341 (98%)	329 (98%)	5 (2%)	67	56
All	All	668/682 (98%)	658 (98%)	10 (2%)	67	56

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	435	ARG
1	B	585	TRP
1	B	747	TYR
1	A	280	PRO
1	B	601	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	349	GLN
1	B	511	GLN
1	B	828	ASN
1	A	343	HIS
1	B	705	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

23 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	XYS	A	451	2	9,9,10	0.69	0	10,12,14	0.75	0
2	XYS	A	452	2	9,9,10	1.80	2 (22%)	10,12,14	1.78	2 (20%)
2	XYS	A	453	2	9,9,10	0.98	0	10,12,14	1.08	1 (10%)
2	XYS	A	454	2	9,9,10	0.78	0	10,12,14	0.97	0
2	XYS	A	455	2	10,10,10	0.58	0	14,14,14	0.60	0
2	XYS	A	456	2	9,9,10	0.67	0	10,12,14	0.83	1 (10%)
2	XYS	A	457	2	9,9,10	0.72	0	10,12,14	0.78	0
2	XYS	A	458	2	9,9,10	0.64	0	10,12,14	0.78	1 (10%)
2	XYS	A	459	2	10,10,10	0.61	0	14,14,14	0.60	0
2	XYS	A	956	2	9,9,10	0.70	0	10,12,14	0.83	1 (10%)
2	XYS	A	957	2	10,10,10	0.59	0	14,14,14	0.60	0
3	SO4	A	970	-	4,4,4	0.35	0	6,6,6	0.07	0
3	SO4	A	971	-	4,4,4	0.35	0	6,6,6	0.09	0
2	XYS	B	951	2	9,9,10	0.62	0	10,12,14	0.78	0
2	XYS	B	952	2	9,9,10	1.83	3 (33%)	10,12,14	1.81	3 (30%)
2	XYS	B	953	2	9,9,10	1.01	1 (11%)	10,12,14	1.13	1 (10%)
2	XYS	B	954	2	9,9,10	0.82	0	10,12,14	1.06	1 (10%)
2	XYS	B	955	2	10,10,10	0.53	0	14,14,14	0.65	0
2	XYS	B	958	2	9,9,10	0.64	0	10,12,14	0.82	1 (10%)
2	XYS	B	959	2	10,10,10	0.55	0	14,14,14	0.58	0
4	GOL	B	980	-	5,5,5	0.15	0	5,5,5	0.31	0
4	GOL	B	981	-	5,5,5	0.13	0	5,5,5	0.31	0
4	GOL	B	982	-	5,5,5	0.16	0	5,5,5	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XYS	A	451	2	1/1/3/4	0/0/14/17	0/1/1/1
2	XYS	A	452	2	1/1/3/4	0/0/14/17	1/1/1/1
2	XYS	A	453	2	1/1/3/4	0/0/14/17	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XYS	A	454	2	1/1/3/4	0/0/14/17	0/1/1/1
2	XYS	A	455	2	1/1/4/4	0/0/17/17	0/1/1/1
2	XYS	A	456	2	1/1/3/4	0/0/14/17	0/1/1/1
2	XYS	A	457	2	1/1/3/4	0/0/14/17	0/1/1/1
2	XYS	A	458	2	1/1/3/4	0/0/14/17	0/1/1/1
2	XYS	A	459	2	1/1/4/4	0/0/17/17	0/1/1/1
2	XYS	A	956	2	1/1/3/4	0/0/14/17	0/1/1/1
2	XYS	A	957	2	1/1/4/4	0/0/17/17	0/1/1/1
3	SO4	A	970	-	-	0/0/0/0	0/0/0/0
3	SO4	A	971	-	-	0/0/0/0	0/0/0/0
2	XYS	B	951	2	1/1/3/4	0/0/14/17	0/1/1/1
2	XYS	B	952	2	1/1/3/4	0/0/14/17	1/1/1/1
2	XYS	B	953	2	1/1/3/4	0/0/14/17	0/1/1/1
2	XYS	B	954	2	1/1/3/4	0/0/14/17	0/1/1/1
2	XYS	B	955	2	1/1/4/4	0/0/17/17	0/1/1/1
2	XYS	B	958	2	1/1/3/4	0/0/14/17	0/1/1/1
2	XYS	B	959	2	1/1/4/4	0/0/17/17	0/1/1/1
4	GOL	B	980	-	-	0/4/4/4	0/0/0/0
4	GOL	B	981	-	-	0/4/4/4	0/0/0/0
4	GOL	B	982	-	-	0/4/4/4	0/0/0/0

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	953	XYS	O5-C5	2.01	1.46	1.42
2	B	952	XYS	C5-C4	2.06	1.57	1.52
2	A	452	XYS	O5-C5	2.48	1.47	1.42
2	B	952	XYS	O5-C5	2.48	1.47	1.42
2	B	952	XYS	C2-C3	3.36	1.57	1.52

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	452	XYS	C5-C4-C3	-3.74	104.93	109.66
2	B	952	XYS	C5-C4-C3	-3.66	105.03	109.66
2	B	952	XYS	C4-C3-C2	-3.46	106.84	110.86
2	A	452	XYS	C4-C3-C2	-3.37	106.94	110.86
2	B	958	XYS	C4-C3-C2	-2.26	108.24	110.86

5 of 18 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	452	XYS	C1
2	B	953	XYS	C1
2	B	958	XYS	C1
2	B	951	XYS	C1
2	A	957	XYS	C1

There are no torsion outliers.

All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	952	XYS	C1-C2-C3-C4-C5-O5
2	A	452	XYS	C1-C2-C3-C4-C5-O5

8 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	453	XYS	1	0
2	A	456	XYS	1	0
2	A	457	XYS	1	0
2	A	458	XYS	1	0
2	A	956	XYS	1	0
2	A	957	XYS	4	0
2	B	953	XYS	1	0
2	B	959	XYS	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	427/436 (97%)	0.02	13 (3%) 50 48	8, 14, 27, 46	0
1	B	427/436 (97%)	0.11	30 (7%) 16 16	6, 12, 38, 54	0
All	All	854/872 (97%)	0.07	43 (5%) 29 27	6, 13, 32, 54	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	936	THR	7.3
1	A	303	GLY	7.2
1	B	818	VAL	7.1
1	B	501	ALA	5.8
1	A	1	ALA	5.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	XYS	A	456	9/10	0.51	0.60	50,54,55,55	0
2	XYS	A	956	9/10	0.52	0.53	40,42,44,45	0
4	GOL	B	981	6/6	0.61	0.16	42,43,43,44	0
2	XYS	A	457	9/10	0.65	0.24	31,34,39,45	0
4	GOL	B	982	6/6	0.68	0.22	41,43,43,43	0
4	GOL	B	980	6/6	0.69	0.19	34,35,35,35	0
2	XYS	B	955	10/10	0.71	0.35	22,36,41,43	0
2	XYS	A	455	10/10	0.73	0.38	29,40,43,44	0
3	SO4	A	970	5/5	0.73	0.34	77,77,78,78	0
2	XYS	B	959	10/10	0.75	0.22	42,44,46,47	0
2	XYS	B	958	9/10	0.78	0.26	48,49,50,50	0
2	XYS	A	459	10/10	0.83	0.16	26,29,31,35	0
2	XYS	A	957	10/10	0.85	0.18	21,28,31,33	0
2	XYS	A	453	9/10	0.90	0.14	16,19,20,21	0
3	SO4	A	971	5/5	0.90	0.23	56,56,56,56	0
2	XYS	B	953	9/10	0.92	0.14	13,13,15,16	0
2	XYS	A	454	9/10	0.93	0.14	18,20,22,24	0
2	XYS	A	452	9/10	0.94	0.14	12,13,16,16	0
2	XYS	B	952	9/10	0.95	0.10	7,11,13,13	0
2	XYS	A	458	9/10	0.95	0.11	17,21,23,26	0
2	XYS	A	451	9/10	0.96	0.11	12,13,14,14	0
2	XYS	B	954	9/10	0.96	0.10	12,13,16,18	0
2	XYS	B	951	9/10	0.97	0.09	8,9,10,12	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.