



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2018 – 07:58 am GMT

PDB ID : 1D3U
Title : TATA-BINDING PROTEIN / TRANSCRIPTION FACTOR
(II)B/BRE+TATA-BOX COMPLEX FROM PYROCOCCUS WOESEI
Authors : Littlefield, O.; Korkhin, Y.; Sigler, P.B.
Deposited on : 1999-10-01
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	3481 (2.40-2.40)
Clashscore	122126	3956 (2.40-2.40)
Ramachandran outliers	120053	3897 (2.40-2.40)
Sidechain outliers	120020	3898 (2.40-2.40)
RSRZ outliers	108989	3386 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	24	<div> <div>4%</div> <div>42%</div> <div>46%</div> <div>13%</div> </div>
2	D	23	<div> <div>9%</div> <div>22%</div> <div>61%</div> <div>17%</div> </div>
3	A	181	<div> <div>10%</div> <div>71%</div> <div>27%</div> <div>..</div> </div>
4	B	201	<div> <div>4%</div> <div>79%</div> <div>18%</div> <div>.</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4147 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA 24-MER: BRE+TATA-BOX.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	24	Total	C	N	O	P	0	0	0
			493	239	91	140	23			

- Molecule 2 is a DNA chain called DNA 23-MER: BRE+TATA-BOX.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	23	Total	C	N	O	P	0	0	0
			464	227	76	139	22			

- Molecule 3 is a protein called TATA-BINDING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	181	Total	C	N	O	S	0	0	0
			1415	910	235	263	7			

- Molecule 4 is a protein called TRANSCRIPTION INITIATION FACTOR IIB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	201	Total	C	N	O	S	0	0	0
			1595	1006	296	289	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1100	MET	ARG	ENGINEERED	UNP P61999

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	16	Total	O	0	0
			16	16		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	19	Total	O	0	0
			19	19		
5	A	72	Total	O	0	0
			72	72		
5	B	73	Total	O	0	0
			73	73		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA 24-MER: BRE+TATA-BOX





- Molecule 2: DNA 23-MER: BRE+TATA-BOX



- Molecule 3: TATA-BINDING PROTEIN



- Molecule 4: TRANSCRIPTION INITIATION FACTOR IIB



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	174.20Å 174.20Å 174.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.40 28.26 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.7 (20.00-2.40) 98.8 (28.26-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.19 (at 2.42Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.206 , 0.245 0.205 , 0.243	Depositor DCC
R_{free} test set	3325 reflections (9.83%)	wwPDB-VP
Wilson B-factor (Å ²)	50.9	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 59.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.021 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4147	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.66	0/554	1.01	0/854
2	D	0.69	0/518	0.99	1/797 (0.1%)
3	A	0.64	0/1438	0.77	1/1939 (0.1%)
4	B	0.68	0/1612	0.77	0/2165
All	All	0.66	0/4122	0.84	2/5755 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	4
2	D	0	3
All	All	0	7

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	52	ASP	C-N-CD	5.49	139.94	128.40
2	D	1440	DC	C1'-O4'-C4'	-5.13	104.97	110.10

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	1409	DG	Sidechain
1	C	1414	DA	Sidechain
1	C	1415	DA	Sidechain
1	C	1418	DC	Sidechain
2	D	1431	DG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	493	0	275	14	0
2	D	464	0	266	25	1
3	A	1415	0	1473	41	1
4	B	1595	0	1695	38	0
5	A	72	0	0	0	0
5	B	73	0	0	6	0
5	C	16	0	0	0	0
5	D	19	0	0	1	0
All	All	4147	0	3709	109	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 109 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:1149:ARG:HH11	4:B:1149:ARG:HB3	0.99	1.11
1:C:1415:DA:H2''	1:C:1416:DT:H5'	1.31	1.11
3:A:15:VAL:HG22	3:A:68:THR:HG22	1.45	0.98
4:B:1149:ARG:HB3	4:B:1149:ARG:NH1	1.82	0.93
4:B:1149:ARG:HH11	4:B:1149:ARG:CB	1.83	0.91

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:1426:DT:O4'	2:D:1426:DT:O4'[16_546]	2.09	0.11
3:A:172:ARG:NH1	3:A:172:ARG:NH1[15_556]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	179/181 (99%)	163 (91%)	10 (6%)	6 (3%)	4	3
4	B	199/201 (99%)	193 (97%)	3 (2%)	3 (2%)	11	15
All	All	378/382 (99%)	356 (94%)	13 (3%)	9 (2%)	6	7

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	52	ASP
4	B	1265	LYS
4	B	1298	PRO
3	A	30	LEU
3	A	34	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	159/159 (100%)	151 (95%)	8 (5%)	27	43
4	B	167/167 (100%)	162 (97%)	5 (3%)	44	65
All	All	326/326 (100%)	313 (96%)	13 (4%)	34	53

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	127	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	A	142	LYS
4	B	1208	PRO
3	A	74	GLN
4	B	1195	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
4	B	1122	GLN
4	B	1213	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	24/24 (100%)	0.01	1 (4%) 36 34	34, 59, 87, 100	0
2	D	23/23 (100%)	0.05	2 (8%) 10 9	34, 59, 93, 100	0
3	A	181/181 (100%)	0.30	18 (9%) 7 6	27, 53, 97, 100	0
4	B	201/201 (100%)	0.04	9 (4%) 33 31	25, 44, 77, 100	0
All	All	429/429 (100%)	0.15	30 (6%) 16 14	25, 49, 95, 100	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	B	1100	MET	14.8
4	B	1300	ALA	9.6
3	A	1	MET	8.8
4	B	1299	ILE	6.4
3	A	2	VAL	5.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.