



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 19, 2019 – 01:27 PM EDT

PDB ID : 3D4O  
Title : Crystal structure of dipicolinate synthase subunit A (NP\_243269.1) from BACILLUS HALODURANS at 2.10 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2008-05-14  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : 2.3.2  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.3.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4608 (2.10-2.10)
Clashscore	122126	5109 (2.10-2.10)
Ramachandran outliers	120053	5059 (2.10-2.10)
Sidechain outliers	120020	5060 (2.10-2.10)
RSRZ outliers	108989	4497 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	<div> <div>8%</div> <div>86%13% .</div> </div>
1	B	293	<div> <div>7%</div> <div>86%13% .</div> </div>
1	C	293	<div> <div>8%</div> <div>87%12% .</div> </div>
1	D	293	<div> <div>6%</div> <div>88%10% .</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TAR	A	295	X	-	-	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9415 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dipicolinate synthase subunit A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	290	Total	C	N	O	S	Se	0	4	0
			2208	1407	378	409	3	11			
1	B	291	Total	C	N	O	S	Se	0	3	0
			2188	1393	374	406	3	12			
1	C	291	Total	C	N	O	S	Se	0	2	0
			2153	1371	362	406	3	11			
1	D	290	Total	C	N	O	S	Se	0	2	0
			2159	1375	367	402	3	12			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	leader sequence	UNP Q9KA87
B	0	GLY	-	leader sequence	UNP Q9KA87
C	0	GLY	-	leader sequence	UNP Q9KA87
D	0	GLY	-	leader sequence	UNP Q9KA87

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is D(-)-TARTARIC ACID (three-letter code: TAR) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	4	6		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	182	Total	O	0	4
			186	186		
4	B	154	Total	O	0	4
			158	158		
4	C	132	Total	O	0	4
			136	136		
4	D	164	Total	O	0	1
			165	165		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dipicolinate synthase subunit A




- Molecule 1: Dipicolinate synthase subunit A




- Molecule 1: Dipicolinate synthase subunit A



- Molecule 1: Dipicolinate synthase subunit A







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.65Å 142.42Å 74.86Å 90.00° 93.49° 90.00°	Depositor
Resolution (Å)	29.77 – 2.10 29.77 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.2 (29.77-2.10) 99.3 (29.77-2.10)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.50 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.4.0067, PHENIX	Depositor
R, $R_{free}$	0.189 , 0.240 0.194 , 0.241	Depositor DCC
$R_{free}$ test set	3729 reflections (5.18%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.2	Xtriage
Anisotropy	0.314	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 59.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9415	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 49.31 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.6001e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, TAR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.74	0/2239	0.85	3/3013 (0.1%)
1	B	0.75	0/2215	0.88	2/2981 (0.1%)
1	C	0.64	0/2178	0.84	1/2940 (0.0%)
1	D	0.66	0/2182	0.83	2/2939 (0.1%)
All	All	0.70	0/8814	0.85	8/11873 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	143	MSE	CG-SE-CE	-8.73	79.69	98.90
1	C	143	MSE	CG-SE-CE	-8.52	80.15	98.90
1	B	171[A]	ARG	NE-CZ-NH1	-8.34	116.13	120.30
1	B	171[B]	ARG	NE-CZ-NH1	-8.34	116.13	120.30
1	A	171	ARG	NE-CZ-NH1	-6.46	117.07	120.30
1	A	149	ASP	CB-CG-OD2	-5.74	113.14	118.30
1	D	125	ARG	NE-CZ-NH2	-5.22	117.69	120.30
1	A	149	ASP	CB-CG-OD1	5.13	122.92	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2208	0	2309	26	0
1	B	2188	0	2264	31	0
1	C	2153	0	2191	27	0
1	D	2159	0	2216	27	0
2	A	8	0	12	0	0
2	B	16	0	24	0	0
2	C	12	0	18	0	0
2	D	16	0	24	1	0
3	A	10	0	4	0	0
4	A	186	0	0	0	0
4	B	158	0	0	2	0
4	C	136	0	0	1	0
4	D	165	0	0	1	0
All	All	9415	0	9062	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:69:ASN:ND2	1:B:73:LYS:HB3	1.40	1.32
1:B:69:ASN:HD21	1:B:73:LYS:CB	1.54	1.20
1:B:128:ILE:HD11	1:B:283:VAL:HG21	1.44	1.00
1:C:171[A]:ARG:HH21	1:D:171[A]:ARG:HH12	1.12	0.97
1:A:257[B]:ARG:HD3	1:A:259[B]:ILE:HG13	1.52	0.89
1:B:69:ASN:ND2	1:B:73:LYS:CB	2.18	0.87
1:D:90:ILE:HD13	1:D:112:MSE:HE1	1.55	0.86
1:D:141:ILE:HD11	1:D:172:LYS:HB3	1.58	0.85
1:A:257[B]:ARG:CD	1:A:259[B]:ILE:HG13	2.05	0.85
1:B:69:ASN:HD21	1:B:73:LYS:HB2	1.40	0.84
1:B:65:ILE:HG23	1:B:123:MSE:HE2	1.59	0.84
1:C:90:ILE:HD13	1:C:112:MSE:HE1	1.60	0.82
1:B:142[B]:MSE:HG2	1:D:142[B]:MSE:HG2	1.66	0.77
1:B:90:ILE:HD13	1:B:112:MSE:CE	2.16	0.76
1:D:233:MSE:HE2	1:D:259:ILE:HD13	1.71	0.72
1:C:38:LEU:O	1:C:47:LYS:NZ	2.18	0.70
1:C:90:ILE:HD13	1:C:112:MSE:CE	2.20	0.70
1:B:65:ILE:CG2	1:B:123:MSE:HE2	2.21	0.70
1:C:171[A]:ARG:NH2	1:D:171[A]:ARG:HH12	1.88	0.69
1:C:141:ILE:HD11	1:C:172:LYS:HB3	1.74	0.69
1:A:141:ILE:HD11	1:A:172:LYS:HB3	1.75	0.68

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:90:ILE:HD13	1:B:112:MSE:HE1	1.74	0.68
1:B:128:ILE:HD11	1:B:283:VAL:CG2	2.21	0.68
1:A:26:PHE:CD2	1:A:285[A]:VAL:HG21	2.28	0.67
1:D:1:MSE:HE2	1:D:285:VAL:HG12	1.78	0.65
1:B:128:ILE:CD1	1:B:283:VAL:HG21	2.24	0.64
1:C:171[A]:ARG:HH21	1:D:171[A]:ARG:NH1	1.89	0.64
1:D:139:GLY:O	1:D:143:MSE:HG3	1.96	0.64
1:C:233:MSE:HE3	1:C:237:THR:HB	1.79	0.62
1:D:123:MSE:HE1	1:D:128:ILE:HG22	1.80	0.62
1:D:141:ILE:HG23	1:D:176:LEU:HD11	1.82	0.62
1:D:112:MSE:HG3	1:D:119:LEU:HB2	1.81	0.62
1:A:68:THR:HG22	1:A:74:VAL:HG22	1.82	0.61
1:B:225:VAL:HA	1:B:229[A]:VAL:HG21	1.84	0.60
1:A:175:ALA:HB1	1:C:175:ALA:HB1	1.84	0.59
1:D:1:MSE:HE3	1:D:289:ALA:CB	2.32	0.59
1:D:141:ILE:HD11	1:D:172:LYS:CB	2.32	0.59
1:B:69:ASN:HD21	1:B:73:LYS:HB3	1.08	0.58
1:D:141:ILE:CD1	1:D:172:LYS:HB3	2.33	0.58
1:C:162:LEU:HD21	1:C:167:MSE:CE	2.35	0.57
1:A:126:ASP:HA	1:A:164:ARG:HH11	1.70	0.56
1:C:55:TRP:HB2	1:C:92:LYS:HB3	1.88	0.56
1:A:108:LEU:O	1:A:112:MSE:HG2	2.06	0.56
1:A:90:ILE:HD13	1:A:112:MSE:CE	2.36	0.56
1:B:126:ASP:HA	1:B:164:ARG:HH11	1.72	0.55
1:C:90:ILE:HG21	1:C:112:MSE:HE1	1.88	0.54
1:B:141:ILE:HD11	1:B:172:LYS:HB3	1.89	0.54
1:C:233:MSE:HE3	1:C:237:THR:CB	2.38	0.54
1:D:42:PHE:HB2	1:D:45:VAL:CG2	2.38	0.54
1:A:128:ILE:HD11	1:A:283:VAL:HG21	1.89	0.53
1:B:112:MSE:HG3	1:B:119:LEU:HB2	1.91	0.53
1:A:90:ILE:HD13	1:A:112:MSE:HE1	1.91	0.52
1:D:233:MSE:HE3	1:D:237:THR:HB	1.90	0.52
1:D:22:LYS:NZ	1:D:282:ASP:OD1	2.30	0.52
1:A:245:LYS:HE2	1:A:265:PRO:HG3	1.90	0.52
1:A:210:LEU:HD13	1:A:233:MSE:HE3	1.90	0.52
1:A:257[B]:ARG:HD2	1:A:259[B]:ILE:CD1	2.39	0.52
1:B:226:THR:O	1:B:229[A]:VAL:HG22	2.10	0.52
1:D:1:MSE:O	1:D:26:PHE:O	2.28	0.51
1:C:123:MSE:HE1	4:C:316:HOH:O	2.10	0.51
1:C:22:LYS:NZ	1:C:282:ASP:OD1	2.36	0.51
1:B:167:MSE:HE2	1:B:196:MSE:SE	2.60	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:26:PHE:CE2	1:A:285[A]:VAL:HG21	2.47	0.49
1:A:257[B]:ARG:HD2	1:A:259[B]:ILE:HG13	1.93	0.49
1:A:196:MSE:HB2	1:A:198:MSE:HE2	1.95	0.48
1:D:90:ILE:HD13	1:D:112:MSE:CE	2.34	0.48
1:D:1:MSE:HE3	1:D:289:ALA:HB2	1.96	0.48
1:C:186:GLU:OE1	1:C:186:GLU:N	2.47	0.47
1:C:68:THR:HG22	1:C:74:VAL:HG22	1.96	0.47
1:C:190:LEU:HA	1:C:193:ILE:HD12	1.98	0.46
1:C:1:MSE:HG3	1:C:26:PHE:CD2	2.50	0.45
1:B:253:TYR:HA	1:B:256:LYS:HG2	1.99	0.45
1:B:162:LEU:HD21	1:B:167:MSE:CE	2.47	0.45
2:D:295:EDO:H22	4:D:452:HOH:O	2.17	0.44
4:B:394[B]:HOH:O	1:D:171[B]:ARG:NH2	1.99	0.44
1:A:267:LEU:HD23	1:A:267:LEU:HA	1.91	0.44
1:C:167:MSE:HB2	1:C:167:MSE:HE3	1.73	0.44
1:A:203:ILE:HA	1:A:203:ILE:HD12	1.83	0.44
1:C:158:ALA:HA	1:C:181:LYS:O	2.18	0.44
1:A:2:LEU:O	1:A:5:LYS:HB2	2.17	0.44
1:B:142[B]:MSE:HE3	1:B:142[B]:MSE:HB3	1.68	0.44
1:C:223:LEU:HG	1:C:226:THR:HG22	1.99	0.43
1:A:26:PHE:CD2	1:A:285[A]:VAL:CG2	2.97	0.43
1:D:1:MSE:O	1:D:2:LEU:HB2	2.18	0.43
1:B:8:VAL:HG22	1:B:31:SER:HB2	2.00	0.43
1:C:1:MSE:HB2	1:C:26:PHE:HD2	1.84	0.43
1:A:33:VAL:HG12	1:A:83:ILE:HD13	2.00	0.43
1:B:123:MSE:HE1	4:B:400:HOH:O	2.18	0.42
1:B:55:TRP:O	1:B:93:THR:HA	2.18	0.42
1:C:42:PHE:HB2	1:C:45:VAL:CG2	2.49	0.42
1:B:175:ALA:HB1	1:D:175:ALA:HB1	2.02	0.42
1:A:65:ILE:HD13	1:A:65:ILE:HA	1.86	0.42
1:B:229[B]:VAL:HG12	1:B:233:MSE:HE3	2.02	0.41
1:B:34:GLY:O	1:B:76:THR:OG1	2.32	0.41
1:D:68:THR:HG22	1:D:74:VAL:HG22	2.02	0.41
1:A:158:ALA:HA	1:A:181:LYS:O	2.20	0.41
1:C:8:VAL:CG1	1:C:61:ILE:HG12	2.51	0.41
1:B:32:LEU:HG	1:B:35:PHE:CD1	2.56	0.41
1:D:267:LEU:HD23	1:D:267:LEU:HA	1.94	0.41
1:B:108:LEU:O	1:B:112:MSE:HG2	2.21	0.41
1:D:65:ILE:HD13	1:D:65:ILE:HA	1.97	0.40
1:B:143:MSE:SE	1:B:264:VAL:HG11	2.69	0.40
1:A:102:GLY:CA	1:A:123:MSE:HG2	2.51	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:VAL:HG13	1:A:62:LEU:HD13	2.02	0.40
1:C:139:GLY:O	1:C:143:MSE:HG3	2.22	0.40
1:C:112:MSE:HG3	1:C:119:LEU:HB2	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	290/293 (99%)	284 (98%)	6 (2%)	0	100	100
1	B	290/293 (99%)	284 (98%)	6 (2%)	0	100	100
1	C	291/293 (99%)	285 (98%)	6 (2%)	0	100	100
1	D	288/293 (98%)	277 (96%)	11 (4%)	0	100	100
All	All	1159/1172 (99%)	1130 (98%)	29 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/230 (105%)	239 (99%)	2 (1%)	83	88
1	B	235/230 (102%)	233 (99%)	2 (1%)	81	85

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	227/230 (99%)	226 (100%)	1 (0%)	92	95
1	D	229/230 (100%)	228 (100%)	1 (0%)	92	95
All	All	932/920 (101%)	926 (99%)	6 (1%)	85	91

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	ASP
1	A	65	ILE
1	B	80	ASN
1	B	257	ARG
1	C	73	LYS
1	D	257	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	ASN
1	A	109	ASN
1	B	208	GLN
1	C	6	HIS
1	C	37	GLN
1	C	109	ASN
1	C	153	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EDO	A	293	-	3,3,3	0.33	0	2,2,2	0.07	0
2	EDO	A	294	-	3,3,3	0.52	0	2,2,2	0.23	0
3	TAR	A	295	-	3,9,9	0.86	0	6,12,12	0.80	0
2	EDO	B	293	-	3,3,3	0.46	0	2,2,2	0.45	0
2	EDO	B	294	-	3,3,3	0.41	0	2,2,2	0.48	0
2	EDO	B	295	-	3,3,3	0.35	0	2,2,2	0.59	0
2	EDO	B	296	-	3,3,3	0.43	0	2,2,2	0.38	0
2	EDO	C	293	-	3,3,3	0.49	0	2,2,2	0.44	0
2	EDO	C	294	-	3,3,3	0.58	0	2,2,2	0.29	0
2	EDO	C	295	-	3,3,3	0.49	0	2,2,2	0.33	0
2	EDO	D	293	-	3,3,3	0.43	0	2,2,2	0.37	0
2	EDO	D	294	-	3,3,3	0.48	0	2,2,2	0.76	0
2	EDO	D	295	-	3,3,3	0.75	0	2,2,2	0.18	0
2	EDO	D	296	-	3,3,3	0.41	0	2,2,2	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	293	-	-	0/1/1/1	-
2	EDO	A	294	-	-	1/1/1/1	-
3	TAR	A	295	-	2/2/4/4	0/4/12/12	-
2	EDO	B	293	-	-	1/1/1/1	-
2	EDO	B	294	-	-	1/1/1/1	-
2	EDO	B	295	-	-	1/1/1/1	-
2	EDO	B	296	-	-	1/1/1/1	-
2	EDO	C	293	-	-	1/1/1/1	-
2	EDO	C	294	-	-	0/1/1/1	-
2	EDO	C	295	-	-	1/1/1/1	-

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	D	293	-	-	1/1/1/1	-
2	EDO	D	294	-	-	0/1/1/1	-
2	EDO	D	295	-	-	0/1/1/1	-
2	EDO	D	296	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	295	TAR	C2
3	A	295	TAR	C3

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	293	EDO	O1-C1-C2-O2
2	D	296	EDO	O1-C1-C2-O2
2	B	296	EDO	O1-C1-C2-O2
2	A	294	EDO	O1-C1-C2-O2
2	C	295	EDO	O1-C1-C2-O2
2	B	295	EDO	O1-C1-C2-O2
2	B	294	EDO	O1-C1-C2-O2
2	B	293	EDO	O1-C1-C2-O2
2	D	293	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	295	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	279/293 (95%)	0.60	24 (8%)	10 13	31, 39, 52, 72	0
1	B	280/293 (95%)	0.61	20 (7%)	16 20	29, 39, 54, 71	0
1	C	280/293 (95%)	0.71	22 (7%)	12 16	30, 39, 48, 65	0
1	D	279/293 (95%)	0.56	18 (6%)	19 24	31, 39, 49, 65	0
All	All	1118/1172 (95%)	0.62	84 (7%)	14 18	29, 39, 51, 72	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	41	GLY	6.3
1	A	43	ILE	5.8
1	C	106	THR	5.8
1	C	41	GLY	5.4
1	C	40	ASP	5.2
1	B	82	SER	5.0
1	D	71	ALA	4.5
1	C	70	GLU	4.4
1	C	56	ASN	4.4
1	B	0	GLY	4.3
1	B	80	ASN	4.2
1	B	51	ASP	4.1
1	B	38	LEU	3.9
1	B	41	GLY	3.9
1	D	86	THR	3.7
1	C	51	ASP	3.6
1	C	55	TRP	3.5
1	A	71	ALA	3.5
1	C	3	THR	3.4
1	A	80	ASN	3.3
1	D	258	GLY	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	39	ASP	3.1
1	B	9	ILE	3.0
1	C	0	GLY	2.9
1	B	71	ALA	2.9
1	A	9	ILE	2.9
1	D	43	ILE	2.8
1	C	4	GLY	2.8
1	C	105	ASN	2.8
1	C	62	LEU	2.8
1	D	116	ASN	2.7
1	A	35	PHE	2.7
1	D	42	PHE	2.7
1	C	92	LYS	2.7
1	A	106	THR	2.7
1	B	106	THR	2.7
1	A	173	PHE	2.7
1	D	137	ALA	2.6
1	D	93	THR	2.6
1	A	174	ALA	2.5
1	D	254	ALA	2.5
1	C	93	THR	2.5
1	B	141	ILE	2.5
1	B	291	PRO	2.5
1	B	42	PHE	2.5
1	A	65	ILE	2.5
1	C	49	ARG	2.4
1	D	107	TYR	2.4
1	A	78	PHE	2.4
1	B	39	ASP	2.4
1	C	176	LEU	2.4
1	B	81	GLU	2.4
1	A	39	ASP	2.4
1	D	51	ASP	2.4
1	C	116	ASN	2.3
1	A	30	ILE	2.3
1	C	113	LYS	2.3
1	A	290	GLU	2.3
1	B	105	ASN	2.3
1	D	159	VAL	2.3
1	C	290	GLU	2.3
1	A	8	VAL	2.3
1	B	159	VAL	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	98	VAL	2.2
1	A	141	ILE	2.2
1	A	105	ASN	2.2
1	A	134	ILE	2.2
1	C	80	ASN	2.2
1	B	169	VAL	2.2
1	A	69[A]	ASN	2.2
1	A	88	GLU	2.1
1	B	73	LYS	2.1
1	D	290	GLU	2.1
1	A	42	PHE	2.1
1	D	251	PHE	2.1
1	B	208	GLN	2.1
1	A	0	GLY	2.1
1	D	0	GLY	2.1
1	D	53	VAL	2.1
1	A	172	LYS	2.1
1	B	36	ASP	2.0
1	D	259	ILE	2.0
1	A	61	ILE	2.0
1	A	219	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EDO	B	295	4/4	0.62	0.37	67,72,75,80	0
2	EDO	D	296	4/4	0.78	0.18	68,68,70,72	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	TAR	A	295	10/10	0.82	0.30	30,47,53,58	0
2	EDO	D	295	4/4	0.87	0.17	29,36,40,46	0
2	EDO	C	295	4/4	0.88	0.13	59,61,61,61	0
2	EDO	A	294	4/4	0.89	0.32	65,67,68,68	0
2	EDO	B	294	4/4	0.90	0.17	25,27,31,32	0
2	EDO	B	296	4/4	0.90	0.35	55,59,61,66	0
2	EDO	B	293	4/4	0.91	0.15	47,47,48,57	0
2	EDO	C	293	4/4	0.92	0.20	42,44,48,49	0
2	EDO	D	293	4/4	0.92	0.13	48,53,56,60	0
2	EDO	A	293	4/4	0.93	0.18	26,27,28,28	0
2	EDO	C	294	4/4	0.94	0.12	29,30,33,39	0
2	EDO	D	294	4/4	0.95	0.11	27,32,32,33	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.