



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 06:51 pm GMT

PDB ID : 5D4C
Title : Crystal structure of Thermus thermophilus product complex for transcription initiation with ATP and CTP
Authors : Zhang, Y.; Ebright, R.H.
Deposited on : 2015-08-07
Resolution : 3.28 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1138 (3.34-3.22)
Clashscore	122126	1055 (3.32-3.24)
Ramachandran outliers	120053	1037 (3.32-3.24)
Sidechain outliers	120020	1036 (3.32-3.24)
RSRZ outliers	108989	1104 (3.34-3.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background-color: green;"></div> <div style="position: absolute; top: 0; left: 60%; width: 10%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; top: 0; left: 73%; width: 27%; height: 10px; background-color: grey;"></div> </div> <div>60% 13% 27%</div> </div>
1	B	315	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background-color: green;"></div> <div style="position: absolute; top: 0; left: 59%; width: 11%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; top: 0; left: 70%; width: 30%; height: 10px; background-color: grey;"></div> </div> <div>59% 11% 30%</div> </div>
1	K	315	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background-color: green;"></div> <div style="position: absolute; top: 0; left: 61%; width: 12%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; top: 0; left: 73%; width: 27%; height: 10px; background-color: grey;"></div> </div> <div>61% 12% 27%</div> </div>
1	L	315	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background-color: green;"></div> <div style="position: absolute; top: 0; left: 58%; width: 12%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; top: 0; left: 70%; width: 30%; height: 10px; background-color: grey;"></div> </div> <div>58% 12% 30%</div> </div>
2	C	1119	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background-color: green;"></div> <div style="position: absolute; top: 0; left: 82%; width: 17%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; top: 0; left: 99%; width: 1%; height: 10px; background-color: grey;"></div> </div> <div>82% 17% .</div> </div>
2	M	1119	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; height: 10px; background-color: red;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background-color: green;"></div> <div style="position: absolute; top: 0; left: 5%; width: 5%; height: 10px; background-color: red;"></div> <div style="position: absolute; top: 0; left: 76%; width: 19%; height: 10px; background-color: yellow;"></div> <div style="position: absolute; top: 0; left: 95%; width: 5%; height: 10px; background-color: grey;"></div> </div> <div>5% 76% 19% . .</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	1524	
3	N	1524	
4	E	99	
4	O	99	
5	F	443	
5	P	443	
6	G	19	
6	R	19	
7	H	27	
7	S	27	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	MG	B	2001	-	-	-	X

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 56600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	231	Total	C	N	O	S	0	0	0
			1809	1155	315	337	2			
1	B	222	Total	C	N	O	S	0	0	0
			1750	1120	303	325	2			
1	K	231	Total	C	N	O	S	0	0	0
			1809	1155	315	337	2			
1	L	222	Total	C	N	O	S	0	0	0
			1750	1120	303	325	2			

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	1112	Total	C	N	O	S	0	0	0
			8774	5550	1565	1635	24			
2	M	1080	Total	C	N	O	S	0	0	0
			8508	5375	1522	1587	24			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	1486	Total	C	N	O	S	0	0	0
			11738	7441	2067	2195	35			
3	N	1486	Total	C	N	O	S	0	0	0
			11738	7441	2067	2195	35			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	94	Total	C	N	O	S	0	0	0
			758	483	132	139	4			
4	O	94	Total	C	N	O	S	0	0	0
			758	483	132	139	4			

- Molecule 5 is a protein called RNA polymerase sigma factor SigA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	346	Total	C	N	O	S	0	0	0
			2807	1770	509	524	4			
5	P	346	Total	C	N	O	S	0	0	0
			2807	1770	509	524	4			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-19	MET	-	initiating methionine	UNP Q5SKW1
F	-18	GLY	-	expression tag	UNP Q5SKW1
F	-17	SER	-	expression tag	UNP Q5SKW1
F	-16	SER	-	expression tag	UNP Q5SKW1
F	-15	HIS	-	expression tag	UNP Q5SKW1
F	-14	HIS	-	expression tag	UNP Q5SKW1
F	-13	HIS	-	expression tag	UNP Q5SKW1
F	-12	HIS	-	expression tag	UNP Q5SKW1
F	-11	HIS	-	expression tag	UNP Q5SKW1
F	-10	HIS	-	expression tag	UNP Q5SKW1
F	-9	SER	-	expression tag	UNP Q5SKW1
F	-8	SER	-	expression tag	UNP Q5SKW1
F	-7	GLY	-	expression tag	UNP Q5SKW1
F	-6	LEU	-	expression tag	UNP Q5SKW1
F	-5	VAL	-	expression tag	UNP Q5SKW1
F	-4	PRO	-	expression tag	UNP Q5SKW1
F	-3	ARG	-	expression tag	UNP Q5SKW1
F	-2	GLY	-	expression tag	UNP Q5SKW1
F	-1	SER	-	expression tag	UNP Q5SKW1
F	0	HIS	-	expression tag	UNP Q5SKW1
P	-19	MET	-	initiating methionine	UNP Q5SKW1
P	-18	GLY	-	expression tag	UNP Q5SKW1
P	-17	SER	-	expression tag	UNP Q5SKW1
P	-16	SER	-	expression tag	UNP Q5SKW1
P	-15	HIS	-	expression tag	UNP Q5SKW1
P	-14	HIS	-	expression tag	UNP Q5SKW1
P	-13	HIS	-	expression tag	UNP Q5SKW1
P	-12	HIS	-	expression tag	UNP Q5SKW1
P	-11	HIS	-	expression tag	UNP Q5SKW1
P	-10	HIS	-	expression tag	UNP Q5SKW1
P	-9	SER	-	expression tag	UNP Q5SKW1
P	-8	SER	-	expression tag	UNP Q5SKW1
P	-7	GLY	-	expression tag	UNP Q5SKW1
P	-6	LEU	-	expression tag	UNP Q5SKW1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
P	-5	VAL	-	expression tag	UNP Q5SKW1
P	-4	PRO	-	expression tag	UNP Q5SKW1
P	-3	ARG	-	expression tag	UNP Q5SKW1
P	-2	GLY	-	expression tag	UNP Q5SKW1
P	-1	SER	-	expression tag	UNP Q5SKW1
P	0	HIS	-	expression tag	UNP Q5SKW1

- Molecule 6 is a DNA chain called DNA (5'-D(*CP*CP*T*GP*CP*AP*TP*CP*CP*GP*T
P*GP*AP*GP*TP*AP*GP*AP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	G	15	Total	C	N	O	P	0	0	0
			308	147	60	87	14			
6	R	15	Total	C	N	O	P	0	0	0
			308	147	60	87	14			

- Molecule 7 is a DNA chain called DNA (27-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	H	20	Total	C	N	O	P	0	0	0
			414	197	82	116	19			
7	S	18	Total	C	N	O	P	0	0	0
			371	177	72	105	17			


- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	P	1	Total	Mg	0	0
			1	1		
8	D	3	Total	Mg	0	0
			3	3		
8	K	1	Total	Mg	0	0
			1	1		
8	B	1	Total	Mg	0	0
			1	1		
8	N	3	Total	Mg	0	0
			3	3		
8	L	1	Total	Mg	0	0
			1	1		
8	F	1	Total	Mg	0	0
			1	1		

- Molecule 9 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	D	2	Total	Zn	0	0
			2	2		
9	N	2	Total	Zn	0	0
			2	2		

- Molecule 10 is CYTIDINE-5'-MONOPHOSPHATE (three-letter code: C) (formula: $C_9H_{14}N_3O_8P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	D	1	Total	C	N	O	P	0	0
			20	9	3	7	1		
10	N	1	Total	C	N	O	P	0	0
			20	9	3	7	1		

- Molecule 11 is CYTIDINE-5'-TRIPHOSPHATE (three-letter code: CTP) (formula: $C_9H_{16}N_3O_{14}P_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	D	1	Total	O	P		0	0
			9	7	2			
11	N	1	Total	O	P		0	0
			9	7	2			

- Molecule 12 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


Continued on next page...

Continued from previous page...



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	N	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	3	Total	O	0	0
			3	3		
13	B	2	Total	O	0	0
			2	2		
13	C	11	Total	O	0	0
			11	11		
13	D	18	Total	O	0	0
			18	18		
13	E	1	Total	O	0	0
			1	1		
13	G	2	Total	O	0	0
			2	2		
13	K	3	Total	O	0	0
			3	3		
13	L	1	Total	O	0	0
			1	1		
13	M	1	Total	O	0	0
			1	1		
13	N	14	Total	O	0	0
			14	14		
13	O	1	Total	O	0	0
			1	1		
13	P	1	Total	O	0	0
			1	1		





• Molecule 3: DNA-directed RNA polymerase subunit beta'






- Molecule 3: DNA-directed RNA polymerase subunit beta'




- Molecule 4: DNA-directed RNA polymerase subunit omega




- Molecule 4: DNA-directed RNA polymerase subunit omega


- Chain F:



- Chain P: 



- Chain G: 47% 21% 11% 21%



- Molecule 6: DNA (5'-D(*CP*CP*T*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*AP*GP*AP*G)-3')

Chain R: 





- Molecule 7: DNA (27-MER)

Chain H: 



- Molecule 7: DNA (27-MER)

Chain S: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	185.96Å 103.64Å 297.42Å 90.00° 98.30° 90.00°	Depositor
Resolution (Å)	49.20 – 3.28 49.50 – 3.28	Depositor EDS
% Data completeness (in resolution range)	89.5 (49.20-3.28) 89.7 (49.50-3.28)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.91 (at 3.25Å)	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
R, R_{free}	0.211 , 0.257 0.213 , 0.259	Depositor DCC
R_{free} test set	7741 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	57.7	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 31.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	56600	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 52.73 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.6669e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CTP, ZN, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/1841	0.45	0/2504
1	B	0.23	0/1781	0.47	0/2420
1	K	0.24	0/1841	0.44	0/2504
1	L	0.24	0/1781	0.45	0/2420
2	C	0.24	0/8941	0.44	0/12092
2	M	0.24	0/8669	0.45	0/11724
3	D	0.24	0/11944	0.44	0/16149
3	N	0.24	0/11944	0.44	0/16149
4	E	0.23	0/772	0.42	0/1040
4	O	0.22	0/772	0.42	0/1040
5	F	0.24	0/2852	0.40	0/3837
5	P	0.24	0/2852	0.43	0/3837
6	G	0.51	0/346	1.14	2/533 (0.4%)
6	R	0.51	0/346	1.09	1/533 (0.2%)
7	H	0.59	0/465	1.06	0/715
7	S	0.51	0/416	1.06	1/639 (0.2%)
All	All	0.25	0/57563	0.47	4/78136 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	D	0	1
3	N	0	1
All	All	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	S	1	DT	O4'-C1'-N1	6.12	112.28	108.00
6	G	15	DT	O4'-C4'-C3'	-5.81	102.17	104.50
6	G	5	DC	O4'-C1'-N1	5.27	111.69	108.00
6	R	15	DT	O4'-C4'-C3'	-5.10	102.46	104.50

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	D	633	VAL	Peptide
3	N	138	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1809	0	1863	26	0
1	B	1750	0	1802	25	0
1	K	1809	0	1863	25	0
1	L	1750	0	1802	25	0
2	C	8774	0	8877	109	0
2	M	8508	0	8605	136	0
3	D	11738	0	11972	146	0
3	N	11738	0	11971	163	0
4	E	758	0	770	9	0
4	O	758	0	770	5	0
5	F	2807	0	2882	30	0
5	P	2807	0	2882	54	0
6	G	308	0	170	5	0
6	R	308	0	170	3	0
7	H	414	0	227	12	0
7	S	371	0	205	13	0
8	B	1	0	0	0	0
8	D	3	0	0	0	0
8	F	1	0	0	0	0
8	K	1	0	0	0	0
8	L	1	0	0	0	0
8	N	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	P	1	0	0	0	0
9	D	2	0	0	0	0
9	N	2	0	0	0	0
10	D	20	0	11	0	0
10	N	20	0	11	1	0
11	D	9	0	0	0	0
11	N	9	0	0	0	0
12	D	31	0	11	0	0
12	N	31	0	11	0	0
13	A	3	0	0	0	0
13	B	2	0	0	0	0
13	C	11	0	0	0	0
13	D	18	0	0	0	0
13	E	1	0	0	0	0
13	G	2	0	0	0	0
13	K	3	0	0	0	0
13	L	1	0	0	0	0
13	M	1	0	0	0	0
13	N	14	0	0	0	0
13	O	1	0	0	0	0
13	P	1	0	0	0	0
All	All	56600	0	56875	707	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 707 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:243:ARG:NH1	7:H:9:DG:O6	1.99	0.93
2:M:165:LEU:HB2	2:M:168:ARG:HG3	1.59	0.84
2:M:758:ARG:HH21	2:M:788:THR:HB	1.45	0.80
3:D:1495:ILE:HG12	4:E:88:GLU:HG3	1.67	0.77
2:M:770:GLU:HB3	5:P:354:LEU:HG	1.65	0.77

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/315 (73%)	226 (99%)	3 (1%)	0	100	100
1	B	218/315 (69%)	212 (97%)	5 (2%)	1 (0%)	31	65
1	K	229/315 (73%)	226 (99%)	3 (1%)	0	100	100
1	L	218/315 (69%)	212 (97%)	6 (3%)	0	100	100
2	C	1108/1119 (99%)	1085 (98%)	22 (2%)	1 (0%)	53	84
2	M	1074/1119 (96%)	1049 (98%)	25 (2%)	0	100	100
3	D	1482/1524 (97%)	1451 (98%)	31 (2%)	0	100	100
3	N	1482/1524 (97%)	1446 (98%)	35 (2%)	1 (0%)	53	84
4	E	92/99 (93%)	89 (97%)	3 (3%)	0	100	100
4	O	92/99 (93%)	89 (97%)	3 (3%)	0	100	100
5	F	344/443 (78%)	340 (99%)	4 (1%)	0	100	100
5	P	344/443 (78%)	338 (98%)	5 (2%)	1 (0%)	43	74
All	All	6912/7630 (91%)	6763 (98%)	145 (2%)	4 (0%)	53	84

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	P	361	LEU
3	N	207	PHE
1	B	8	ALA
2	C	23	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/273 (73%)	199 (100%)	1 (0%)	90	94
1	B	195/273 (71%)	193 (99%)	2 (1%)	78	87
1	K	200/273 (73%)	197 (98%)	3 (2%)	67	84
1	L	195/273 (71%)	191 (98%)	4 (2%)	56	78
2	C	936/941 (100%)	910 (97%)	26 (3%)	47	73
2	M	908/941 (96%)	876 (96%)	32 (4%)	39	69
3	D	1253/1279 (98%)	1225 (98%)	28 (2%)	55	78
3	N	1253/1279 (98%)	1221 (97%)	32 (3%)	49	75
4	E	82/88 (93%)	80 (98%)	2 (2%)	52	76
4	O	82/88 (93%)	76 (93%)	6 (7%)	15	44
5	F	301/388 (78%)	294 (98%)	7 (2%)	53	77
5	P	301/388 (78%)	280 (93%)	21 (7%)	16	47
All	All	5906/6484 (91%)	5742 (97%)	164 (3%)	47	73

5 of 164 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	M	118	ILE
2	M	513	VAL
5	P	232	ARG
2	M	162	ILE
2	M	368	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
2	M	219	GLN
2	M	1047	HIS
5	P	218	GLN
2	M	343	GLN
2	M	538	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 15 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
10	C	D	2005	8,12	14,21,22	0.81	0	15,30,33	0.58	0
11	CTP	D	2006	8	8,8,30	1.43	1 (12%)	9,13,47	1.21	1 (11%)
12	ATP	D	2008	10	27,33,33	1.55	5 (18%)	27,52,52	2.23	4 (14%)
10	C	N	1605	8,12	14,21,22	0.87	0	15,30,33	0.51	0
11	CTP	N	1606	8	8,8,30	1.01	0	9,13,47	1.03	1 (11%)
12	ATP	N	1608	10	27,33,33	1.57	6 (22%)	27,52,52	2.24	5 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	C	D	2005	8,12	-	0/3/25/26	0/2/2/2
11	CTP	D	2006	8	-	0/6/6/38	0/0/0/2
12	ATP	D	2008	10	-	0/18/38/38	0/3/3/3
10	C	N	1605	8,12	-	0/3/25/26	0/2/2/2
11	CTP	N	1606	8	-	0/6/6/38	0/0/0/2
12	ATP	N	1608	10	-	0/18/38/38	0/3/3/3

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	D	2008	ATP	PG-O3B	-2.25	1.56	1.60
12	N	1608	ATP	PG-O3B	-2.24	1.56	1.60
12	N	1608	ATP	O3'-C3'	-2.12	1.37	1.43
12	N	1608	ATP	C2'-C3'	-2.06	1.48	1.53
12	D	2008	ATP	O3'-C3'	-2.03	1.38	1.43

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	N	1608	ATP	N3-C2-N1	-9.35	120.86	128.86
12	D	2008	ATP	N3-C2-N1	-9.04	121.12	128.86
12	D	2008	ATP	PA-O3A-PB	-3.70	120.19	132.63
12	N	1608	ATP	PA-O3A-PB	-3.55	120.70	132.63
12	N	1608	ATP	PB-O3B-PG	-3.51	120.85	132.63

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	N	1605	C	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/315 (73%)	-0.32	3 (1%) 77 76	13, 36, 63, 128	1 (0%)
1	B	222/315 (70%)	-0.25	0 100 100	16, 42, 80, 94	0
1	K	231/315 (73%)	-0.14	4 (1%) 70 67	18, 47, 73, 123	1 (0%)
1	L	222/315 (70%)	-0.27	0 100 100	17, 46, 87, 114	0
2	C	1112/1119 (99%)	-0.28	11 (0%) 82 82	4, 31, 89, 122	3 (0%)
2	M	1080/1119 (96%)	0.18	54 (5%) 29 27	5, 58, 121, 135	2 (0%)
3	D	1486/1524 (97%)	-0.23	15 (1%) 82 82	3, 32, 90, 122	4 (0%)
3	N	1486/1524 (97%)	-0.08	36 (2%) 59 56	3, 40, 103, 143	4 (0%)
4	E	94/99 (94%)	-0.43	0 100 100	11, 29, 67, 94	0
4	O	94/99 (94%)	-0.40	0 100 100	13, 40, 81, 99	0
5	F	346/443 (78%)	-0.10	9 (2%) 56 53	13, 47, 108, 124	0
5	P	346/443 (78%)	0.30	27 (7%) 13 12	29, 74, 142, 167	0
6	G	15/19 (78%)	-0.33	0 100 100	15, 34, 132, 135	0
6	R	15/19 (78%)	-0.41	0 100 100	24, 43, 131, 133	0
7	H	20/27 (74%)	-0.30	0 100 100	36, 66, 117, 125	0
7	S	18/27 (66%)	-0.21	0 100 100	63, 80, 125, 140	0
All	All	7018/7722 (90%)	-0.12	159 (2%) 60 58	3, 42, 105, 167	15 (0%)

The worst 5 of 159 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	P	377	ASP	7.9
5	P	391	GLY	7.5
5	P	381	HIS	5.9
5	P	392	VAL	5.9
3	N	70	GLY	5.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
8	MG	B	2001	1/1	0.37	0.49	70,70,70,70	0
8	MG	L	2001	1/1	0.70	0.28	72,72,72,72	0
9	ZN	N	1603	1/1	0.91	0.06	128,128,128,128	0
12	ATP	N	1608	31/31	0.92	0.17	28,39,97,99	0
10	C	N	1605	20/21	0.93	0.17	23,34,41,41	0
11	CTP	D	2006	9/29	0.94	0.30	27,37,51,69	0
10	C	D	2005	20/21	0.94	0.17	16,22,28,34	0
8	MG	D	2004	1/1	0.95	0.18	29,29,29,29	0
8	MG	D	2007	1/1	0.95	0.34	17,17,17,17	0
12	ATP	D	2008	31/31	0.95	0.15	18,28,68,77	0
8	MG	N	1607	1/1	0.95	0.30	30,30,30,30	0
8	MG	F	2001	1/1	0.95	0.05	31,31,31,31	0
8	MG	K	901	1/1	0.96	0.26	36,36,36,36	0
11	CTP	N	1606	9/29	0.97	0.32	34,40,54,72	0
8	MG	P	2001	1/1	0.98	0.06	77,77,77,77	0
9	ZN	D	2002	1/1	0.98	0.03	92,92,92,92	0
8	MG	N	1601	1/1	0.99	0.16	25,25,25,25	0
8	MG	D	2003	1/1	0.99	0.26	6,6,6,6	0
8	MG	N	1604	1/1	0.99	0.21	12,12,12,12	0
9	ZN	N	1602	1/1	0.99	0.14	8,8,8,8	0
9	ZN	D	2001	1/1	0.99	0.12	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.