



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 02:38 pm GMT

PDB ID : 1D5L
Title : CRYSTAL STRUCTURE OF CYANIDE-BOUND HUMAN MYELOPER-
OXIDASE ISOFORM C AT PH 5.5
Authors : Fiedler, T.J.; Davey, C.A.; Fenna, R.E.
Deposited on : 1999-10-07
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	6115 (1.90-1.90)
Ramachandran outliers	120053	6048 (1.90-1.90)
Sidechain outliers	120020	6048 (1.90-1.90)
RSRZ outliers	108989	5379 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	104	<div> <div>2%</div> <div>89%</div> <div>11%</div> </div>
1	B	104	<div> <div>96%</div> <div>.</div> </div>
2	C	466	<div> <div>%</div> <div>88%</div> <div>11%</div> </div>
2	D	466	<div> <div>3%</div> <div>89%</div> <div>9%</div> <div>.</div> </div>

2 Entry composition [i](#)

There are 13 unique types of molecules in this entry. The entry contains 10347 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MYELOPEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	104	Total	C	N	O	S	0	0	0
			838	529	148	156	5			
1	B	104	Total	C	N	O	S	0	0	0
			838	529	148	156	5			

- Molecule 2 is a protein called MYELOPEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	466	Total	C	N	O	S	0	0	0
			3733	2351	687	668	27			
2	D	466	Total	C	N	O	S	0	0	0
			3733	2351	687	668	27			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	150	CSO	CYS	MODIFIED RESIDUE	UNP P05164
D	150	CSO	CYS	MODIFIED RESIDUE	UNP P05164

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is CYANIDE ION (three-letter code: CYN) (formula: CN).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	N	0	0
			2	1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	N	0	0
			2	1	1		
5	B	1	Total	C	N	0	0
			2	1	1		
5	D	1	Total	C	N	0	0
			2	1	1		

- Molecule 6 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).






Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	C	1	Total	C	N	O	0	0
			14	8	1	5		
7	C	1	Total	C	N	O	0	0
			14	8	1	5		
7	C	1	Total	C	N	O	0	0
			14	8	1	5		
7	C	1	Total	C	N	O	0	0
			14	8	1	5		
7	D	1	Total	C	N	O	0	0
			14	8	1	5		
7	D	1	Total	C	N	O	0	0
			14	8	1	5		
7	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 8 is BETA-D-MANNOSE (three-letter code: BMA) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	C	1	Total	C	O	0	0
			11	6	5		
8	D	1	Total	C	O	0	0
			11	6	5		

- Molecule 9 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: $C_6H_{12}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	C	1	Total	C	O	0	0
			11	6	5		
9	C	1	Total	C	O	0	0
			11	6	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	D	1	Total	C	O	0	0
			11	6	5		
9	D	1	Total	C	O	0	0
			11	6	5		

- Molecule 10 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula: $C_6H_{12}O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	C	1	Total	C	O	0	0
			10	6	4		
10	D	1	Total	C	O	0	0
			10	6	4		

- Molecule 11 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	D	1	Total	Ca	0	0
			1	1		
11	C	1	Total	Ca	0	0
			1	1		

- Molecule 12 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	C	1	Total	C	O	0	0
			4	2	2		
12	C	1	Total	C	O	0	0
			4	2	2		
12	C	1	Total	C	O	0	0
			4	2	2		
12	C	1	Total	C	O	0	0
			4	2	2		
12	D	1	Total	C	O	0	0
			4	2	2		
12	D	1	Total	C	O	0	0
			4	2	2		
12	D	1	Total	C	O	0	0
			4	2	2		



- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	96	Total	O	0	0
			96	96		
13	C	338	Total	O	0	0
			338	338		
13	B	105	Total	O	0	0
			105	105		
13	D	327	Total	O	0	0
			327	327		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: MYELOPEROXIDASE





- Molecule 1: MYELOPEROXIDASE



- Molecule 2: MYELOPEROXIDASE



- Molecule 2: MYELOPEROXIDASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	111.22Å 63.51Å 92.34Å 90.00° 97.43° 90.00°	Depositor
Resolution (Å)	30.00 – 1.90 45.88 – 1.91	Depositor EDS
% Data completeness (in resolution range)	91.3 (30.00-1.90) 96.0 (45.88-1.91)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.42 (at 1.91Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.172 , 0.215 0.170 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	13.2	Xtriage
Anisotropy	0.393	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 55.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10347	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, BMA, NAG, CL, SO4, CA, FUC, ACT, HEM, CYN, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/863	0.67	0/1174
1	B	0.57	0/863	0.66	0/1174
2	C	0.54	0/3811	0.59	0/5168
2	D	0.52	0/3811	0.57	0/5168
All	All	0.54	0/9348	0.60	0/12684

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	838	0	798	8	0
1	B	838	0	798	3	0
2	C	3733	0	3725	34	0
2	D	3733	0	3725	31	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	5	0	0	0	0
4	C	5	0	0	0	0
4	D	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	2	0	0	0	0
5	B	2	0	0	0	0
5	C	2	0	0	1	0
5	D	2	0	0	0	0
6	A	43	0	30	3	0
6	B	43	0	30	3	0
7	C	56	0	49	0	0
7	D	56	0	49	0	0
8	C	11	0	8	0	0
8	D	11	0	8	0	0
9	C	22	0	20	0	0
9	D	22	0	20	0	0
10	C	10	0	10	0	0
10	D	10	0	10	0	0
11	C	1	0	0	0	0
11	D	1	0	0	0	0
12	C	16	0	12	0	0
12	D	12	0	9	1	0
13	A	96	0	0	2	0
13	B	105	0	0	0	0
13	C	338	0	0	2	0
13	D	327	0	0	3	0
All	All	10347	0	9301	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 74 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:411:MET:HE2	2:D:415:LEU:HD21	1.76	0.68
2:D:113:VAL:HG21	2:D:122:GLN:HB2	1.81	0.62
2:C:465:MET:HE1	2:C:471:PRO:HG3	1.81	0.62
13:A:1243(A):HOH:O	2:C:129:LYS:HD3	1.99	0.61
2:C:333:ARG:HH11	2:C:421:ASN:HD22	1.48	0.60

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	102/104 (98%)	100 (98%)	2 (2%)	0	100	100
1	B	102/104 (98%)	99 (97%)	3 (3%)	0	100	100
2	C	463/466 (99%)	453 (98%)	9 (2%)	1 (0%)	49	40
2	D	463/466 (99%)	453 (98%)	10 (2%)	0	100	100
All	All	1130/1140 (99%)	1105 (98%)	24 (2%)	1 (0%)	53	44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	114	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	90/90 (100%)	89 (99%)	1 (1%)	76	75
1	B	90/90 (100%)	89 (99%)	1 (1%)	76	75
2	C	410/410 (100%)	402 (98%)	8 (2%)	58	53
2	D	410/410 (100%)	402 (98%)	8 (2%)	58	53
All	All	1000/1000 (100%)	982 (98%)	18 (2%)	62	58

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	523	GLN
1	B	54	ASN
2	D	267	LEU
2	C	318	ASP
2	C	497	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	421	ASN
2	D	549	ASN
1	B	54	ASN
2	C	200	ASN
2	C	526	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CSO	C	150	2	4,6,7	1.50	1 (25%)	1,6,8	1.75	0
2	CSO	D	150	2	4,6,7	1.99	1 (25%)	1,6,8	1.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSO	C	150	2	-	0/1/5/7	0/0/0/0
2	CSO	D	150	2	-	0/1/5/7	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	150	CSO	CA-C	2.70	1.53	1.50
2	D	150	CSO	CA-C	3.64	1.55	1.50

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 36 ligands modelled in this entry, 4 are monoatomic - leaving 32 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	1602	-	4,4,4	0.21	0	6,6,6	0.13	0
5	CYN	A	1844	6	0,1,1	0.00	-	0,0,0	0.00	-
6	HEM	A	605	1,2,5	27,50,50	1.47	5 (18%)	17,82,82	1.12	1 (5%)
5	CYN	B	2844	6	0,1,1	0.00	-	0,0,0	0.00	-
6	HEM	B	605	1,2,5	27,50,50	1.44	5 (18%)	17,82,82	1.20	2 (11%)
4	SO4	C	1603	-	4,4,4	0.30	0	6,6,6	0.11	0
12	ACT	C	1604	-	1,3,3	2.39	1 (100%)	0,3,3	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
12	ACT	C	1606	-	1,3,3	2.28	1 (100%)	0,3,3	0.00	-
12	ACT	C	1607	-	1,3,3	3.39	1 (100%)	0,3,3	0.00	-
12	ACT	C	1612	-	1,3,3	3.09	1 (100%)	0,3,3	0.00	-
7	NAG	C	1620	2	14,14,15	0.83	1 (7%)	17,19,21	0.86	0
7	NAG	C	1630	2	14,14,15	0.46	0	17,19,21	0.69	0
7	NAG	C	1640	10,2,7	14,14,15	0.70	0	17,19,21	0.85	0
7	NAG	C	1641	8,7	14,14,15	0.51	0	17,19,21	0.71	0
8	BMA	C	1642	9,7	11,11,12	0.49	0	15,15,17	0.51	0
9	MAN	C	1643	8	11,11,12	0.64	0	15,15,17	0.54	0
9	MAN	C	1644	8	11,11,12	0.49	0	15,15,17	0.55	0
10	FUC	C	1645	7	9,10,11	0.47	0	13,14,16	0.69	0
5	CYN	C	1889	-	0,1,1	0.00	-	0,0,0	0.00	-
4	SO4	D	2602	-	4,4,4	0.34	0	6,6,6	0.06	0
12	ACT	D	2604	-	1,3,3	2.94	1 (100%)	0,3,3	0.00	-
12	ACT	D	2606	-	1,3,3	2.30	1 (100%)	0,3,3	0.00	-
12	ACT	D	2607	-	1,3,3	2.80	1 (100%)	0,3,3	0.00	-
7	NAG	D	2620	2	14,14,15	0.49	0	17,19,21	0.79	1 (5%)
7	NAG	D	2630	2	14,14,15	0.44	0	17,19,21	0.75	1 (5%)
7	NAG	D	2640	10,2,7	14,14,15	0.50	0	17,19,21	0.99	1 (5%)
7	NAG	D	2641	8,7	14,14,15	0.42	0	17,19,21	0.74	0
8	BMA	D	2642	9,7	11,11,12	0.34	0	15,15,17	0.60	0
9	MAN	D	2643	8	11,11,12	0.67	0	15,15,17	0.63	0
9	MAN	D	2644	8	11,11,12	0.53	0	15,15,17	0.53	0
10	FUC	D	2645	7	9,10,11	0.58	0	13,14,16	0.59	0
5	CYN	D	2889	-	0,1,1	0.00	-	0,0,0	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SO4	A	1602	-	-	0/0/0/0	0/0/0/0
5	CYN	A	1844	6	-	0/0/0/0	0/0/0/0
6	HEM	A	605	1,2,5	-	0/6/54/54	0/0/8/8
5	CYN	B	2844	6	-	0/0/0/0	0/0/0/0
6	HEM	B	605	1,2,5	-	0/6/54/54	0/0/8/8
4	SO4	C	1603	-	-	0/0/0/0	0/0/0/0
12	ACT	C	1604	-	-	0/0/0/0	0/0/0/0
12	ACT	C	1606	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	ACT	C	1607	-	-	0/0/0/0	0/0/0/0
12	ACT	C	1612	-	-	0/0/0/0	0/0/0/0
7	NAG	C	1620	2	-	0/6/23/26	0/1/1/1
7	NAG	C	1630	2	-	0/6/23/26	0/1/1/1
7	NAG	C	1640	10,2,7	-	0/6/23/26	0/1/1/1
7	NAG	C	1641	8,7	-	0/6/23/26	0/1/1/1
8	BMA	C	1642	9,7	-	0/2/19/22	0/1/1/1
9	MAN	C	1643	8	-	0/2/19/22	0/1/1/1
9	MAN	C	1644	8	-	0/2/19/22	0/1/1/1
10	FUC	C	1645	7	-	0/0/17/20	0/1/1/1
5	CYN	C	1889	-	-	0/0/0/0	0/0/0/0
4	SO4	D	2602	-	-	0/0/0/0	0/0/0/0
12	ACT	D	2604	-	-	0/0/0/0	0/0/0/0
12	ACT	D	2606	-	-	0/0/0/0	0/0/0/0
12	ACT	D	2607	-	-	0/0/0/0	0/0/0/0
7	NAG	D	2620	2	-	0/6/23/26	0/1/1/1
7	NAG	D	2630	2	-	0/6/23/26	0/1/1/1
7	NAG	D	2640	10,2,7	-	0/6/23/26	0/1/1/1
7	NAG	D	2641	8,7	-	0/6/23/26	0/1/1/1
8	BMA	D	2642	9,7	-	0/2/19/22	0/1/1/1
9	MAN	D	2643	8	-	0/2/19/22	0/1/1/1
9	MAN	D	2644	8	-	0/2/19/22	0/1/1/1
10	FUC	D	2645	7	-	0/0/17/20	0/1/1/1
5	CYN	D	2889	-	-	0/0/0/0	0/0/0/0

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	605	HEM	C3B-CAB	-3.33	1.41	1.47
6	B	605	HEM	C3C-CAC	-3.16	1.41	1.47
6	A	605	HEM	C3C-CAC	-3.06	1.41	1.47
6	B	605	HEM	C3B-CAB	-2.98	1.41	1.47
6	B	605	HEM	C3B-C2B	-2.86	1.36	1.40

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	D	2630	NAG	C2-N2-C7	-2.14	119.82	122.94
7	D	2640	NAG	O5-C1-C2	-2.06	108.68	111.52
7	D	2620	NAG	C1-O5-C5	2.19	115.20	112.19
6	A	605	HEM	CMC-C2C-C3C	2.21	128.91	124.88
6	B	605	HEM	CMC-C2C-C3C	2.22	128.93	124.88

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	605	HEM	3	0
6	B	605	HEM	3	0
5	C	1889	CYN	1	0
12	D	2606	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	104/104 (100%)	-0.12	2 (1%) 66 70	5, 9, 24, 54	0
1	B	104/104 (100%)	-0.21	0 100 100	6, 10, 20, 28	0
2	C	465/466 (99%)	-0.12	5 (1%) 80 83	4, 10, 24, 40	0
2	D	465/466 (99%)	-0.01	14 (3%) 50 53	5, 13, 30, 53	0
All	All	1138/1140 (99%)	-0.08	21 (1%) 68 71	4, 11, 28, 54	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	113	VAL	8.1
2	D	217	HIS	5.0
2	D	578	ALA	3.7
2	D	355	PRO	3.5
1	A	4	GLN	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CSO	D	150	7/8	0.94	0.09	5,6,9,13	0
2	CSO	C	150	7/8	0.96	0.08	6,9,10,12	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	CYN	D	2889	2/2	0.52	0.20	27,27,27,28	0
12	ACT	D	2604	4/4	0.70	0.28	46,48,48,48	0
4	SO4	D	2602	5/5	0.86	0.18	36,37,37,37	5
7	NAG	C	1630	14/15	0.87	0.20	20,26,34,34	0
12	ACT	C	1612	4/4	0.87	0.25	4,6,7,20	4
7	NAG	D	2630	14/15	0.88	0.19	33,35,37,40	0
9	MAN	D	2643	11/12	0.89	0.14	16,21,24,25	0
12	ACT	C	1607	4/4	0.90	0.10	23,24,24,24	0
12	ACT	C	1604	4/4	0.90	0.19	17,20,20,23	0
12	ACT	C	1606	4/4	0.90	0.11	27,29,29,30	0
12	ACT	D	2606	4/4	0.91	0.13	23,24,25,27	0
7	NAG	D	2620	14/15	0.92	0.18	20,24,28,31	0
9	MAN	C	1643	11/12	0.93	0.13	18,21,22,26	0
7	NAG	C	1620	14/15	0.93	0.12	13,16,20,20	0
12	ACT	D	2607	4/4	0.94	0.13	29,32,32,33	0
10	FUC	C	1645	10/11	0.95	0.09	11,12,13,14	0
8	BMA	C	1642	11/12	0.95	0.09	9,10,13,16	0
5	CYN	C	1889	2/2	0.95	0.08	17,17,17,20	0
9	MAN	D	2644	11/12	0.96	0.09	8,12,13,13	0
9	MAN	C	1644	11/12	0.96	0.11	10,12,15,16	0
7	NAG	C	1640	14/15	0.96	0.08	8,10,14,16	0
7	NAG	C	1641	14/15	0.96	0.10	6,9,10,11	0
4	SO4	C	1603	5/5	0.97	0.14	26,26,29,31	0
10	FUC	D	2645	10/11	0.97	0.08	9,12,13,14	0
7	NAG	D	2641	14/15	0.97	0.08	5,8,11,12	0
8	BMA	D	2642	11/12	0.97	0.07	8,10,11,12	0
6	HEM	B	605	43/43	0.97	0.09	4,7,8,10	0
7	NAG	D	2640	14/15	0.97	0.08	7,12,17,18	0
5	CYN	B	2844	2/2	0.98	0.15	14,14,14,15	0
6	HEM	A	605	43/43	0.98	0.10	3,7,8,10	0
5	CYN	A	1844	2/2	0.99	0.09	10,10,10,12	0
4	SO4	A	1602	5/5	0.99	0.07	25,25,27,28	0
3	CL	A	1601	1/1	1.00	0.12	8,8,8,8	0
11	CA	D	2600	1/1	1.00	0.10	9,9,9,9	0
3	CL	B	2601	1/1	1.00	0.12	10,10,10,10	0
11	CA	C	1600	1/1	1.00	0.07	8,8,8,8	0

6.5 Other polymers [i](#)

There are no such residues in this entry.