



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 10:46 am GMT

PDB ID : 3D6Y
Title : Crystal structure of R275E mutant of BMRR bound to DNA and berberine
Authors : Newberry, K.J.; Brennan, R.G.
Deposited on : 2008-05-20
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	2756 (2.70-2.70)
Ramachandran outliers	120053	2716 (2.70-2.70)
Sidechain outliers	120020	2716 (2.70-2.70)
RSRZ outliers	108989	2376 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	24	
2	A	284	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	965	-	X	-	-
4	GOL	A	966	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2847 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called BMR promoter DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	24	Total	C	N	O	P	0	0	0
			488	232	89	144	23			


- Molecule 2 is a protein called Multidrug-efflux transporter 1 regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	277	Total	C	N	O	S	0	0	0
			2288	1478	361	441	8			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	275	GLU	ARG	ENGINEERED	UNP P39075
A	277	LEU	ALA	ENGINEERED	UNP P39075
A	278	ASP	GLU	ENGINEERED	UNP P39075
A	279	HIS	-	EXPRESSION TAG	UNP P39075
A	280	HIS	-	EXPRESSION TAG	UNP P39075
A	281	HIS	-	EXPRESSION TAG	UNP P39075
A	282	HIS	-	EXPRESSION TAG	UNP P39075
A	283	HIS	-	EXPRESSION TAG	UNP P39075
A	284	HIS	-	EXPRESSION TAG	UNP P39075

- Molecule 3 is BERBERINE (three-letter code: BER) (formula: C₂₀H₁₈NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			25	20	1	4		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	34	Total 34	O 34	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BMR promoter DNA



- Molecule 2: Multidrug-efflux transporter 1 regulator



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, α , β , γ	106.33Å 106.33Å 146.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	86.10 – 2.70 86.10 – 2.70	Depositor EDS
% Data completeness (in resolution range)	98.3 (86.10-2.70) 98.4 (86.10-2.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 2.69Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.227 , 0.259 0.224 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	76.0	Xtriage
Anisotropy	0.397	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 60.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2847	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BER, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.36	0/546	0.67	0/841
2	A	0.44	0/2337	0.68	1/3157 (0.0%)
All	All	0.42	0/2883	0.68	1/3998 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	212	ILE	N-CA-C	-5.81	95.30	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	488	0	271	16	0
2	A	2288	0	2285	101	0
3	A	25	0	18	3	0
4	A	12	0	8	0	0
5	A	34	0	0	1	0
All	All	2847	0	2582	113	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 21.

All (113) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:133:ARG:HD2	2:A:197:ASN:HD21	1.26	0.97
2:A:247:VAL:HB	2:A:274:ILE:HD11	1.47	0.97
2:A:163:ASP:OD1	2:A:196:THR:HG22	1.81	0.80
2:A:156:LYS:HE2	2:A:167:ASN:HD21	1.44	0.80
2:A:255:ILE:HG13	2:A:270:VAL:HG22	1.64	0.78
2:A:133:ARG:HD2	2:A:197:ASN:ND2	2.00	0.77
2:A:162:ALA:O	2:A:196:THR:HG21	1.85	0.76
2:A:239:TYR:O	2:A:243:ARG:HG2	1.93	0.68
1:B:-7:DT:H2''	1:B:-6:DC:H5'	1.75	0.67
2:A:65:PRO:HG2	2:A:68:GLU:HB2	1.75	0.67
2:A:144:PRO:HB3	3:A:964:BER:H102	1.76	0.67
2:A:223:ASN:HD21	2:A:269:ARG:NH1	1.94	0.66
2:A:119:PRO:HB2	2:A:125:PHE:CE2	2.31	0.66
2:A:87:GLN:O	2:A:91:ILE:HG12	1.96	0.65
2:A:158:PHE:CE2	2:A:203:ILE:HA	2.32	0.64
2:A:87:GLN:NE2	2:A:87:GLN:HA	2.14	0.61
2:A:76:GLU:O	2:A:78:GLU:N	2.29	0.61
1:B:5:DG:H2''	1:B:6:DG:O5'	2.00	0.61
2:A:113:LYS:HE2	2:A:117:GLU:OE2	2.02	0.59
2:A:190:ILE:HD13	2:A:190:ILE:H	1.67	0.59
2:A:147:VAL:CG2	2:A:187:TYR:HE1	2.15	0.58
2:A:256:ILE:HG23	2:A:269:ARG:HB3	1.87	0.56
2:A:108:VAL:O	2:A:112:MET:HB2	2.05	0.56
2:A:24:TYR:O	2:A:28:ILE:HG12	2.07	0.55
2:A:163:ASP:CG	2:A:196:THR:HG22	2.27	0.55
2:A:55:LEU:HD22	2:A:59:LEU:HG	1.89	0.54
2:A:158:PHE:HE2	2:A:203:ILE:HA	1.73	0.54
2:A:137:THR:HG23	2:A:190:ILE:HD11	1.89	0.54
2:A:139:ALA:O	2:A:142:ILE:HB	2.08	0.53
2:A:107:LEU:HD13	2:A:195:LEU:CD2	2.39	0.53
1:B:-8:DC:OP1	2:A:43:ARG:HG3	2.09	0.53
2:A:76:GLU:C	2:A:78:GLU:H	2.12	0.53
2:A:258:ILE:HG13	2:A:267:GLU:HG2	1.90	0.53
2:A:8:ILE:HG23	2:A:9:GLY:N	2.24	0.53
2:A:87:GLN:HE21	2:A:87:GLN:HA	1.73	0.52
2:A:8:ILE:HG12	2:A:19:ILE:HG23	1.91	0.51
2:A:40:THR:O	2:A:41:SER:HB2	2.11	0.51
1:B:-7:DT:C2'	1:B:-6:DC:H5'	2.41	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:147:VAL:HG21	2:A:187:TYR:HE1	1.75	0.51
1:B:-8:DC:H2'	1:B:-7:DT:H71	1.92	0.51
2:A:159:ILE:HD12	2:A:159:ILE:N	2.26	0.51
2:A:8:ILE:HD11	2:A:22:LEU:CB	2.41	0.51
2:A:133:ARG:CD	2:A:197:ASN:HD21	2.12	0.50
2:A:268:TYR:CE2	3:A:964:BER:H203	2.47	0.50
2:A:188:ARG:NH1	2:A:250:ASP:OD1	2.45	0.50
2:A:181:SER:O	2:A:184:GLU:HB2	2.12	0.49
2:A:71:LYS:HG3	2:A:72:ALA:N	2.27	0.49
2:A:6:TYR:HE1	2:A:47:ASP:HA	1.77	0.49
2:A:65:PRO:O	2:A:68:GLU:HB3	2.13	0.49
2:A:256:ILE:HD13	2:A:256:ILE:C	2.33	0.49
1:B:9:DG:H2''	1:B:10:DG:OP2	2.12	0.49
1:B:-11:DA:H2''	1:B:-10:DC:O5'	2.14	0.48
2:A:276:ILE:HD12	2:A:276:ILE:C	2.34	0.48
2:A:167:ASN:N	2:A:167:ASN:HD22	2.12	0.48
2:A:156:LYS:HE3	2:A:170:TYR:OH	2.14	0.47
1:B:-13:DT:H2''	1:B:-12:DG:H5''	1.95	0.47
2:A:8:ILE:HD11	2:A:22:LEU:HB3	1.96	0.47
2:A:94:LYS:HB3	2:A:94:LYS:NZ	2.28	0.47
2:A:227:GLU:HG2	2:A:227:GLU:O	2.15	0.47
2:A:127:LEU:CD2	2:A:129:GLU:HG2	2.45	0.46
1:B:-9:DC:H3'	2:A:8:ILE:CG2	2.45	0.46
2:A:107:LEU:HD13	2:A:195:LEU:HD23	1.96	0.46
2:A:203:ILE:HD11	2:A:207:MET:HB2	1.97	0.46
2:A:137:THR:CG2	2:A:190:ILE:HD11	2.45	0.46
1:B:-9:DC:H3'	2:A:8:ILE:HG21	1.97	0.46
2:A:198:LYS:O	2:A:200:ILE:HD13	2.16	0.46
2:A:112:MET:O	2:A:116:MET:HG3	2.16	0.45
2:A:133:ARG:CD	2:A:197:ASN:ND2	2.73	0.45
2:A:158:PHE:CZ	2:A:203:ILE:HA	2.51	0.45
2:A:75:LEU:HD13	2:A:80:LEU:CA	2.47	0.45
2:A:163:ASP:OD1	2:A:196:THR:CG2	2.60	0.45
1:B:3:DG:H2''	1:B:4:DG:C8	2.51	0.45
2:A:65:PRO:HG2	2:A:68:GLU:CB	2.45	0.44
1:B:-12:DG:H2''	1:B:-11:DA:C8	2.52	0.44
2:A:144:PRO:O	3:A:964:BER:H71	2.17	0.44
2:A:64:THR:HG23	2:A:68:GLU:HG2	2.00	0.44
2:A:28:ILE:O	2:A:70:LYS:NZ	2.47	0.44
2:A:163:ASP:C	2:A:165:PHE:H	2.22	0.43
2:A:103:GLN:OE1	2:A:103:GLN:HA	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:158:PHE:CE2	2:A:203:ILE:HG13	2.53	0.43
2:A:201:SER:O	2:A:202:SER:C	2.57	0.43
2:A:204:THR:HG23	2:A:206:ASP:OD2	2.19	0.43
2:A:261:SER:HA	2:A:262:PRO:HD3	1.88	0.43
1:B:-8:DC:H5''	2:A:42:TYR:HA	2.01	0.43
1:B:3:DG:OP1	1:B:3:DG:H4'	2.19	0.42
2:A:163:ASP:OD1	2:A:195:LEU:HB2	2.18	0.42
2:A:264:LYS:H	2:A:264:LYS:HG2	1.70	0.42
2:A:114:ARG:HG2	2:A:114:ARG:NH1	2.34	0.42
2:A:223:ASN:ND2	2:A:269:ARG:NH1	2.65	0.42
1:B:6:DG:H4'	1:B:6:DG:OP1	2.19	0.42
2:A:34:ALA:CB	2:A:46:THR:HG23	2.49	0.42
2:A:114:ARG:HG2	2:A:114:ARG:HH11	1.85	0.42
2:A:256:ILE:HD13	2:A:257:PRO:O	2.19	0.42
2:A:94:LYS:O	2:A:98:LEU:HD23	2.20	0.41
2:A:118:TYR:HA	2:A:119:PRO:HD3	1.92	0.41
2:A:107:LEU:HD13	2:A:195:LEU:HD22	2.02	0.41
2:A:83:PHE:O	2:A:87:GLN:HG2	2.20	0.41
2:A:111:ARG:HG2	2:A:111:ARG:HH11	1.86	0.41
2:A:8:ILE:CG2	2:A:9:GLY:N	2.83	0.41
1:B:-8:DC:H2''	1:B:-7:DT:C6	2.56	0.41
2:A:188:ARG:HH11	2:A:250:ASP:CG	2.24	0.41
2:A:188:ARG:HG2	2:A:189:HIS:ND1	2.35	0.41
2:A:139:ALA:HB3	2:A:187:TYR:O	2.21	0.41
2:A:90:GLN:O	2:A:94:LYS:HG2	2.20	0.41
2:A:98:LEU:HD13	2:A:98:LEU:HA	1.94	0.41
2:A:75:LEU:HD13	2:A:80:LEU:HB2	2.02	0.41
2:A:28:ILE:O	2:A:28:ILE:HG22	2.20	0.40
2:A:138:GLU:HG2	5:A:967:HOH:O	2.19	0.40
2:A:198:LYS:O	2:A:199:GLN:C	2.59	0.40
2:A:11:VAL:HG21	2:A:45:TYR:CD2	2.56	0.40
2:A:277:LEU:HA	2:A:277:LEU:HD23	1.78	0.40
2:A:87:GLN:CA	2:A:87:GLN:HE21	2.32	0.40
2:A:188:ARG:HD2	2:A:250:ASP:OD2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	275/284 (97%)	252 (92%)	20 (7%)	3 (1%)	16	38

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	202	SER
2	A	77	MET
2	A	201	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	255/262 (97%)	236 (92%)	19 (8%)	15	34

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	16	ASN
2	A	51	ILE
2	A	55	LEU
2	A	75	LEU
2	A	76	GLU
2	A	94	LYS
2	A	107	LEU
2	A	118	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	A	125	PHE
2	A	127	LEU
2	A	142	ILE
2	A	165	PHE
2	A	167	ASN
2	A	190	ILE
2	A	204	THR
2	A	219	CYS
2	A	243	ARG
2	A	256	ILE
2	A	269	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	16	ASN
2	A	73	GLN
2	A	87	GLN
2	A	136	GLN
2	A	149	ASN
2	A	167	ASN
2	A	197	ASN
2	A	223	ASN
2	A	265	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	BER	A	964	-	29,29,29	4.80	22 (75%)	43,43,43	2.17	14 (32%)
4	GOL	A	965	-	5,5,5	4.72	5 (100%)	5,5,5	5.70	3 (60%)
4	GOL	A	966	-	5,5,5	4.87	5 (100%)	5,5,5	5.62	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BER	A	964	-	-	0/4/19/19	0/5/5/5
4	GOL	A	965	-	-	0/4/4/4	0/0/0/0
4	GOL	A	966	-	-	0/4/4/4	0/0/0/0

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	966	GOL	C3-C2	-8.45	1.19	1.52
4	A	965	GOL	C3-C2	-7.83	1.22	1.52
3	A	964	BER	C7-C10	-4.85	1.44	1.51
4	A	966	GOL	C1-C2	-3.48	1.38	1.52
4	A	965	GOL	C1-C2	-2.76	1.41	1.52
4	A	965	GOL	O2-C2	-2.69	1.35	1.43
4	A	966	GOL	O2-C2	-2.63	1.35	1.43
3	A	964	BER	C6-C12	2.32	1.45	1.41
3	A	964	BER	C15-C12	2.58	1.48	1.43
3	A	964	BER	O1-C17	2.63	1.48	1.43
4	A	966	GOL	O3-C3	3.00	1.55	1.42
3	A	964	BER	C5-C11	3.12	1.44	1.38
3	A	964	BER	C10-C4	3.19	1.56	1.51
3	A	964	BER	C13-C16	3.34	1.43	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	964	BER	C12-C8	3.63	1.49	1.43
3	A	964	BER	C9-C4	3.66	1.45	1.39
4	A	965	GOL	O3-C3	3.73	1.58	1.42
3	A	964	BER	C14-C11	3.82	1.49	1.39
3	A	964	BER	O3-C15	4.07	1.44	1.38
3	A	964	BER	C5-C2	4.11	1.46	1.39
3	A	964	BER	C16-C18	4.37	1.48	1.39
4	A	966	GOL	O1-C1	4.37	1.60	1.42
3	A	964	BER	C3-C1	4.49	1.45	1.36
4	A	965	GOL	O1-C1	4.63	1.62	1.42
3	A	964	BER	C7-N1	4.78	1.55	1.48
3	A	964	BER	O4-C18	5.04	1.45	1.37
3	A	964	BER	C9-C14	5.23	1.48	1.38
3	A	964	BER	C1-N1	6.17	1.48	1.37
3	A	964	BER	C13-C8	7.18	1.59	1.42
3	A	964	BER	C18-C15	8.33	1.53	1.38
3	A	964	BER	C2-C4	8.41	1.52	1.40
3	A	964	BER	C6-N1	12.90	1.44	1.33

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	964	BER	C6-N1-C1	-4.26	120.36	122.39
3	A	964	BER	C1-C2-C4	-3.65	117.03	120.17
3	A	964	BER	O4-C18-C16	-3.03	119.26	124.36
3	A	964	BER	C12-C6-N1	-3.03	119.83	121.90
3	A	964	BER	C3-C1-C2	-2.22	122.40	124.60
3	A	964	BER	C3-C8-C12	2.15	121.94	119.24
3	A	964	BER	O1-C11-C5	2.36	131.05	127.86
3	A	964	BER	C5-C2-C1	2.52	122.06	118.55
3	A	964	BER	O2-C14-C9	2.76	131.59	127.86
3	A	964	BER	O4-C18-C15	3.15	121.24	116.50
4	A	966	GOL	O1-C1-C2	3.19	125.56	110.11
3	A	964	BER	C10-C7-N1	3.22	113.51	109.42
4	A	965	GOL	O1-C1-C2	3.40	126.59	110.11
3	A	964	BER	C3-C1-N1	4.14	120.49	117.20
3	A	964	BER	C20-O4-C18	4.41	124.07	117.53
3	A	964	BER	C19-O3-C15	4.62	126.44	114.89
4	A	965	GOL	O2-C2-C3	6.44	138.44	109.00
4	A	966	GOL	O2-C2-C3	6.76	139.88	109.00
4	A	966	GOL	O3-C3-C2	10.06	158.92	110.11
4	A	965	GOL	O3-C3-C2	10.42	160.69	110.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	964	BER	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	24/24 (100%)	0.26	1 (4%) 36 34	74, 110, 180, 188	0
2	A	277/284 (97%)	0.85	21 (7%) 14 12	47, 70, 112, 142	0
All	All	301/308 (97%)	0.80	22 (7%) 15 13	47, 72, 119, 188	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	125	PHE	3.4
2	A	127	LEU	3.2
2	A	5	TYR	3.2
2	A	195	LEU	2.7
2	A	229	TYR	2.5
2	A	276	ILE	2.4
2	A	124	VAL	2.4
2	A	190	ILE	2.4
2	A	55	LEU	2.3
2	A	56	ILE	2.3
2	A	77	MET	2.3
2	A	115	GLN	2.2
2	A	118	TYR	2.2
2	A	274	ILE	2.2
2	A	159	ILE	2.1
2	A	270	VAL	2.1
2	A	209	ILE	2.1
2	A	158	PHE	2.1
1	B	-13	DT	2.1
2	A	182	ILE	2.1
2	A	201	SER	2.1
2	A	105	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	A	965	6/6	0.51	0.25	92,108,111,113	0
3	BER	A	964	25/25	0.82	0.34	112,120,124,130	0
4	GOL	A	966	6/6	0.91	0.54	102,113,120,124	0

6.5 Other polymers [i](#)

There are no such residues in this entry.