



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 01:56 pm GMT

PDB ID : 3D8F
Title : Crystal structure of the human Fe65-PTB1 domain with bound phosphate (trigonal crystal form)
Authors : Radzimanowski, J.; Ravaud, S.; Sinning, I.; Wild, K.
Deposited on : 2008-05-23
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2449 (2.70-2.70)
Clashscore	122126	2756 (2.70-2.70)
Ramachandran outliers	120053	2716 (2.70-2.70)
Sidechain outliers	120020	2716 (2.70-2.70)
RSRZ outliers	108989	2376 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 53%, yellow 32%, orange 5%, grey 14%);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> % 53% 32% • 14% </div> </div>
1	B	148	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 57%, yellow 26%, orange 1%, grey 15%);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> % 57% 26% • 15% </div> </div>
1	C	148	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, green 59%, yellow 22%, orange 5%, grey 14%);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> % 59% 22% 5% • 14% </div> </div>
1	D	148	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 52%, yellow 26%, orange 6%, grey 16%);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> 52% 26% 6% 16% </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4000 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Amyloid beta A4 precursor protein-binding family B member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	127	Total	C	N	O	S	0	0	0
			1003	630	180	185	8			
1	B	126	Total	C	N	O	S	0	0	0
			986	622	174	182	8			
1	C	127	Total	C	N	O	S	0	0	0
			1003	630	180	185	8			
1	D	125	Total	C	N	O	S	0	0	0
			976	616	171	181	8			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	506	LEU	-	EXPRESSION TAG	UNP O00213
A	507	GLU	-	EXPRESSION TAG	UNP O00213
A	508	HIS	-	EXPRESSION TAG	UNP O00213
A	509	HIS	-	EXPRESSION TAG	UNP O00213
A	510	HIS	-	EXPRESSION TAG	UNP O00213
A	511	HIS	-	EXPRESSION TAG	UNP O00213
A	512	HIS	-	EXPRESSION TAG	UNP O00213
A	513	HIS	-	EXPRESSION TAG	UNP O00213
B	506	LEU	-	EXPRESSION TAG	UNP O00213
B	507	GLU	-	EXPRESSION TAG	UNP O00213
B	508	HIS	-	EXPRESSION TAG	UNP O00213
B	509	HIS	-	EXPRESSION TAG	UNP O00213
B	510	HIS	-	EXPRESSION TAG	UNP O00213
B	511	HIS	-	EXPRESSION TAG	UNP O00213
B	512	HIS	-	EXPRESSION TAG	UNP O00213
B	513	HIS	-	EXPRESSION TAG	UNP O00213
C	506	LEU	-	EXPRESSION TAG	UNP O00213
C	507	GLU	-	EXPRESSION TAG	UNP O00213
C	508	HIS	-	EXPRESSION TAG	UNP O00213
C	509	HIS	-	EXPRESSION TAG	UNP O00213

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	510	HIS	-	EXPRESSION TAG	UNP O00213
C	511	HIS	-	EXPRESSION TAG	UNP O00213
C	512	HIS	-	EXPRESSION TAG	UNP O00213
C	513	HIS	-	EXPRESSION TAG	UNP O00213
D	506	LEU	-	EXPRESSION TAG	UNP O00213
D	507	GLU	-	EXPRESSION TAG	UNP O00213
D	508	HIS	-	EXPRESSION TAG	UNP O00213
D	509	HIS	-	EXPRESSION TAG	UNP O00213
D	510	HIS	-	EXPRESSION TAG	UNP O00213
D	511	HIS	-	EXPRESSION TAG	UNP O00213
D	512	HIS	-	EXPRESSION TAG	UNP O00213
D	513	HIS	-	EXPRESSION TAG	UNP O00213

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	O	0	0
			4	4		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	5	Total 5	O 5	0	0
3	C	9	Total 9	O 9	0	0
3	D	4	Total 4	O 4	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Amyloid beta A4 precursor protein-binding family B member 1




- Molecule 1: Amyloid beta A4 precursor protein-binding family B member 1



- Molecule 1: Amyloid beta A4 precursor protein-binding family B member 1



- Molecule 1: Amyloid beta A4 precursor protein-binding family B member 1



R463	D464	V468	D471	T474	Q475	N476	L477	K478	C479	C484	E485	A486	P487	A488	K489	N490	H496	E497	S500	K501	I502	N503	A504	E505	L506	E507	HIS	HIS	HIS	HIS	HIS	HIS	HIS
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	146.04Å 146.04Å 78.29Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.21 – 2.70 49.19 – 2.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.21-2.70) 100.0 (49.19-2.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.55 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.242 , 0.313 0.234 , 0.300	Depositor DCC
R_{free} test set	865 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	64.7	Xtriage
Anisotropy	0.025	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.027 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4000	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 81.30 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.6653e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	1/1018 (0.1%)	0.78	0/1371
1	B	0.66	0/1001	0.78	0/1348
1	C	0.72	0/1018	0.77	1/1371 (0.1%)
1	D	0.69	0/990	0.77	0/1333
All	All	0.69	1/4027 (0.0%)	0.78	1/5423 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	479	CYS	CB-SG	-5.09	1.73	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	442	LEU	CA-CB-CG	5.71	128.43	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1003	0	1017	31	0
1	B	986	0	994	29	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1003	0	1017	30	0
1	D	976	0	987	32	0
2	A	5	0	0	0	0
2	C	5	0	0	1	0
3	A	4	0	0	0	0
3	B	5	0	0	1	0
3	C	9	0	0	1	0
3	D	4	0	0	1	0
All	All	4000	0	4015	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 118 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:487:PRO:HG2	1:D:490:ASN:ND2	1.95	0.81
1:B:500:SER:O	1:B:503:MET:HB3	1.80	0.81
1:D:502:ILE:O	1:D:505:GLU:HG3	1.81	0.80
1:A:501:LYS:O	1:A:505:GLU:HB2	1.81	0.80
1:B:398:ILE:O	1:B:402:SER:HB2	1.82	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	123/148 (83%)	108 (88%)	14 (11%)	1 (1%)	21	47
1	B	120/148 (81%)	103 (86%)	14 (12%)	3 (2%)	6	16
1	C	123/148 (83%)	102 (83%)	20 (16%)	1 (1%)	21	47

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	119/148 (80%)	97 (82%)	19 (16%)	3 (2%)	6	16
All	All	485/592 (82%)	410 (84%)	67 (14%)	8 (2%)	11	27

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	367	ILE
1	B	506	LEU
1	D	488	ALA
1	B	441	LEU
1	D	455	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	110/127 (87%)	102 (93%)	8 (7%)	15	36
1	B	107/127 (84%)	98 (92%)	9 (8%)	12	27
1	C	110/127 (87%)	97 (88%)	13 (12%)	6	13
1	D	106/127 (84%)	91 (86%)	15 (14%)	3	9
All	All	433/508 (85%)	388 (90%)	45 (10%)	8	17

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	395	ASN
1	C	461	ARG
1	D	497	GLU
1	C	437	GLN
1	C	462	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	425	GLN
1	C	437	GLN
1	D	437	GLN
1	D	445	GLN
1	D	490	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	1	-	4,4,4	0.71	0	6,6,6	0.54	0
2	PO4	C	2	-	4,4,4	0.85	0	6,6,6	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	1	-	-	0/0/0/0	0/0/0/0
2	PO4	C	2	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	127/148 (85%)	0.11	2 (1%) 72 73	41, 60, 77, 85	0
1	B	126/148 (85%)	0.01	1 (0%) 86 87	38, 60, 80, 83	0
1	C	127/148 (85%)	-0.01	2 (1%) 72 73	37, 57, 76, 79	0
1	D	125/148 (84%)	-0.11	0 100 100	38, 58, 79, 85	0
All	All	505/592 (85%)	0.00	5 (0%) 82 83	37, 59, 78, 85	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	366	GLY	5.2
1	A	367	ILE	3.6
1	C	367	ILE	3.0
1	C	366	GLY	2.4
1	B	378	VAL	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PO4	C	2	5/5	0.90	0.14	89,89,89,90	0
2	PO4	A	1	5/5	0.95	0.11	83,84,84,84	0

6.5 Other polymers [i](#)

There are no such residues in this entry.