



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 12:57 am GMT

PDB ID : 2DB4  
Title : Crystal structure of rotor ring with DCCD of the V- ATPase from *Enterococcus hirae*  
Authors : Murata, T.; Yamato, I.; Kakinuma, Y.; Shirouzu, M.; Walker, J.E.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2005-12-15  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	3481 (2.40-2.40)
Clashscore	122126	3956 (2.40-2.40)
Ramachandran outliers	120053	3897 (2.40-2.40)
Sidechain outliers	120020	3898 (2.40-2.40)
RSRZ outliers	108989	3386 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	156	<div> <div>13%</div> <div>91%</div> <div>8%</div> <div>•</div> </div>
1	B	156	<div> <div>17%</div> <div>92%</div> <div>6%</div> <div>•</div> </div>
1	C	156	<div> <div>13%</div> <div>93%</div> <div>6%</div> <div>•</div> </div>
1	D	156	<div> <div>19%</div> <div>93%</div> <div>6%</div> <div>•</div> </div>
1	E	156	<div> <div>16%</div> <div>93%</div> <div>6%</div> <div>•</div> </div>
1	F	156	<div> <div>13%</div> <div>93%</div> <div>5%</div> <div>•</div> </div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	156	
1	H	156	
1	I	156	
1	J	156	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	LHG	A	204	-	-	-	X
4	LHG	B	204	-	-	-	X
4	LHG	C	204	-	-	-	X
4	LHG	D	203	-	-	-	X
4	LHG	E	203	-	-	-	X
4	LHG	F	204	-	-	-	X
4	LHG	G	203	-	-	-	X
4	LHG	H	203	-	-	-	X
4	LHG	I	204	-	-	-	X
5	UMQ	C	206	X	-	-	-
5	UMQ	F	206	X	-	-	-

## 2 Entry composition


There are 6 unique types of molecules in this entry. The entry contains 12910 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called V-Type Sodium ATPase Subunit K.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	156	Total	C	N	O	S	0	9	0
			1154	760	179	207	8			
1	B	156	Total	C	N	O	S	0	9	0
			1154	760	179	207	8			
1	C	156	Total	C	N	O	S	0	9	0
			1154	760	179	207	8			
1	D	156	Total	C	N	O	S	0	9	0
			1154	760	179	207	8			
1	E	156	Total	C	N	O	S	0	9	0
			1154	760	179	207	8			
1	F	156	Total	C	N	O	S	0	9	0
			1154	760	179	207	8			
1	G	156	Total	C	N	O	S	0	9	0
			1154	760	179	207	8			
1	H	156	Total	C	N	O	S	0	9	0
			1154	760	179	207	8			
1	I	156	Total	C	N	O	S	0	7	0
			1147	757	178	204	8			
1	J	156	Total	C	N	O	S	0	8	0
			1152	759	180	206	7			

- Molecule 2 is DICYCLOHEXYLUREA (three-letter code: DCW) (formula: C<sub>13</sub>H<sub>24</sub>N<sub>2</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			16	13	2	1		
2	B	1	Total	C	N	O	0	0
			16	13	2	1		
2	C	1	Total	C	N	O	0	0
			16	13	2	1		
2	D	1	Total	C	N	O	0	0
			16	13	2	1		
2	E	1	Total	C	N	O	0	0
			16	13	2	1		
2	F	1	Total	C	N	O	0	0
			16	13	2	1		
2	G	1	Total	C	N	O	0	0
			16	13	2	1		
2	H	1	Total	C	N	O	0	0
			16	13	2	1		
2	I	1	Total	C	N	O	0	0
			16	13	2	1		
2	J	1	Total	C	N	O	0	0
			16	13	2	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	1	Total	Na	0	0
			1	1		
3	J	1	Total	Na	0	0
			1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	1	Total	Na	0	0
			1	1		
3	E	1	Total	Na	0	0
			1	1		
3	H	1	Total	Na	0	0
			1	1		
3	B	1	Total	Na	0	0
			1	1		
3	I	1	Total	Na	0	0
			1	1		
3	C	1	Total	Na	0	0
			1	1		
3	A	1	Total	Na	0	0
			1	1		
3	F	1	Total	Na	0	0
			1	1		

- Molecule 4 is 1,2-DIPALMITOYL-PHOSPHATIDYL-GLYCEROLE (three-letter code: LHG) (formula: C<sub>38</sub>H<sub>75</sub>O<sub>10</sub>P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			40	35	5		
4	A	1	Total	C	O	0	0
			40	35	5		
4	B	1	Total	C	O	0	0
			40	35	5		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			40	35	5		
4	C	1	Total	C	O	0	0
			40	35	5		
4	C	1	Total	C	O	0	0
			40	35	5		
4	C	1	Total	C	O	0	0
			40	35	5		
4	D	1	Total	C	O	0	0
			40	35	5		
4	D	1	Total	C	O	0	0
			40	35	5		
4	E	1	Total	C	O	0	0
			40	35	5		
4	F	1	Total	C	O	0	0
			40	35	5		
4	F	1	Total	C	O	0	0
			40	35	5		
4	F	1	Total	C	O	0	0
			40	35	5		
4	G	1	Total	C	O	0	0
			40	35	5		
4	G	1	Total	C	O	0	0
			40	35	5		
4	H	1	Total	C	O	0	0
			40	35	5		
4	I	1	Total	C	O	0	0
			40	35	5		
4	I	1	Total	C	O	0	0
			40	35	5		
4	J	1	Total	C	O	0	0
			40	35	5		
4	J	1	Total	C	O	0	0
			40	35	5		

- Molecule 5 is UNDECYL-MALTOSIDE (three-letter code: UMQ) (formula: C<sub>23</sub>H<sub>44</sub>O<sub>11</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total 12	C 11	O 1	0	0
5	A	1	Total 12	C 11	O 1	0	0
5	C	1	Total 34	C 23	O 11	0	0
5	D	1	Total 12	C 11	O 1	0	0
5	E	1	Total 12	C 11	O 1	0	0
5	F	1	Total 34	C 23	O 11	0	0
5	H	1	Total 12	C 11	O 1	0	0
5	I	1	Total 12	C 11	O 1	0	0
5	J	1	Total 12	C 11	O 1	0	0
5	J	1	Total 12	C 11	O 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	25	Total O 25 25	0	0
6	B	22	Total O 22 22	0	0

*Continued on next page...*



*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	14	Total 14	O 14	0	0
6	D	14	Total 14	O 14	0	0
6	E	17	Total 17	O 17	0	0
6	F	20	Total 20	O 20	0	0
6	G	24	Total 24	O 24	0	0
6	H	32	Total 32	O 32	0	0
6	I	47	Total 47	O 47	0	0
6	J	30	Total 30	O 30	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: V-Type Sodium ATPase Subunit K

Chain A: 





#### • Molecule 1: V-Type Sodium ATPase Subunit K

Chain B: 





#### • Molecule 1: V-Type Sodium ATPase Subunit K

Chain C: 





#### • Molecule 1: V-Type Sodium ATPase Subunit K

Chain D: 




#### • Molecule 1: V-Type Sodium ATPase Subunit K


Chain E: 




#### • Molecule 1: V-Type Sodium ATPase Subunit K




- Molecule 1: V-Type Sodium ATPase Subunit K




- Molecule 1: V-Type Sodium ATPase Subunit K



- Molecule 1: V-Type Sodium ATPase Subunit K



- Molecule 1: V-Type Sodium ATPase Subunit K



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.21Å 125.23Å 211.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	107.83 – 2.40 48.41 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (107.83-2.40) 99.7 (48.41-2.40)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.63 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.225 , 0.231 0.227 , 0.232	Depositor DCC
$R_{free}$ test set	6272 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.3	Xtriage
Anisotropy	0.315	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 63.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	12910	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LHG, NA, UMQ, DCW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	0/1218	0.56	1/1647 (0.1%)
1	B	0.46	0/1218	0.57	1/1647 (0.1%)
1	C	0.44	0/1218	0.57	1/1647 (0.1%)
1	D	0.41	0/1218	0.56	1/1647 (0.1%)
1	E	0.43	0/1218	0.56	1/1647 (0.1%)
1	F	0.46	0/1218	0.57	1/1647 (0.1%)
1	G	0.48	0/1218	0.57	1/1647 (0.1%)
1	H	0.49	0/1218	0.58	1/1647 (0.1%)
1	I	0.55	0/1201	0.58	1/1624 (0.1%)
1	J	0.51	0/1211	0.56	1/1638 (0.1%)
All	All	0.47	0/12156	0.57	10/16438 (0.1%)

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	139	GLU	N-CA-CB	-6.34	99.19	110.60
1	B	139	GLU	N-CA-CB	-6.09	99.64	110.60
1	J	139	GLU	N-CA-CB	-6.02	99.76	110.60
1	C	139	GLU	N-CA-CB	-6.01	99.78	110.60
1	E	139	GLU	N-CA-CB	-5.70	100.35	110.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1154	0	1218	15	0
1	B	1154	0	1218	14	0
1	C	1154	0	1218	14	0
1	D	1154	0	1218	13	0
1	E	1154	0	1218	13	0
1	F	1154	0	1218	15	0
1	G	1154	0	1218	13	0
1	H	1154	0	1218	14	0
1	I	1147	0	1214	13	0
1	J	1152	0	1218	11	0
2	A	16	0	23	0	0
2	B	16	0	23	0	0
2	C	16	0	23	0	0
2	D	16	0	23	0	0
2	E	16	0	23	0	0
2	F	16	0	23	0	0
2	G	16	0	23	0	0
2	H	16	0	23	0	0
2	I	16	0	23	0	0
2	J	16	0	23	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
3	I	1	0	0	0	0
3	J	1	0	0	0	0
4	A	80	0	134	7	0
4	B	80	0	134	6	0
4	C	120	0	201	6	0
4	D	80	0	134	6	0
4	E	40	0	67	6	0
4	F	120	0	201	6	0
4	G	80	0	134	7	0
4	H	40	0	67	5	0
4	I	80	0	134	4	0
4	J	80	0	134	4	0
5	A	24	0	46	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	34	0	41	1	0
5	D	12	0	23	0	0
5	E	12	0	23	0	0
5	F	34	0	40	2	0
5	H	12	0	23	0	0
5	I	12	0	23	0	0
5	J	24	0	46	1	0
6	A	25	0	0	0	0
6	B	22	0	0	0	0
6	C	14	0	0	0	0
6	D	14	0	0	0	0
6	E	17	0	0	0	0
6	F	20	0	0	1	0
6	G	24	0	0	0	0
6	H	32	0	0	0	0
6	I	47	0	0	0	0
6	J	30	0	0	0	0
All	All	12910	0	14011	134	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 134 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:54[B]:GLN:HE22	5:C:206:UMQ:HA1	1.46	0.80
1:G:2:MET:HG2	1:H:1:MET:HB2	1.73	0.70
1:F:2:MET:HG2	1:G:1:MET:HB2	1.80	0.64
1:A:2:MET:HG2	1:B:1:MET:HB2	1.79	0.64
1:C:2:MET:HG2	1:D:1:MET:HB2	1.81	0.62

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	163/156 (104%)	161 (99%)	2 (1%)	0	100	100
1	B	163/156 (104%)	161 (99%)	2 (1%)	0	100	100
1	C	163/156 (104%)	160 (98%)	3 (2%)	0	100	100
1	D	163/156 (104%)	161 (99%)	2 (1%)	0	100	100
1	E	163/156 (104%)	161 (99%)	2 (1%)	0	100	100
1	F	163/156 (104%)	161 (99%)	2 (1%)	0	100	100
1	G	163/156 (104%)	161 (99%)	2 (1%)	0	100	100
1	H	163/156 (104%)	161 (99%)	2 (1%)	0	100	100
1	I	161/156 (103%)	160 (99%)	1 (1%)	0	100	100
1	J	162/156 (104%)	161 (99%)	1 (1%)	0	100	100
All	All	1627/1560 (104%)	1608 (99%)	19 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	122/113 (108%)	119 (98%)	3 (2%)	50	71
1	B	122/113 (108%)	119 (98%)	3 (2%)	50	71
1	C	122/113 (108%)	119 (98%)	3 (2%)	50	71
1	D	122/113 (108%)	119 (98%)	3 (2%)	50	71
1	E	122/113 (108%)	119 (98%)	3 (2%)	50	71
1	F	122/113 (108%)	119 (98%)	3 (2%)	50	71
1	G	122/113 (108%)	119 (98%)	3 (2%)	50	71
1	H	122/113 (108%)	119 (98%)	3 (2%)	50	71
1	I	120/113 (106%)	117 (98%)	3 (2%)	50	71

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	J	121/113 (107%)	118 (98%)	3 (2%)	50	71
All	All	1217/1130 (108%)	1187 (98%)	30 (2%)	60	71

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	54[B]	GLN
1	F	54[B]	GLN
1	J	54[A]	GLN
1	F	54[A]	GLN
1	F	61	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	I	9	ASN
1	J	8	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 50 ligands modelled in this entry, 10 are monoatomic - leaving 40 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	DCW	A	201	1	17,17,17	1.46	2 (11%)	21,21,21	1.55	3 (14%)
4	LHG	A	203	-	39,39,48	1.07	2 (5%)	41,41,54	1.14	3 (7%)
4	LHG	A	204	-	39,39,48	1.04	2 (5%)	41,41,54	1.07	2 (4%)
5	UMQ	A	205	-	11,11,35	0.91	0	10,10,46	0.82	0
5	UMQ	A	206	-	11,11,35	0.29	0	10,10,46	0.47	0
2	DCW	B	201	1	17,17,17	1.48	2 (11%)	21,21,21	1.56	3 (14%)
4	LHG	B	203	-	39,39,48	1.08	2 (5%)	41,41,54	1.14	3 (7%)
4	LHG	B	204	-	39,39,48	1.04	2 (5%)	41,41,54	1.07	2 (4%)
2	DCW	C	201	1	17,17,17	1.49	2 (11%)	21,21,21	1.53	3 (14%)
4	LHG	C	203	-	39,39,48	1.09	2 (5%)	41,41,54	1.12	2 (4%)
4	LHG	C	204	-	39,39,48	1.03	2 (5%)	41,41,54	1.06	2 (4%)
4	LHG	C	205	-	39,39,48	1.07	2 (5%)	41,41,54	1.12	3 (7%)
5	UMQ	C	206	-	35,35,35	1.06	2 (5%)	46,46,46	1.64	6 (13%)
2	DCW	D	201	1	17,17,17	1.47	2 (11%)	21,21,21	1.55	3 (14%)
4	LHG	D	203	-	39,39,48	1.04	2 (5%)	41,41,54	1.07	3 (7%)
4	LHG	D	204	-	39,39,48	1.08	2 (5%)	41,41,54	1.13	3 (7%)
5	UMQ	D	205	-	11,11,35	0.79	1 (9%)	10,10,46	0.75	0
2	DCW	E	201	1	17,17,17	1.49	2 (11%)	21,21,21	1.59	3 (14%)
4	LHG	E	203	-	39,39,48	1.04	2 (5%)	41,41,54	1.07	2 (4%)
5	UMQ	E	204	-	11,11,35	0.25	0	10,10,46	0.54	0
2	DCW	F	201	1	17,17,17	1.46	2 (11%)	21,21,21	1.52	4 (19%)
4	LHG	F	203	-	39,39,48	1.07	2 (5%)	41,41,54	1.11	3 (7%)
4	LHG	F	204	-	39,39,48	1.03	2 (5%)	41,41,54	1.07	2 (4%)
4	LHG	F	205	-	39,39,48	1.07	2 (5%)	41,41,54	1.13	3 (7%)
5	UMQ	F	206	-	35,35,35	1.10	3 (8%)	46,46,46	1.64	6 (13%)
2	DCW	G	201	1	17,17,17	1.43	2 (11%)	21,21,21	1.43	3 (14%)
4	LHG	G	203	-	39,39,48	1.03	2 (5%)	41,41,54	1.07	3 (7%)
4	LHG	G	204	-	39,39,48	1.07	2 (5%)	41,41,54	1.15	3 (7%)
2	DCW	H	201	1	17,17,17	1.43	2 (11%)	21,21,21	1.51	3 (14%)
4	LHG	H	203	-	39,39,48	1.04	2 (5%)	41,41,54	1.08	3 (7%)
5	UMQ	H	204	-	11,11,35	0.72	0	10,10,46	0.70	0
2	DCW	I	201	1	17,17,17	1.55	2 (11%)	21,21,21	1.56	3 (14%)
4	LHG	I	203	-	39,39,48	1.07	2 (5%)	41,41,54	1.13	3 (7%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	LHG	I	204	-	39,39,48	1.04	2 (5%)	41,41,54	1.07	2 (4%)
5	UMQ	I	205	-	11,11,35	0.27	0	10,10,46	0.53	0
2	DCW	J	201	1	17,17,17	1.48	2 (11%)	21,21,21	1.51	4 (19%)
4	LHG	J	203	-	39,39,48	1.07	2 (5%)	41,41,54	1.13	3 (7%)
4	LHG	J	204	-	39,39,48	1.04	2 (5%)	41,41,54	1.03	2 (4%)
5	UMQ	J	205	-	11,11,35	0.25	0	10,10,46	0.48	0
5	UMQ	J	206	-	11,11,35	1.17	1 (9%)	10,10,46	1.14	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DCW	A	201	1	-	0/8/24/24	0/2/2/2
4	LHG	A	203	-	-	0/41/41/53	0/0/0/0
4	LHG	A	204	-	-	0/41/41/53	0/0/0/0
5	UMQ	A	205	-	-	0/9/9/60	0/0/0/2
5	UMQ	A	206	-	-	0/9/9/60	0/0/0/2
2	DCW	B	201	1	-	0/8/24/24	0/2/2/2
4	LHG	B	203	-	-	0/41/41/53	0/0/0/0
4	LHG	B	204	-	-	0/41/41/53	0/0/0/0
2	DCW	C	201	1	-	0/8/24/24	0/2/2/2
4	LHG	C	203	-	-	0/41/41/53	0/0/0/0
4	LHG	C	204	-	-	0/41/41/53	0/0/0/0
4	LHG	C	205	-	-	0/41/41/53	0/0/0/0
5	UMQ	C	206	-	1/1/10/10	0/20/60/60	0/2/2/2
2	DCW	D	201	1	-	0/8/24/24	0/2/2/2
4	LHG	D	203	-	-	0/41/41/53	0/0/0/0
4	LHG	D	204	-	-	0/41/41/53	0/0/0/0
5	UMQ	D	205	-	-	0/9/9/60	0/0/0/2
2	DCW	E	201	1	-	0/8/24/24	0/2/2/2
4	LHG	E	203	-	-	0/41/41/53	0/0/0/0
5	UMQ	E	204	-	-	0/9/9/60	0/0/0/2
2	DCW	F	201	1	-	0/8/24/24	0/2/2/2
4	LHG	F	203	-	-	0/41/41/53	0/0/0/0
4	LHG	F	204	-	-	0/41/41/53	0/0/0/0
4	LHG	F	205	-	-	0/41/41/53	0/0/0/0
5	UMQ	F	206	-	1/1/10/10	0/20/60/60	0/2/2/2
2	DCW	G	201	1	-	0/8/24/24	0/2/2/2
4	LHG	G	203	-	-	0/41/41/53	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	LHG	G	204	-	-	0/41/41/53	0/0/0/0
2	DCW	H	201	1	-	0/8/24/24	0/2/2/2
4	LHG	H	203	-	-	0/41/41/53	0/0/0/0
5	UMQ	H	204	-	-	0/9/9/60	0/0/0/2
2	DCW	I	201	1	-	0/8/24/24	0/2/2/2
4	LHG	I	203	-	-	0/41/41/53	0/0/0/0
4	LHG	I	204	-	-	0/41/41/53	0/0/0/0
5	UMQ	I	205	-	-	0/9/9/60	0/0/0/2
2	DCW	J	201	1	-	0/8/24/24	0/2/2/2
4	LHG	J	203	-	-	0/41/41/53	0/0/0/0
4	LHG	J	204	-	-	0/41/41/53	0/0/0/0
5	UMQ	J	205	-	-	0/9/9/60	0/0/0/2
5	UMQ	J	206	-	-	0/9/9/60	0/0/0/2

The worst 5 of 67 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	206	UMQ	O3-C3	-4.15	1.32	1.43
5	C	206	UMQ	O3-C3	-4.13	1.32	1.43
5	F	206	UMQ	O2-C2	-3.92	1.33	1.43
5	C	206	UMQ	O2-C2	-3.79	1.33	1.43
5	D	205	UMQ	CL-CK	2.03	1.66	1.49

The worst 5 of 98 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	201	DCW	C2-N1-C1	-4.07	115.42	122.93
2	E	201	DCW	C2-N1-C1	-4.06	115.43	122.93
2	C	201	DCW	C2-N1-C1	-4.04	115.46	122.93
2	B	201	DCW	C2-N1-C1	-4.04	115.46	122.93
2	F	201	DCW	C2-N1-C1	-4.01	115.53	122.93

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	F	206	UMQ	C2
5	C	206	UMQ	C3

There are no torsion outliers.

There are no ring outliers.

15 monomers are involved in 48 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	204	LHG	7	0
5	A	205	UMQ	1	0
4	B	204	LHG	6	0
4	C	204	LHG	6	0
5	C	206	UMQ	1	0
4	D	203	LHG	6	0
4	E	203	LHG	6	0
4	F	204	LHG	6	0
5	F	206	UMQ	2	0
4	G	203	LHG	7	0
4	H	203	LHG	5	0
4	I	204	LHG	4	0
4	J	204	LHG	4	0
5	J	205	UMQ	1	0
5	J	206	UMQ	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	156/156 (100%)	0.74	21 (13%) 3 2	46, 52, 69, 81	9 (5%)
1	B	156/156 (100%)	0.87	26 (16%) 1 1	46, 52, 69, 81	9 (5%)
1	C	156/156 (100%)	0.45	20 (12%) 3 3	46, 52, 69, 81	9 (5%)
1	D	156/156 (100%)	0.90	29 (18%) 1 1	46, 52, 69, 81	9 (5%)
1	E	156/156 (100%)	0.79	25 (16%) 1 1	46, 52, 69, 81	9 (5%)
1	F	156/156 (100%)	0.72	21 (13%) 3 2	46, 52, 69, 81	9 (5%)
1	G	156/156 (100%)	0.92	23 (14%) 2 2	46, 52, 69, 81	9 (5%)
1	H	156/156 (100%)	0.58	24 (15%) 2 1	46, 52, 69, 81	9 (5%)
1	I	156/156 (100%)	0.23	14 (8%) 9 8	46, 52, 69, 81	7 (4%)
1	J	156/156 (100%)	-0.03	12 (7%) 13 12	46, 51, 60, 72	8 (5%)
All	All	1560/1560 (100%)	0.62	215 (13%) 3 2	46, 52, 69, 81	87 (5%)

The worst 5 of 215 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	83[A]	ASP	11.2
1	B	82	SER	10.9
1	D	81	GLY	10.8
1	G	82	SER	10.5
1	G	81	GLY	10.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	LHG	B	204	40/49	0.15	0.52	135,142,146,146	0
4	LHG	A	204	40/49	0.19	0.52	135,142,146,146	0
4	LHG	H	203	40/49	0.28	0.51	135,142,146,146	0
4	LHG	G	203	40/49	0.30	0.51	135,142,146,146	0
4	LHG	I	204	40/49	0.31	0.51	135,142,146,146	0
4	LHG	J	204	40/49	0.39	0.37	135,142,146,146	0
4	LHG	C	204	40/49	0.40	0.49	135,142,146,146	0
4	LHG	E	203	40/49	0.50	0.43	135,142,146,146	0
4	LHG	F	204	40/49	0.53	0.49	135,142,146,146	0
4	LHG	D	203	40/49	0.54	0.45	135,142,146,146	0
5	UMQ	H	204	12/34	0.56	0.28	87,87,89,89	0
4	LHG	A	203	40/49	0.57	0.33	97,110,121,121	0
5	UMQ	A	205	12/34	0.65	0.34	94,96,97,97	0
5	UMQ	D	205	12/34	0.67	0.31	98,100,102,102	0
4	LHG	D	204	40/49	0.68	0.35	97,110,121,121	0
5	UMQ	A	206	12/34	0.69	0.26	93,94,95,95	0
4	LHG	F	205	40/49	0.69	0.33	97,110,121,121	0
4	LHG	G	204	40/49	0.70	0.30	97,110,121,121	0
4	LHG	C	205	40/49	0.70	0.30	97,110,121,121	0
5	UMQ	E	204	12/34	0.71	0.39	95,97,98,98	0
4	LHG	F	203	40/49	0.72	0.34	97,110,121,121	0
4	LHG	I	203	40/49	0.72	0.23	97,110,121,121	0
5	UMQ	J	205	12/34	0.74	0.21	93,94,95,95	0
4	LHG	B	203	40/49	0.74	0.30	97,110,121,121	0
4	LHG	C	203	40/49	0.74	0.34	97,110,121,121	0
4	LHG	J	203	40/49	0.76	0.26	97,110,121,121	0
5	UMQ	J	206	12/34	0.81	0.21	89,90,91,91	0
5	UMQ	F	206	34/34	0.84	0.27	80,87,93,94	0
3	NA	H	202	1/1	0.85	0.10	62,62,62,62	0
3	NA	D	202	1/1	0.85	0.28	62,62,62,62	0
5	UMQ	C	206	34/34	0.85	0.39	90,93,96,96	0
2	DCW	D	201	16/16	0.88	0.29	64,65,69,69	0

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NA	B	202	1/1	0.89	0.33	62,62,62,62	0
2	DCW	A	201	16/16	0.89	0.36	64,65,69,69	0
5	UMQ	I	205	12/34	0.90	0.13	77,78,79,80	0
2	DCW	E	201	16/16	0.90	0.27	64,65,69,69	0
3	NA	E	202	1/1	0.90	0.37	62,62,62,62	0
3	NA	G	202	1/1	0.90	0.33	62,62,62,62	0
2	DCW	G	201	16/16	0.90	0.19	64,65,69,69	0
2	DCW	H	201	16/16	0.90	0.15	63,65,69,69	0
2	DCW	J	201	16/16	0.92	0.30	64,65,69,69	0
2	DCW	C	201	16/16	0.92	0.32	64,65,69,69	0
3	NA	I	202	1/1	0.92	0.23	63,63,63,63	0
2	DCW	F	201	16/16	0.93	0.19	64,65,69,69	0
3	NA	A	202	1/1	0.93	0.13	62,62,62,62	0
2	DCW	B	201	16/16	0.94	0.28	64,65,69,69	0
3	NA	F	202	1/1	0.94	0.36	62,62,62,62	0
3	NA	C	202	1/1	0.94	0.54	62,62,62,62	0
2	DCW	I	201	16/16	0.95	0.17	64,65,69,69	0
3	NA	J	202	1/1	0.95	0.07	61,61,61,61	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.