



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 06:31 pm GMT

PDB ID : 4DBR
Title : Myosin VI D179Y (MD) pre-powerstroke state
Authors : Pylypenko, O.; Sweeney, H.L.; Houdusse, A.
Deposited on : 2012-01-16
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2220 (1.96-1.96)
Clashscore	122126	2333 (1.96-1.96)
Ramachandran outliers	120053	2314 (1.96-1.96)
Sidechain outliers	120020	2314 (1.96-1.96)
RSRZ outliers	108989	2174 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	786	<div> <div>3%</div> <div>89%</div> <div>6%</div> <div>.</div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6862 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myosin-VI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	754	6070	3878	1037	1125	30	0	14	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	4	ALA	-	EXPRESSION TAG	UNP F1RQI7
A	179	TYR	ASP	ENGINEERED MUTATION	UNP F1RQI7

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is VANADATE ION (three-letter code: VO4) (formula: O_4V).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	V	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	727	Total	O	0	0
			727	727		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Myosin-VI



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	91.94Å 93.53Å 102.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.36 – 1.95 29.36 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.36-1.95) 99.6 (29.36-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 1.91Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.2_869)	Depositor
R, R_{free}	0.174 , 0.216 0.170 , 0.212	Depositor DCC
R_{free} test set	3474 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	30.1	Xtriage
Anisotropy	0.508	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6862	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: VO4, MG, EDO, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/6234	0.58	0/8415

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6070	0	5954	33	0
2	A	32	0	48	3	0
3	A	27	0	12	0	0
4	A	1	0	0	0	0
5	A	5	0	0	0	0
6	A	727	0	0	8	0
All	All	6862	0	6014	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:508:TYR:H	2:A:808:EDO:H11	1.43	0.80
1:A:533:ASN:ND2	1:A:597:ASN:HD22	1.99	0.59
1:A:326:LYS:NZ	6:A:1560:HOH:O	2.20	0.59
1:A:78:LYS:NZ	6:A:1563:HOH:O	2.35	0.58
1:A:227:HIS:O	2:A:807:EDO:H22	2.06	0.56
1:A:180:ASP:O	1:A:184:GLU:HB3	2.07	0.54
1:A:738:LEU:HD11	1:A:780:LEU:HD23	1.90	0.52
1:A:655:LEU:HD23	1:A:658:LEU:HD12	1.92	0.51
1:A:712[A]:HIS:CE1	1:A:716:ASN:HD21	2.31	0.49
1:A:515:ILE:O	1:A:519:GLU:HB2	2.13	0.48
1:A:169[B]:THR:HG21	1:A:179:TYR:HA	1.96	0.47
1:A:561:ARG:HA	1:A:569:ARG:HA	1.97	0.47
1:A:742:LEU:HD11	1:A:780:LEU:HD21	1.98	0.46
1:A:384:ASP:OD1	1:A:387:ARG:NH1	2.42	0.45
1:A:85:ARG:HA	2:A:802:EDO:H12	1.99	0.45
1:A:186[B]:ASN:OD1	6:A:1213:HOH:O	2.21	0.45
1:A:259:ILE:HD12	6:A:1295:HOH:O	2.16	0.45
1:A:722:MET:HG2	1:A:726:LEU:HD23	1.99	0.45
1:A:721:TYR:O	1:A:781:VAL:HG11	2.16	0.44
1:A:208:LYS:HE2	6:A:1299:HOH:O	2.18	0.43
1:A:314:HIS:CE1	1:A:318:ILE:HD11	2.53	0.43
1:A:208:LYS:NZ	6:A:1311:HOH:O	2.46	0.42
1:A:162:LYS:O	1:A:166:ARG:HG3	2.19	0.42
1:A:18:VAL:HG12	1:A:34:LEU:HB2	2.02	0.42
1:A:162:LYS:HD3	1:A:186[B]:ASN:ND2	2.35	0.42
1:A:54:ASP:HB3	1:A:57:LYS:HG2	2.01	0.42
1:A:726:LEU:HB2	1:A:785:ASN:OD1	2.20	0.41
1:A:720:LYS:HE3	1:A:720:LYS:HB2	1.80	0.41
1:A:135:PHE:HA	1:A:452:ILE:HD11	2.01	0.41
1:A:491:GLU:OE2	6:A:1317:HOH:O	2.22	0.41
1:A:229:LEU:HA	1:A:229:LEU:HD12	1.94	0.40
1:A:426[A]:THR:HG21	1:A:601:LEU:HD22	2.04	0.40
1:A:702:GLN:NE2	6:A:1286:HOH:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	760/786 (97%)	747 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	659/694 (95%)	654 (99%)	5 (1%)	83	82

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	276	ARG
1	A	377	GLU
1	A	387	ARG
1	A	539	SER
1	A	708	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	533	ASN
1	A	549	GLN
1	A	645	ASN
1	A	702	GLN
1	A	716	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	801	-	3,3,3	0.49	0	2,2,2	0.30	0
2	EDO	A	802	-	3,3,3	0.56	0	2,2,2	0.34	0
2	EDO	A	803	-	3,3,3	0.59	0	2,2,2	0.30	0
2	EDO	A	804	-	3,3,3	0.58	0	2,2,2	0.59	0
2	EDO	A	805	-	3,3,3	0.59	0	2,2,2	0.12	0
2	EDO	A	806	-	3,3,3	0.45	0	2,2,2	0.32	0
2	EDO	A	807	-	3,3,3	0.51	0	2,2,2	0.52	0
2	EDO	A	808	-	3,3,3	0.37	0	2,2,2	0.66	0
3	ADP	A	809	5,4	25,29,29	1.06	1 (4%)	25,45,45	1.83	6 (24%)
5	VO4	A	811	3,4	1,4,4	5.51	1 (100%)	0,6,6	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	801	-	-	0/1/1/1	0/0/0/0
2	EDO	A	802	-	-	0/1/1/1	0/0/0/0
2	EDO	A	803	-	-	0/1/1/1	0/0/0/0
2	EDO	A	804	-	-	0/1/1/1	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	805	-	-	0/1/1/1	0/0/0/0
2	EDO	A	806	-	-	0/1/1/1	0/0/0/0
2	EDO	A	807	-	-	0/1/1/1	0/0/0/0
2	EDO	A	808	-	-	0/1/1/1	0/0/0/0
3	ADP	A	809	5,4	-	0/12/32/32	0/3/3/3
5	VO4	A	811	3,4	-	0/0/0/0	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	809	ADP	C5-C4	3.39	1.48	1.40
5	A	811	VO4	O1-V	5.51	1.95	1.63

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	809	ADP	N3-C2-N1	-6.54	123.27	128.86
3	A	809	ADP	C4-C5-N7	-2.37	107.12	109.41
3	A	809	ADP	PA-O3A-PB	-2.29	124.92	132.63
3	A	809	ADP	C1'-N9-C4	-2.23	122.78	126.64
3	A	809	ADP	O2B-PB-O1B	2.22	119.25	110.60
3	A	809	ADP	C2-N1-C6	2.23	122.54	118.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	802	EDO	1	0
2	A	807	EDO	1	0
2	A	808	EDO	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	754/786 (95%)	0.04	26 (3%) 45 55	8, 22, 49, 75	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	603	MET	4.5
1	A	393	ARG	4.0
1	A	407	ILE	4.0
1	A	409	VAL	4.0
1	A	4	ALA	3.8
1	A	394	VAL	3.8
1	A	176	GLN	3.7
1	A	27	ASP	3.1
1	A	178[A]	ILE	2.9
1	A	36	GLN	2.9
1	A	177	ASP	2.9
1	A	26	PRO	2.7
1	A	561	ARG	2.7
1	A	247	ILE	2.6
1	A	366	ASN	2.5
1	A	410	PRO	2.5
1	A	784	VAL	2.4
1	A	450	TYR	2.4
1	A	408	LYS	2.4
1	A	747	ILE	2.3
1	A	35	ASN	2.3
1	A	362	CYS	2.3
1	A	175	GLY	2.2
1	A	602	HIS	2.1
1	A	789	ILE	2.1
1	A	42	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	EDO	A	807	4/4	0.93	0.20	22,37,41,42	0
2	EDO	A	808	4/4	0.93	0.25	29,35,36,38	0
2	EDO	A	803	4/4	0.95	0.11	17,18,22,23	0
2	EDO	A	802	4/4	0.95	0.14	18,26,28,30	0
2	EDO	A	805	4/4	0.96	0.16	14,25,26,31	0
2	EDO	A	804	4/4	0.96	0.10	22,24,26,27	0
2	EDO	A	806	4/4	0.96	0.14	25,25,31,35	0
2	EDO	A	801	4/4	0.98	0.07	13,14,14,20	0
4	MG	A	810	1/1	0.99	0.20	14,14,14,14	0
3	ADP	A	809	27/27	0.99	0.11	7,10,14,16	0
5	VO4	A	811	5/5	0.99	0.13	7,9,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.