



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 01:00 am GMT

PDB ID : 4DDS  
Title : CTX-M-9 class A beta-lactamase complexed with compound 11  
Authors : Nichols, D.A.; Chen, Y.  
Deposited on : 2012-01-19  
Resolution : 1.36 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1197 (1.38-1.34)
Clashscore	122126	1232 (1.38-1.34)
Ramachandran outliers	120053	1215 (1.38-1.34)
Sidechain outliers	120020	1215 (1.38-1.34)
RSRZ outliers	108989	1177 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	<div> <div>6%</div> <div> <div></div> <div>90%</div> <div>8%</div> <div>..</div> </div> </div>
1	B	263	<div> <div>8%</div> <div> <div></div> <div>90%</div> <div>9%</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	0J7	A	303	-	-	X	-
2	0J7	B	302	-	-	X	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4617 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	9	0
			2007	1244	360	396	7			
1	B	261	Total	C	N	O	S	0	6	0
			1983	1229	355	391	8			

- Molecule 2 is 3-(pyrimidin-2-yl)-N-[3-(1H-tetrazol-5-yl)phenyl]benzamide (three-letter code: OJ7) (formula: C<sub>18</sub>H<sub>13</sub>N<sub>7</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			26	18	7	1		
2	A	1	Total	C	N	O	0	0
			26	18	7	1		
2	A	1	Total	C	N	O	0	0
			26	18	7	1		
2	B	1	Total	C	N	O	0	0
			26	18	7	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			26	18	7	1		

- Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C<sub>2</sub>H<sub>6</sub>OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	269	Total	O	0	0
			269	269		
4	B	224	Total	O	0	0
			224	224		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-lactamase



- Molecule 1: Beta-lactamase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.24Å 106.94Å 47.68Å 90.00° 100.97° 90.00°	Depositor
Resolution (Å)	25.26 – 1.36 25.26 – 1.36	Depositor EDS
% Data completeness (in resolution range)	93.0 (25.26-1.36) 93.0 (25.26-1.36)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.64 (at 1.36Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.156 , 0.199 0.151 , 0.195	Depositor DCC
$R_{free}$ test set	4395 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	6.6	Xtriage
Anisotropy	0.835	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 52.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4617	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 0J7, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.54	0/2039	0.67	0/2774
1	B	0.52	0/2018	0.68	1/2744 (0.0%)
All	All	0.53	0/4057	0.67	1/5518 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	212	LYS	CD-CE-NZ	-5.18	99.78	111.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2007	0	2019	24	0
1	B	1983	0	2001	30	0
2	A	78	0	39	9	0
2	B	52	0	26	17	0
3	B	4	0	6	0	0
4	A	269	0	0	4	0
4	B	224	0	0	9	0
All	All	4617	0	4091	60	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:GLN:HG3	4:A:665:HOH:O	1.37	1.22
1:B:105:TYR:CD2	2:B:302:OJ7:H4	1.86	1.10
1:A:105:TYR:CD1	2:A:303:OJ7:H4	1.89	1.08
1:A:105:TYR:CG	2:A:303:OJ7:H4	1.92	1.05
2:B:302:OJ7:O26	2:B:302:OJ7:H3	1.59	1.00
1:B:105:TYR:CG	2:B:302:OJ7:H4	1.97	1.00
1:B:105:TYR:CE2	2:B:302:OJ7:H4	2.05	0.91
1:A:44[B]:LEU:HD11	1:A:46:VAL:HG23	1.54	0.87
1:A:44[B]:LEU:CD1	1:A:46:VAL:HG23	2.05	0.87
2:A:303:OJ7:O26	2:A:303:OJ7:H3	1.74	0.85
1:A:44[B]:LEU:HD11	1:A:46:VAL:CG2	2.08	0.83
1:A:44[B]:LEU:HD13	1:A:45:GLY:N	1.97	0.78
1:A:167:PRO:HG2	2:A:301:OJ7:H13	1.70	0.72
1:B:75[B]:MET:HE2	1:B:75[B]:MET:HA	1.69	0.72
1:A:105:TYR:CE1	2:A:303:OJ7:H4	2.24	0.71
1:A:44[B]:LEU:C	1:A:44[B]:LEU:HD13	2.12	0.69
1:B:105:TYR:CD1	2:B:302:OJ7:H4	2.29	0.67
1:B:105:TYR:CG	2:B:302:OJ7:C10	2.78	0.66
2:B:302:OJ7:O26	2:B:302:OJ7:C9	2.34	0.66
1:A:105:TYR:CD2	2:A:303:OJ7:H4	2.31	0.65
2:B:302:OJ7:C18	4:B:623:HOH:O	2.45	0.65
1:B:201:GLU:HG3	4:B:598:HOH:O	1.97	0.63
2:B:302:OJ7:H8	4:B:623:HOH:O	1.98	0.62
1:B:105:TYR:CZ	2:B:302:OJ7:H4	2.35	0.62
1:B:27:SER:O	1:B:30:GLN:HB2	2.00	0.61
1:A:273:GLU:CD	1:A:273:GLU:H	2.04	0.59
1:A:240[A]:ASP:OD1	1:A:270[A]:GLN:O	2.21	0.58
1:A:186[A]:MET:HE1	4:A:537:HOH:O	2.04	0.56
1:A:44[B]:LEU:C	1:A:44[B]:LEU:CD1	2.74	0.55
1:A:102:LEU:HD11	1:A:113[A]:VAL:HG11	1.88	0.54
1:B:186[A]:MET:CE	4:B:535:HOH:O	2.55	0.54
1:B:105:TYR:CD1	2:B:302:OJ7:C10	2.92	0.53
1:B:212:LYS:HZ3	1:B:212:LYS:CB	2.22	0.52
1:B:105:TYR:CE1	2:B:302:OJ7:H4	2.46	0.51
1:A:56:GLN:NE2	4:A:570:HOH:O	2.44	0.51
1:B:167:PRO:HG2	2:B:301:OJ7:H13	1.94	0.50
1:B:252:PRO:HB2	1:B:256:ARG:HG2	1.93	0.50

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:27:SER:HB3	1:B:30:GLN:CG	2.43	0.49
2:A:303:OJ7:O26	2:A:303:OJ7:C9	2.46	0.49
1:A:105:TYR:CG	2:A:303:OJ7:C10	2.81	0.48
1:B:27:SER:HB3	1:B:30:GLN:HG2	1.94	0.48
1:B:88:LYS:NZ	4:B:558:HOH:O	2.46	0.48
1:B:111:LYS:HE2	4:B:620:HOH:O	2.13	0.47
1:B:186[A]:MET:HE3	4:B:535:HOH:O	2.15	0.47
1:B:89:GLN:NE2	4:B:604:HOH:O	2.48	0.46
1:A:240[A]:ASP:OD1	1:A:270[A]:GLN:CG	2.64	0.45
1:A:240[A]:ASP:OD1	1:A:270[A]:GLN:HG2	2.17	0.44
1:B:111:LYS:HE3	4:B:599:HOH:O	2.17	0.44
1:B:105:TYR:CE2	2:B:302:OJ7:C10	2.91	0.43
1:A:96:GLU:OE1	4:A:655:HOH:O	2.21	0.43
1:B:28:ALA:HA	1:B:31:GLN:HE21	1.83	0.42
1:A:186[A]:MET:HA	1:A:189[A]:THR:HG22	2.01	0.42
1:B:259:LEU:HD11	1:B:290:LEU:HD11	2.02	0.42
1:B:75[B]:MET:CA	1:B:75[B]:MET:HE2	2.42	0.41
1:B:105:TYR:CZ	2:B:302:OJ7:C10	3.03	0.41
1:B:105:TYR:CE1	2:B:302:OJ7:C10	3.04	0.41
1:A:91:LEU:HB3	1:A:120:ALA:HB2	2.03	0.41
1:A:105:TYR:CD2	2:A:303:OJ7:C10	3.02	0.41
1:B:91:LEU:HB3	1:B:120:ALA:HB2	2.03	0.41
1:B:105:TYR:CE1	2:B:302:OJ7:H5	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/263 (102%)	258 (97%)	8 (3%)	1 (0%)	36	13
1	B	265/263 (101%)	259 (98%)	5 (2%)	1 (0%)	36	13

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	532/526 (101%)	517 (97%)	13 (2%)	2 (0%)	36 13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	103	VAL
1	B	103	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	211/204 (103%)	202 (96%)	9 (4%)	32 4
1	B	209/204 (102%)	203 (97%)	6 (3%)	45 11
All	All	420/408 (103%)	405 (96%)	15 (4%)	44 7

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	106	ASN
1	A	130	SER
1	A	240[A]	ASP
1	A	240[B]	ASP
1	A	254	GLN
1	A	269	GLN
1	A	273	GLU
1	A	274[A]	SER
1	A	274[B]	SER
1	B	38	LYS
1	B	130	SER
1	B	212	LYS
1	B	240	ASP
1	B	274[A]	SER
1	B	274[B]	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	GLN
1	A	93	GLN
1	A	254	GLN
1	A	269	GLN
1	B	31	GLN
1	B	89	GLN
1	B	112	HIS
1	B	197	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	OJ7	A	301	-	29,29,29	1.61	5 (17%)	39,39,39	2.17	14 (35%)
2	OJ7	A	302	-	29,29,29	1.68	3 (10%)	39,39,39	1.90	11 (28%)
2	OJ7	A	303	-	29,29,29	1.71	3 (10%)	39,39,39	2.43	15 (38%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	0J7	B	301	-	29,29,29	1.49	3 (10%)	39,39,39	2.24	10 (25%)
2	0J7	B	302	-	29,29,29	1.69	3 (10%)	39,39,39	2.32	15 (38%)
3	DMS	B	303	-	3,3,3	2.42	1 (33%)	3,3,3	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0J7	A	301	-	-	0/16/16/16	0/4/4/4
2	0J7	A	302	-	-	0/16/16/16	0/4/4/4
2	0J7	A	303	-	-	0/16/16/16	0/4/4/4
2	0J7	B	301	-	-	0/16/16/16	0/4/4/4
2	0J7	B	302	-	-	0/16/16/16	0/4/4/4
3	DMS	B	303	-	-	0/0/0/0	0/0/0/0

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	0J7	C5-N4	2.11	1.36	1.33
2	A	301	0J7	C5-N1	2.54	1.37	1.33
2	A	301	0J7	N1-N2	2.56	1.38	1.34
2	B	301	0J7	N1-N2	3.31	1.40	1.34
2	A	302	0J7	N1-N2	3.59	1.40	1.34
2	A	301	0J7	N4-N3	3.84	1.40	1.34
2	A	302	0J7	N4-N3	3.88	1.40	1.34
2	B	301	0J7	N4-N3	4.01	1.41	1.34
3	B	303	DMS	O-S	4.03	1.77	1.50
2	B	302	0J7	N4-N3	4.04	1.41	1.34
2	B	302	0J7	N1-N2	4.06	1.41	1.34
2	A	303	0J7	N1-N2	4.15	1.41	1.34
2	A	303	0J7	N4-N3	4.23	1.41	1.34
2	B	301	0J7	N3-N2	5.11	1.39	1.32
2	A	301	0J7	N3-N2	5.66	1.40	1.32
2	A	303	0J7	N3-N2	5.83	1.40	1.32
2	B	302	0J7	N3-N2	6.01	1.41	1.32
2	A	302	0J7	N3-N2	6.06	1.41	1.32

All (65) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	OJ7	N1-N2-N3	-6.96	104.84	109.54
2	A	303	OJ7	N4-N3-N2	-5.83	105.60	109.54
2	B	302	OJ7	N4-N3-N2	-5.12	106.08	109.54
2	A	301	OJ7	N4-N3-N2	-4.64	106.40	109.54
2	A	303	OJ7	C11-C6-C5	-4.35	113.48	120.79
2	A	302	OJ7	N1-N2-N3	-3.87	106.93	109.54
2	B	302	OJ7	C11-C6-C5	-3.79	114.42	120.79
2	B	302	OJ7	N1-N2-N3	-3.66	107.07	109.54
2	A	302	OJ7	N4-N3-N2	-3.66	107.07	109.54
2	A	301	OJ7	N1-N2-N3	-3.62	107.09	109.54
2	A	303	OJ7	N1-N2-N3	-3.11	107.44	109.54
2	A	301	OJ7	C23-C24-N25	-3.02	118.45	123.43
2	B	301	OJ7	N4-C5-N1	-2.96	107.52	111.32
2	A	301	OJ7	N4-C5-N1	-2.76	107.78	111.32
2	A	302	OJ7	C23-C24-N25	-2.75	118.89	123.43
2	B	302	OJ7	C8-N12-C13	-2.57	119.99	126.59
2	A	303	OJ7	C23-C24-N25	-2.56	119.22	123.43
2	B	302	OJ7	C23-C24-N25	-2.53	119.27	123.43
2	B	301	OJ7	C23-C24-N25	-2.51	119.29	123.43
2	B	301	OJ7	N21-C20-N25	-2.49	119.84	125.47
2	B	302	OJ7	N21-C20-N25	-2.49	119.85	125.47
2	A	302	OJ7	N21-C20-N25	-2.43	119.99	125.47
2	A	303	OJ7	C23-C22-N21	-2.40	119.48	123.43
2	A	301	OJ7	C10-C9-C8	-2.35	116.88	119.72
2	A	301	OJ7	N21-C20-N25	-2.33	120.22	125.47
2	A	303	OJ7	N4-C5-N1	-2.31	108.35	111.32
2	A	303	OJ7	N21-C20-N25	-2.30	120.28	125.47
2	B	302	OJ7	C23-C22-N21	-2.23	119.76	123.43
2	B	301	OJ7	N4-N3-N2	-2.22	108.04	109.54
2	A	301	OJ7	C23-C22-N21	-2.20	119.81	123.43
2	A	303	OJ7	C17-C16-C20	-2.08	117.29	120.79
2	B	302	OJ7	N4-C5-N1	-2.00	108.75	111.32
2	B	302	OJ7	C14-C13-N12	2.03	120.26	115.91
2	A	302	OJ7	C16-C20-N25	2.04	119.66	117.40
2	A	301	OJ7	C22-C23-C24	2.15	120.03	116.72
2	A	301	OJ7	C14-C13-N12	2.20	120.64	115.91
2	A	303	OJ7	C6-C5-N1	2.33	128.07	124.09
2	A	302	OJ7	C5-N4-N3	2.66	107.62	104.92
2	A	302	OJ7	C16-C20-N21	2.66	120.35	117.40
2	A	302	OJ7	C11-C6-C7	2.71	121.67	118.16
2	B	301	OJ7	C16-C20-N25	2.74	120.42	117.40
2	A	301	OJ7	C16-C20-N25	2.81	120.50	117.40
2	A	302	OJ7	C5-N1-N2	3.09	108.06	104.92

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	OJ7	C22-N21-C20	3.28	119.76	116.12
2	A	302	OJ7	C22-N21-C20	3.42	119.92	116.12
2	A	303	OJ7	C16-C20-N21	3.42	121.19	117.40
2	B	302	OJ7	C16-C20-N25	3.57	121.35	117.40
2	B	301	OJ7	C5-N4-N3	3.62	108.59	104.92
2	B	302	OJ7	C5-N1-N2	3.66	108.64	104.92
2	A	303	OJ7	C5-N1-N2	3.73	108.71	104.92
2	B	301	OJ7	C22-N21-C20	3.88	120.43	116.12
2	B	302	OJ7	C7-C6-C5	3.91	125.68	120.14
2	A	301	OJ7	C5-N4-N3	4.08	109.06	104.92
2	A	303	OJ7	C22-N21-C20	4.12	120.70	116.12
2	B	302	OJ7	C5-N4-N3	4.36	109.34	104.92
2	B	302	OJ7	C22-N21-C20	4.38	120.98	116.12
2	A	301	OJ7	C5-N1-N2	4.41	109.40	104.92
2	A	303	OJ7	C24-N25-C20	4.42	121.02	116.12
2	B	302	OJ7	C24-N25-C20	4.57	121.20	116.12
2	B	301	OJ7	C24-N25-C20	4.58	121.20	116.12
2	A	303	OJ7	C7-C6-C5	4.75	126.86	120.14
2	A	303	OJ7	C5-N4-N3	4.85	109.84	104.92
2	A	302	OJ7	C24-N25-C20	4.97	121.64	116.12
2	A	301	OJ7	C24-N25-C20	5.04	121.72	116.12
2	B	301	OJ7	C5-N1-N2	5.92	110.93	104.92

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 26 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	OJ7	1	0
2	A	303	OJ7	8	0
2	B	301	OJ7	1	0
2	B	302	OJ7	16	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	260/263 (98%)	0.69	17 (6%) 19 21	3, 6, 17, 24	0
1	B	261/263 (99%)	0.71	21 (8%) 12 14	4, 7, 19, 28	0
All	All	521/526 (99%)	0.70	38 (7%) 15 17	3, 7, 18, 28	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	27	SER	7.8
1	B	28	ALA	5.7
1	A	290	LEU	5.5
1	B	227	THR	5.3
1	B	290	LEU	5.0
1	B	254	GLN	4.9
1	A	271[A]	ASN	4.3
1	A	254	GLN	4.1
1	B	271	ASN	3.9
1	A	228	SER	3.9
1	B	228	SER	3.7
1	B	92	ASN	3.7
1	B	289	GLY	3.2
1	A	88	LYS	3.2
1	B	255	GLY	3.1
1	A	269	GLN	2.9
1	A	227	THR	2.8
1	A	247	ILE	2.8
1	A	197	HIS	2.8
1	B	247	ILE	2.7
1	B	55	THR	2.6
1	A	44[A]	LEU	2.5
1	A	74	VAL	2.4
1	B	229	TRP	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	288	GLU	2.4
1	B	269	GLN	2.4
1	B	29	VAL	2.4
1	B	268	PRO	2.3
1	A	162	LEU	2.3
1	A	28	ALA	2.3
1	B	54	ASN	2.3
1	B	88	LYS	2.3
1	B	288	GLU	2.2
1	A	229	TRP	2.2
1	B	52	ALA	2.2
1	A	270[A]	GLN	2.1
1	B	41	GLY	2.1
1	A	142[A]	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	0J7	B	302	26/26	0.67	0.32	29,33,38,39	0
2	0J7	A	303	26/26	0.74	0.29	25,28,34,35	0
2	0J7	A	302	26/26	0.81	0.28	16,18,22,24	0
2	0J7	A	301	26/26	0.90	0.15	5,8,17,18	0
2	0J7	B	301	26/26	0.91	0.16	8,10,20,21	0
3	DMS	B	303	4/4	0.92	0.17	16,17,18,19	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.