



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 05:10 pm GMT

PDB ID : 3DRX  
Title : X-ray crystal structure of human KCTD5 protein crystallized in high-salt buffer  
Authors : Tereshko, V.; Dementieva, I.; Goldstein, S.A.N.  
Deposited on : 2008-07-11  
Resolution : 3.11 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	trunk30967
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30967


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1111 (3.14-3.10)
Clashscore	122126	1202 (3.14-3.10)
Ramachandran outliers	120053	1163 (3.14-3.10)
Sidechain outliers	120020	1163 (3.14-3.10)
RSRZ outliers	108989	1084 (3.14-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	202	<div> <div>4%</div> <div> <div></div> <div>47%</div> <div>26%</div> <div>7%</div> <div>20%</div> </div> </div>
1	B	202	<div> <div>5%</div> <div> <div></div> <div>54%</div> <div>25%</div> <div>6%</div> <div>15%</div> </div> </div>
1	C	202	<div> <div>8%</div> <div> <div></div> <div>52%</div> <div>26%</div> <div>8%</div> <div>12%</div> </div> </div>
1	D	202	<div> <div>13%</div> <div> <div></div> <div>62%</div> <div>22%</div> <div>•</div> <div>12%</div> </div> </div>
1	E	202	<div> <div>5%</div> <div> <div></div> <div>53%</div> <div>24%</div> <div>7%</div> <div>16%</div> </div> </div>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 6907 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BTB/POZ domain-containing protein KCTD5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	161	Total	C	N	O	S	0	0	0
			1306	831	223	246	6			
1	B	172	Total	C	N	O	S	0	0	0
			1385	874	236	269	6			
1	C	177	Total	C	N	O	S	0	0	0
			1430	902	243	279	6			
1	D	177	Total	C	N	O	S	0	0	0
			1427	901	243	277	6			
1	E	169	Total	C	N	O	S	0	0	0
			1359	860	232	261	6			


There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	SER	-	EXPRESSION TAG	UNP Q9NXV2
B	33	SER	-	EXPRESSION TAG	UNP Q9NXV2
C	33	SER	-	EXPRESSION TAG	UNP Q9NXV2
D	33	SER	-	EXPRESSION TAG	UNP Q9NXV2
E	33	SER	-	EXPRESSION TAG	UNP Q9NXV2


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: BTB/POZ domain-containing protein KCTD5




#### • Molecule 1: BTB/POZ domain-containing protein KCTD5






#### • Molecule 1: BTB/POZ domain-containing protein KCTD5







- Molecule 1: BTB/POZ domain-containing protein KCTD5



- Molecule 1: BTB/POZ domain-containing protein KCTD5



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	72.86Å 128.58Å 152.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 3.11 19.94 – 3.11	Depositor EDS
% Data completeness (in resolution range)	98.9 (20.00-3.11) 99.0 (19.94-3.11)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.26 (at 3.09Å)	Xtriage
Refinement program	REFMAC 5.4.0073	Depositor
R, $R_{free}$	0.227 , 0.275 0.225 , 0.275	Depositor DCC
$R_{free}$ test set	1321 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	62.6	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 68.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	6907	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/1329	0.52	0/1793
1	B	0.44	0/1409	0.53	0/1903
1	C	0.45	0/1457	0.61	1/1970 (0.1%)
1	D	0.42	0/1453	0.53	0/1963
1	E	0.43	0/1383	0.56	0/1868
All	All	0.42	0/7031	0.55	1/9497 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	C	143	ARG	NE-CZ-NH2	5.36	122.98	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1306	0	1309	62	0
1	B	1385	0	1371	46	0
1	C	1430	0	1405	67	0
1	D	1427	0	1405	46	0
1	E	1359	0	1353	65	0
All	All	6907	0	6843	274	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 274 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:152:VAL:CG1	1:E:153:PRO:HD2	1.77	1.13
1:E:38:ARG:HG2	1:E:38:ARG:HH11	1.09	1.07
1:E:152:VAL:HG12	1:E:153:PRO:HD2	1.37	1.03
1:E:165:GLU:O	1:E:168:LEU:HB2	1.69	0.92
1:A:72:ARG:HG3	1:A:73:LEU:HD13	1.49	0.91

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	155/202 (77%)	128 (83%)	22 (14%)	5 (3%)	4	24
1	B	168/202 (83%)	152 (90%)	12 (7%)	4 (2%)	6	30
1	C	175/202 (87%)	146 (83%)	23 (13%)	6 (3%)	4	22
1	D	173/202 (86%)	148 (86%)	20 (12%)	5 (3%)	5	26
1	E	165/202 (82%)	144 (87%)	17 (10%)	4 (2%)	6	30
All	All	836/1010 (83%)	718 (86%)	94 (11%)	24 (3%)	5	26

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	43	SER
1	A	74	CYS
1	A	153	PRO
1	B	81	ASP
1	C	198	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	147/181 (81%)	129 (88%)	18 (12%)	5	21
1	B	156/181 (86%)	134 (86%)	22 (14%)	4	16
1	C	161/181 (89%)	141 (88%)	20 (12%)	5	21
1	D	160/181 (88%)	154 (96%)	6 (4%)	36	70
1	E	153/181 (84%)	140 (92%)	13 (8%)	12	41
All	All	777/905 (86%)	698 (90%)	79 (10%)	8	30

5 of 79 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	166	GLU
1	C	46	VAL
1	E	119	GLU
1	B	168	LEU
1	C	35	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	164	GLN
1	E	151	GLN
1	D	156	HIS
1	C	75	GLN
1	C	195	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	161/202 (79%)	0.12	8 (4%) 29 13	53, 66, 82, 85	2 (1%)
1	B	172/202 (85%)	0.14	10 (5%) 23 10	52, 65, 85, 96	1 (0%)
1	C	177/202 (87%)	0.23	17 (9%) 8 3	52, 66, 83, 85	0
1	D	177/202 (87%)	0.58	27 (15%) 2 1	53, 68, 84, 106	0
1	E	169/202 (83%)	0.09	10 (5%) 22 10	42, 64, 79, 84	0
All	All	856/1010 (84%)	0.24	72 (8%) 11 4	42, 66, 83, 106	3 (0%)

The worst 5 of 72 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	36	ALA	8.8
1	D	78	PRO	5.8
1	D	35	LEU	5.8
1	D	41	SER	4.9
1	D	79	ASP	4.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.