



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2018 – 12:30 pm GMT

PDB ID : 1DSS  
Title : STRUCTURE OF ACTIVE-SITE CARBOXYMETHYLATED D-GLYCER  
ALDEHYDE-3-PHOSPHATE DEHYDROGENASE FROM PALINURUS  
VERSICOLOR  
Authors : Song, S.; Lin, Z.  
Deposited on : 1997-06-04  
Resolution : 1.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	9028 (1.90-1.86)
Ramachandran outliers	120053	8930 (1.90-1.86)
Sidechain outliers	120020	8930 (1.90-1.86)
RSRZ outliers	108989	8087 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	333	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> <span>%</span> <span>89%</span> <span>10%</span> <span>.</span> </div> </div>
1	R	333	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> <span>%</span> <span>86%</span> <span>12%</span> <span>.</span> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5528 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called D-GLYCERALDEHYDE-3-PHOSPHATE-DEHYDROGEN ASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	333	Total	C	N	O	S	0	0	0
			2510	1591	421	482	16			
1	R	333	Total	C	N	O	S	0	0	0
			2510	1591	421	482	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	149	CCS	CYS	CONFLICT	UNP P56649
R	149	CCS	CYS	CONFLICT	UNP P56649

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	G	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	G	1	Total	O	S	0	0
			5	4	1		
2	R	1	Total	O	S	0	0
			5	4	1		
2	R	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	G	1	Total	C	N	O	P	
			44	21	7	14	2	0
3	R	1	Total	C	N	O	P	
			44	21	7	14	2	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	226	Total	O	0	0
			226	226		
4	R	174	Total	O	0	0
			174	174		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: D-GLYCERALDEHYDE-3-PHOSPHATE-DEHYDROGENASE



#### • Molecule 1: D-GLYCERALDEHYDE-3-PHOSPHATE-DEHYDROGENASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	128.10Å 99.61Å 80.69Å 90.00° 114.40° 90.00°	Depositor
Resolution (Å)	6.00 – 1.88 6.00 – 1.88	Depositor EDS
% Data completeness (in resolution range)	81.0 (6.00-1.88) 81.0 (6.00-1.88)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.02 (at 1.88Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.172 , 0.218 0.164 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.0	Xtriage
Anisotropy	0.607	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.48 , 103.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5528	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CCS, NAD, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	G	0.71	0/2544	1.34	18/3435 (0.5%)
1	R	0.71	0/2544	1.37	21/3435 (0.6%)
All	All	0.71	0/5088	1.36	39/6870 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	2
1	R	0	4
All	All	0	6

There are no bond length outliers.

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	R	194	ARG	NE-CZ-NH1	16.58	128.59	120.30
1	G	288	ARG	NE-CZ-NH1	15.28	127.94	120.30
1	R	194	ARG	NE-CZ-NH2	-14.54	113.03	120.30
1	G	288	ARG	NE-CZ-NH2	-14.07	113.27	120.30
1	R	13	ARG	NE-CZ-NH2	-12.35	114.13	120.30

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	202	ASN	Peptide

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	G	288	ARG	Sidechain
1	R	13	ARG	Sidechain
1	R	137	TYR	Sidechain
1	R	194	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2510	0	2530	10	0
1	R	2510	0	2530	17	0
2	G	10	0	0	0	0
2	R	10	0	0	0	0
3	G	44	0	26	1	0
3	R	44	0	26	1	0
4	G	226	0	0	4	0
4	R	174	0	0	1	0
All	All	5528	0	5112	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 27 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:249:GLU:HB3	1:G:302:LYS:HE3	1.77	0.66
1:G:172:MET:CE	1:G:208:THR:HG21	2.27	0.65
1:G:286:ASP:OD1	1:G:288:ARG:HD3	2.00	0.61
1:G:136:LYS:HD2	4:G:683:HOH:O	2.01	0.60
1:R:154:LEU:HD13	1:R:214:VAL:HG21	1.85	0.58

There are no symmetry-related clashes.



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	330/333 (99%)	316 (96%)	12 (4%)	2 (1%)	27	15
1	R	330/333 (99%)	314 (95%)	14 (4%)	2 (1%)	27	15
All	All	660/666 (99%)	630 (96%)	26 (4%)	4 (1%)	27	15

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	237	VAL
1	R	237	VAL
1	R	166	GLU
1	G	166	GLU

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	268/268 (100%)	258 (96%)	10 (4%)	37	24
1	R	268/268 (100%)	253 (94%)	15 (6%)	23	11
All	All	536/536 (100%)	511 (95%)	25 (5%)	29	16

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	R	45	LYS
1	R	103	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	R	319	GLN
1	R	58	LYS
1	R	136	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	327	HIS
1	R	319	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CCS	G	149	1	6,9,10	1.24	0	4,10,12	1.64	1 (25%)
1	CCS	R	149	1	6,9,10	1.48	1 (16%)	4,10,12	1.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CCS	G	149	1	-	0/4/8/10	0/0/0/0
1	CCS	R	149	1	-	0/4/8/10	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	R	149	CCS	CA-C	2.88	1.54	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	149	CCS	CE-CD-SG	2.21	118.97	112.85

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAD	G	335	-	40,48,48	0.98	2 (5%)	44,73,73	1.45	4 (9%)
2	SO4	G	501	-	4,4,4	0.90	0	6,6,6	0.76	0
2	SO4	G	502	-	4,4,4	0.72	0	6,6,6	0.45	0
3	NAD	R	335	-	40,48,48	0.95	2 (5%)	44,73,73	1.51	5 (11%)
2	SO4	R	501	-	4,4,4	0.86	0	6,6,6	0.40	0
2	SO4	R	502	-	4,4,4	0.59	0	6,6,6	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.  
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	G	335	-	-	0/22/62/62	0/5/5/5
2	SO4	G	501	-	-	0/0/0/0	0/0/0/0
2	SO4	G	502	-	-	0/0/0/0	0/0/0/0
3	NAD	R	335	-	-	0/22/62/62	0/5/5/5
2	SO4	R	501	-	-	0/0/0/0	0/0/0/0
2	SO4	R	502	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	R	335	NAD	C3N-C7N	2.19	1.54	1.50
3	G	335	NAD	O4D-C1D	2.24	1.44	1.41
3	R	335	NAD	O4D-C1D	2.41	1.44	1.41
3	G	335	NAD	C3N-C7N	2.98	1.55	1.50

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	335	NAD	N3A-C2A-N1A	-6.34	123.44	128.86
3	R	335	NAD	N3A-C2A-N1A	-5.85	123.86	128.86
3	R	335	NAD	C4B-O4B-C1B	-3.77	105.90	109.83
3	G	335	NAD	C3N-C2N-N1N	-2.26	118.15	120.41
3	R	335	NAD	C2N-C3N-C4N	2.23	120.83	118.26

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	G	335	NAD	1	0
3	R	335	NAD	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	G	332/333 (99%)	-1.00	4 (1%) 79 81	7, 18, 36, 70	0
1	R	332/333 (99%)	-0.69	3 (0%) 84 85	8, 25, 50, 83	0
All	All	664/666 (99%)	-0.85	7 (1%) 80 82	7, 20, 47, 83	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	R	333	SER	4.8
1	G	334	ALA	4.7
1	R	212	LYS	3.8
1	R	334	ALA	3.6
1	G	333	SER	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CCS	R	149	10/11	0.99	0.04	10,16,20,22	0
1	CCS	G	149	10/11	0.99	0.05	7,11,20,20	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	R	502	5/5	0.92	0.17	50,51,53,53	0
2	SO4	G	502	5/5	0.93	0.23	49,51,53,53	0
2	SO4	G	501	5/5	0.97	0.10	26,32,34,34	0
3	NAD	R	335	44/44	0.98	0.06	14,20,28,28	0
2	SO4	R	501	5/5	0.98	0.07	24,27,29,31	0
3	NAD	G	335	44/44	0.99	0.05	10,16,21,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.