



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 01:22 pm GMT

PDB ID : 2E37  
Title : Structure of TT0471 protein from *Thermus thermophilus*  
Authors : Lokanath, N.K.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2006-11-21  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4477 (2.30-2.30)
Clashscore	122126	5072 (2.30-2.30)
Ramachandran outliers	120053	5022 (2.30-2.30)
Sidechain outliers	120020	5021 (2.30-2.30)
RSRZ outliers	108989	4374 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	<div> <div>2%</div> <div> <div></div> <div>76%</div> <div>23%</div> <div>..</div> </div> </div>
1	B	310	<div> <div>3%</div> <div> <div></div> <div>70%</div> <div>26%</div> <div>..</div> </div> </div>
1	C	310	<div> <div>%</div> <div> <div></div> <div>77%</div> <div>21%</div> <div>.</div> </div> </div>
1	D	310	<div> <div>8%</div> <div> <div></div> <div>63%</div> <div>32%</div> <div>..</div> </div> </div>
1	E	310	<div> <div>3%</div> <div> <div></div> <div>70%</div> <div>27%</div> <div>..</div> </div> </div>
1	F	310	<div> <div>3%</div> <div> <div></div> <div>69%</div> <div>27%</div> <div>..</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	310	<div><div>%</div><div><div></div><div>76%</div><div>22%</div><div></div></div><div></div></div>
1	H	310	<div><div>%</div><div><div></div><div>76%</div><div>22%</div><div></div></div><div></div></div>

## 2 Entry composition


There are 4 unique types of molecules in this entry. The entry contains 19337 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-lactate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	308	Total	C	N	O	Se	0	0	0
			2300	1460	410	427	3			
1	B	304	Total	C	N	O	Se	0	0	0
			2268	1441	405	419	3			
1	C	310	Total	C	N	O	Se	0	0	0
			2315	1468	415	429	3			
1	D	302	Total	C	N	O	Se	0	0	0
			2253	1433	400	417	3			
1	E	304	Total	C	N	O	Se	0	0	0
			2271	1442	403	423	3			
1	F	308	Total	C	N	O	Se	0	0	0
			2300	1460	410	427	3			
1	G	310	Total	C	N	O	Se	0	0	0
			2315	1468	415	429	3			
1	H	308	Total	C	N	O	Se	0	0	0
			2300	1460	410	427	3			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).



- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	H	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	123	Total	O	0	0
			123	123		
4	B	76	Total	O	0	0
			76	76		
4	C	140	Total	O	0	0
			140	140		
4	D	60	Total	O	0	0
			60	60		
4	E	84	Total	O	0	0
			84	84		
4	F	100	Total	O	0	0
			100	100		
4	G	129	Total	O	0	0
			129	129		
4	H	117	Total	O	0	0
			117	117		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: L-lactate dehydrogenase



#### • Molecule 1: L-lactate dehydrogenase




#### • Molecule 1: L-lactate dehydrogenase







• Molecule 1: L-lactate dehydrogenase



• Molecule 1: L-lactate dehydrogenase




• Molecule 1: L-lactate dehydrogenase




• Molecule 1: L-lactate dehydrogenase





• Molecule 1: L-lactate dehydrogenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	162.74Å 135.64Å 162.06Å 90.00° 113.79° 90.00°	Depositor
Resolution (Å)	44.01 – 2.30 44.01 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.1 (44.01-2.30) 95.2 (44.01-2.30)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.46 (at 2.29Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.215 , 0.255 0.214 , 0.253	Depositor DCC
$R_{free}$ test set	6900 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.2	Xtriage
Anisotropy	0.609	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 45.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	19337	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/2337	0.65	0/3175
1	B	0.41	0/2304	0.69	0/3129
1	C	0.40	0/2353	0.67	0/3197
1	D	0.39	0/2288	0.66	0/3107
1	E	0.39	0/2308	0.66	0/3136
1	F	0.43	0/2337	0.71	0/3175
1	G	0.47	1/2353 (0.0%)	0.71	0/3197
1	H	0.44	0/2337	0.70	0/3175
All	All	0.42	1/18617 (0.0%)	0.68	0/25291

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	180	TRP	NE1-CE2	8.71	1.48	1.37

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2300	0	2350	61	0
1	B	2268	0	2322	71	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2315	0	2367	47	0
1	D	2253	0	2305	89	0
1	E	2271	0	2316	71	0
1	F	2300	0	2350	84	0
1	G	2315	0	2367	50	0
1	H	2300	0	2350	49	0
2	A	44	0	26	0	0
2	C	44	0	26	0	0
2	E	44	0	26	0	0
2	G	44	0	26	1	0
3	B	5	0	0	1	0
3	H	5	0	0	0	0
4	A	123	0	0	4	0
4	B	76	0	0	2	0
4	C	140	0	0	4	0
4	D	60	0	0	0	0
4	E	84	0	0	3	0
4	F	100	0	0	5	0
4	G	129	0	0	2	0
4	H	117	0	0	2	0
All	All	19337	0	18831	484	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 484 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:79:GLN:HE22	1:F:84:THR:HA	1.14	1.09
1:D:156:TYR:CE2	1:D:201:LEU:HD23	1.96	1.01
1:D:69:ARG:HH11	1:D:69:ARG:HB3	1.24	0.99
1:D:192:GLU:HG3	1:D:193:PHE:H	1.32	0.93
1:D:192:GLU:HG3	1:D:193:PHE:N	1.85	0.92

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/310 (98%)	290 (95%)	14 (5%)	0	100	100
1	B	300/310 (97%)	279 (93%)	19 (6%)	2 (1%)	24	29
1	C	308/310 (99%)	295 (96%)	10 (3%)	3 (1%)	17	19
1	D	296/310 (96%)	282 (95%)	13 (4%)	1 (0%)	43	53
1	E	300/310 (97%)	282 (94%)	16 (5%)	2 (1%)	24	29
1	F	304/310 (98%)	288 (95%)	12 (4%)	4 (1%)	13	13
1	G	308/310 (99%)	296 (96%)	10 (3%)	2 (1%)	27	33
1	H	304/310 (98%)	294 (97%)	8 (3%)	2 (1%)	24	29
All	All	2424/2480 (98%)	2306 (95%)	102 (4%)	16 (1%)	24	29

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	264	GLU
1	F	80	ARG
1	F	81	PRO
1	H	228	TYR
1	C	52	ALA

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	231/229 (101%)	224 (97%)	7 (3%)	44 60

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	227/229 (99%)	215 (95%)	12 (5%)	25	34
1	C	232/229 (101%)	219 (94%)	13 (6%)	23	32
1	D	226/229 (99%)	214 (95%)	12 (5%)	25	34
1	E	229/229 (100%)	221 (96%)	8 (4%)	39	53
1	F	231/229 (101%)	216 (94%)	15 (6%)	19	25
1	G	232/229 (101%)	218 (94%)	14 (6%)	21	28
1	H	231/229 (101%)	221 (96%)	10 (4%)	32	43
All	All	1839/1832 (100%)	1748 (95%)	91 (5%)	27	38

5 of 91 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	296	ARG
1	E	292	GLU
1	H	104	LEU
1	D	303	GLU
1	E	178	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	41	HIS
1	E	47	HIS
1	F	79	GLN
1	D	172	HIS
1	E	117	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAD	A	1401	-	40,48,48	2.62	15 (37%)	44,73,73	2.96	11 (25%)
3	SO4	B	1501	-	4,4,4	0.61	0	6,6,6	0.82	0
2	NAD	C	1402	-	40,48,48	2.60	15 (37%)	44,73,73	2.93	12 (27%)
2	NAD	E	1403	-	40,48,48	2.61	15 (37%)	44,73,73	2.93	12 (27%)
2	NAD	G	1404	-	40,48,48	2.63	15 (37%)	44,73,73	2.96	10 (22%)
3	SO4	H	1502	-	4,4,4	0.37	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	1401	-	-	0/22/62/62	0/5/5/5
3	SO4	B	1501	-	-	0/0/0/0	0/0/0/0
2	NAD	C	1402	-	-	0/22/62/62	0/5/5/5
2	NAD	E	1403	-	-	0/22/62/62	0/5/5/5
2	NAD	G	1404	-	-	0/22/62/62	0/5/5/5
3	SO4	H	1502	-	-	0/0/0/0	0/0/0/0

The worst 5 of 60 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1402	NAD	C2D-C3D	-2.81	1.46	1.53
2	E	1403	NAD	C2D-C3D	-2.79	1.46	1.53
2	G	1404	NAD	C2D-C3D	-2.69	1.46	1.53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1401	NAD	C2D-C3D	-2.65	1.46	1.53
2	C	1402	NAD	O4D-C4D	-2.26	1.39	1.45

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1402	NAD	N3A-C2A-N1A	-12.06	118.54	128.86
2	A	1401	NAD	N3A-C2A-N1A	-12.04	118.56	128.86
2	G	1404	NAD	N3A-C2A-N1A	-12.03	118.57	128.86
2	E	1403	NAD	N3A-C2A-N1A	-12.02	118.57	128.86
2	A	1401	NAD	C4B-O4B-C1B	-10.08	99.32	109.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1501	SO4	1	0
2	G	1404	NAD	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	305/310 (98%)	-0.02	7 (2%) 60 67	31, 43, 60, 91	0
1	B	301/310 (97%)	0.18	8 (2%) 54 62	28, 47, 72, 98	0
1	C	307/310 (99%)	-0.18	2 (0%) 87 91	26, 40, 52, 61	0
1	D	299/310 (96%)	0.42	24 (8%) 12 16	32, 55, 83, 99	0
1	E	301/310 (97%)	0.24	10 (3%) 46 53	32, 46, 67, 93	0
1	F	305/310 (98%)	0.16	9 (2%) 50 57	31, 43, 73, 103	0
1	G	307/310 (99%)	-0.19	3 (0%) 82 86	27, 39, 55, 68	0
1	H	305/310 (98%)	-0.07	3 (0%) 82 86	24, 38, 63, 80	0
All	All	2430/2480 (97%)	0.07	66 (2%) 54 62	24, 43, 70, 103	0

The worst 5 of 66 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	296	ARG	5.6
1	E	191	LEU	5.0
1	F	306	PHE	4.9
1	D	306	PHE	4.7
1	D	199	ARG	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	B	1501	5/5	0.89	0.34	112,112,112,113	0
2	NAD	G	1404	44/44	0.90	0.17	50,56,67,68	0
2	NAD	C	1402	44/44	0.91	0.15	59,64,71,72	0
2	NAD	E	1403	44/44	0.92	0.15	50,63,80,81	0
2	NAD	A	1401	44/44	0.92	0.15	48,56,69,70	0
3	SO4	H	1502	5/5	0.92	0.16	100,101,102,102	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.