



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 11:59 am GMT

PDB ID : 3E3Y  
Title : Q138F HincII bound to GTTAAC and cocrystallized with 5 mM Ca<sup>2+</sup>  
Authors : Horton, N.C.; Babic, A.C.; Little, E.J.; Manohar, V.M.  
Deposited on : 2008-08-08  
Resolution : 2.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2128 (2.16-2.12)
Clashscore	122126	2253 (2.16-2.12)
Ramachandran outliers	120053	2223 (2.16-2.12)
Sidechain outliers	120020	2222 (2.16-2.12)
RSRZ outliers	108989	2086 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0;">%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 13%, yellow 13%, yellow 79%, green 79%, green 100%);"></div> <div style="position: absolute; bottom: -10px; left: 0;">79%13%• 6%</div> </div> </div>
1	B	257	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0;">2%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red 0%, red 2%, orange 2%, orange 20%, yellow 20%, yellow 72%, green 72%, green 100%);"></div> <div style="position: absolute; bottom: -10px; left: 0;">72%20%• 6%</div> </div> </div>
2	E	14	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, green 0%, green 57%, yellow 57%, yellow 43%, green 43%, green 100%);"></div> <div style="position: absolute; bottom: -10px; left: 0;">57%43%</div> </div> </div>
2	F	14	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, green 0%, green 79%, yellow 79%, yellow 21%, green 21%, green 100%);"></div> <div style="position: absolute; bottom: -10px; left: 0;">79%21%</div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4873 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type-2 restriction enzyme HindII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	242	Total	C	N	O	S	0	0	0
			1929	1259	302	362	6			
1	B	241	Total	C	N	O	S	0	0	0
			1923	1253	305	359	6			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	67	ASN	LYS	CONFLICT	UNP P44413
A	138	PHE	GLN	ENGINEERED	UNP P44413
B	67	ASN	LYS	CONFLICT	UNP P44413
B	138	PHE	GLN	ENGINEERED	UNP P44413

- Molecule 2 is a DNA chain called 5'-D(\*DGP\*DCP\*DCP\*DGP\*DGP\*DTP\*DTP\*DAP\*DAP\*DCP\*DCP\*DGP\*DGP\*DC)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	14	Total	C	N	O	P	0	0	0
			284	135	54	82	13			
2	F	14	Total	C	N	O	P	0	0	0
			284	135	54	82	13			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ca	0	0
			1	1		
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Na 1	0	0
4	A	1	Total 1	Na 1	0	0
4	F	1	Total 1	Na 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	199	Total 199	O 199	0	0
5	B	161	Total 161	O 161	0	0
5	E	47	Total 47	O 47	0	0
5	F	41	Total 41	O 41	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Type-2 restriction enzyme HindII




- Molecule 1: Type-2 restriction enzyme HindII



- Molecule 2: 5'-D(\*DGP\*DCP\*DCP\*DGP\*DGP\*DTP\*DTP\*DAP\*DAP\*DCP\*DCP\*DGP\*DGP\*DC)-3'



- Molecule 2: 5'-D(\*DGP\*DCP\*DCP\*DGP\*DGP\*DTP\*DTP\*DAP\*DAP\*DCP\*DCP\*DGP\*DGP\*DC)-3'



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.48Å 91.64Å 65.78Å 90.00° 104.92° 90.00°	Depositor
Resolution (Å)	43.06 – 2.13 43.06 – 2.13	Depositor EDS
% Data completeness (in resolution range)	99.9 (43.06-2.13) 99.9 (43.06-2.13)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.92 (at 2.14Å)	Xtriage
Refinement program	PHENIX ?	Depositor
R, $R_{free}$	0.180 , 0.234 0.175 , 0.228	Depositor DCC
$R_{free}$ test set	1606 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.7	Xtriage
Anisotropy	0.124	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 62.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4873	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.31	0/1973	0.46	0/2678
1	B	0.29	0/1967	0.46	0/2666
2	E	0.68	0/318	1.41	3/489 (0.6%)
2	F	0.69	0/318	1.37	3/489 (0.6%)
All	All	0.38	0/4576	0.69	6/6322 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	12	DG	O4'-C1'-N9	10.94	115.66	108.00
2	F	12	DG	O4'-C1'-N9	9.29	114.50	108.00
2	E	12	DG	C1'-O4'-C4'	-6.58	103.52	110.10
2	F	6	DT	C5-C4-O4	-5.63	120.96	124.90
2	E	3	DC	O4'-C1'-N1	-5.23	104.34	108.00
2	F	12	DG	C1'-O4'-C4'	-5.05	105.05	110.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1929	0	1813	34	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1923	0	1822	51	0
2	E	284	0	158	4	0
2	F	284	0	158	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	F	1	0	0	0	0
5	A	199	0	0	2	0
5	B	161	0	0	2	0
5	E	47	0	0	0	0
5	F	41	0	0	0	0
All	All	4873	0	3951	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:70:ILE:HD13	1:B:79:LEU:HD21	1.53	0.90
1:A:11:ILE:HG12	1:A:44:PHE:CE2	2.21	0.76
1:B:11:ILE:HG12	1:B:44:PHE:CE2	2.23	0.74
1:B:141:ASN:ND2	1:B:209:GLN:HE21	1.87	0.72
1:A:10:ASP:O	1:A:14:ILE:HG12	1.89	0.72
1:A:4:ILE:HD11	1:A:126:LEU:HD22	1.72	0.69
1:A:14:ILE:HG13	1:A:44:PHE:CE1	2.27	0.69
1:B:195:PRO:HA	1:B:198:LEU:HD23	1.73	0.69
1:B:170:GLU:HB2	1:B:212:VAL:HB	1.75	0.68
1:B:127:ASP:HB3	1:B:168:TYR:CD1	2.31	0.66
1:A:110:ASN:HD22	1:A:110:ASN:N	1.93	0.65
1:B:38:GLU:C	1:B:40:LEU:H	2.01	0.64
1:A:132:ASN:HD22	1:A:134:SER:H	1.44	0.63
1:A:14:ILE:HG13	1:A:44:PHE:HE1	1.63	0.63
1:A:132:ASN:HD22	1:A:132:ASN:C	2.02	0.63
1:B:73:HIS:HD2	1:B:97:GLU:OE2	1.82	0.62
1:A:132:ASN:ND2	1:A:134:SER:H	2.00	0.60
1:B:52:LEU:HD22	1:B:119:LYS:HB2	1.84	0.59
1:A:205:ALA:HA	1:B:203:ALA:O	2.02	0.59
1:B:67:ASN:HB3	1:B:70:ILE:HD12	1.85	0.59
1:A:127:ASP:HB3	1:A:168:TYR:CD1	2.39	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:5:LYS:HB3	1:B:6:PRO:HD3	1.84	0.58
1:A:4:ILE:CD1	1:A:126:LEU:HD22	2.34	0.57
1:A:203:ALA:O	1:B:205:ALA:HA	2.04	0.57
1:A:108:LYS:HE3	1:A:110:ASN:ND2	2.19	0.57
1:B:195:PRO:HA	1:B:198:LEU:CD2	2.34	0.57
1:B:141:ASN:HD22	1:B:209:GLN:HE21	1.51	0.56
1:B:4:ILE:CD1	1:B:126:LEU:HD22	2.37	0.55
1:B:56:GLN:HE22	1:B:113:ALA:H	1.52	0.55
1:B:38:GLU:HG2	1:B:171:VAL:HG11	1.88	0.55
1:B:133:ILE:HD11	1:B:174:GLU:HB3	1.88	0.54
1:A:71:ILE:HD13	1:A:71:ILE:H	1.72	0.54
1:B:91:ARG:NE	1:B:108:LYS:HD3	2.22	0.54
1:B:91:ARG:CZ	1:B:108:LYS:HD3	2.37	0.54
1:B:109:GLN:NE2	1:B:110:ASN:N	2.56	0.53
1:B:56:GLN:NE2	1:B:113:ALA:H	2.06	0.53
1:B:140:PRO:HG3	5:B:279:HOH:O	2.08	0.52
1:A:194:GLU:HG3	5:A:299:HOH:O	2.08	0.52
1:A:85:LEU:HD22	1:A:163:LEU:HD21	1.92	0.52
1:A:110:ASN:N	5:A:440:HOH:O	2.43	0.51
1:A:123:TYR:O	1:A:164:PHE:HA	2.10	0.51
1:B:11:ILE:O	1:B:14:ILE:HG13	2.12	0.50
1:A:147:LYS:HA	1:A:150:GLN:HE21	1.76	0.50
1:B:108:LYS:HZ1	2:F:11:DC:H5"	1.77	0.50
1:A:173:TRP:CE3	1:A:180:LEU:HG	2.46	0.50
1:B:14:ILE:HD12	1:B:15:LEU:N	2.28	0.49
1:A:52:LEU:O	1:A:117:LEU:HA	2.13	0.49
1:A:4:ILE:HG13	1:A:117:LEU:HD22	1.95	0.49
1:B:108:LYS:NZ	2:F:11:DC:H5"	2.27	0.49
1:B:109:GLN:O	2:E:7:DT:H4'	2.13	0.48
1:B:38:GLU:C	1:B:40:LEU:N	2.66	0.48
1:A:4:ILE:HG23	1:A:11:ILE:HD12	1.96	0.47
1:B:41:VAL:HG21	1:B:169:LEU:CD2	2.45	0.47
1:B:109:GLN:NE2	1:B:110:ASN:H	2.13	0.47
1:B:169:LEU:HD12	1:B:186:SER:O	2.15	0.47
1:B:38:GLU:O	1:B:40:LEU:HD23	2.14	0.47
1:B:38:GLU:O	1:B:40:LEU:N	2.48	0.46
1:B:73:HIS:CD2	1:B:97:GLU:OE2	2.67	0.46
1:A:132:ASN:HD21	1:A:134:SER:HB3	1.82	0.45
1:B:127:ASP:HB3	1:B:168:TYR:CE1	2.51	0.45
1:B:72:GLY:HA2	1:B:97:GLU:HG3	1.99	0.44
1:A:5:LYS:HB3	1:A:6:PRO:HD3	1.99	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:200:ILE:HG12	1:A:208:ILE:HD13	2.00	0.44
1:A:110:ASN:ND2	1:A:110:ASN:N	2.61	0.44
1:B:140:PRO:HD2	1:B:210:PHE:O	2.17	0.44
1:B:139:ALA:HA	1:B:140:PRO:HD3	1.87	0.43
1:A:85:LEU:HA	1:A:85:LEU:HD12	1.90	0.43
1:B:67:ASN:N	1:B:68:PRO:HD3	2.34	0.42
1:B:167:ASN:ND2	5:B:303:HOH:O	2.52	0.42
1:B:14:ILE:HD12	1:B:15:LEU:HG	2.02	0.42
1:B:108:LYS:HZ2	1:B:108:LYS:HB2	1.84	0.41
1:B:123:TYR:O	1:B:164:PHE:HA	2.20	0.41
1:B:109:GLN:N	1:B:109:GLN:HE21	2.19	0.41
1:B:42:TYR:CE2	1:B:55:LYS:HG2	2.55	0.41
1:B:141:ASN:ND2	2:E:9:DA:H62	2.19	0.41
1:A:59:TYR:HH	1:A:123:TYR:HH	1.65	0.41
1:A:3:PHE:CZ	1:A:119:LYS:HB3	2.56	0.41
1:B:70:ILE:CD1	1:B:79:LEU:HD21	2.38	0.41
1:A:93:LYS:HE3	2:E:14:DC:OP2	2.21	0.41
2:E:8:DA:H2'	2:E:9:DA:O4'	2.21	0.41
1:A:127:ASP:HB3	1:A:168:TYR:CE1	2.56	0.40
1:B:39:LYS:HE3	1:B:39:LYS:HB3	1.92	0.40
1:B:57:TYR:CE1	1:B:107:GLU:HB2	2.56	0.40
1:A:132:ASN:ND2	1:A:132:ASN:C	2.73	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/257 (92%)	230 (98%)	6 (2%)	0	100	100
1	B	237/257 (92%)	229 (97%)	7 (3%)	1 (0%)	36	30
All	All	473/514 (92%)	459 (97%)	13 (3%)	1 (0%)	49	47

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	39	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/229 (86%)	189 (96%)	8 (4%)	33	30
1	B	196/229 (86%)	186 (95%)	10 (5%)	26	21
All	All	393/458 (86%)	375 (95%)	18 (5%)	29	25

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	ILE
1	A	85	LEU
1	A	110	ASN
1	A	118	VAL
1	A	132	ASN
1	A	138	PHE
1	A	148	LEU
1	A	231	LEU
1	B	38	GLU
1	B	55	LYS
1	B	86	LEU
1	B	108	LYS
1	B	109	GLN
1	B	138	PHE
1	B	198	LEU
1	B	215	LEU
1	B	216	ASP
1	B	231	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	48	ASN
1	A	98	ASN
1	A	103	ASN
1	A	132	ASN
1	A	150	GLN
1	A	211	HIS
1	B	56	GLN
1	B	73	HIS
1	B	109	GLN
1	B	141	ASN
1	B	167	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	242/257 (94%)	-0.24	3 (1%) 79 82	18, 29, 43, 61	0
1	B	241/257 (93%)	0.00	6 (2%) 57 64	19, 34, 51, 66	0
2	E	14/14 (100%)	-0.48	0 100 100	20, 28, 32, 33	0
2	F	14/14 (100%)	-0.50	0 100 100	19, 28, 37, 39	0
All	All	511/542 (94%)	-0.14	9 (1%) 68 73	18, 31, 49, 66	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	177	GLY	4.1
1	B	40	LEU	3.6
1	B	220	ASN	3.0
1	A	20	VAL	3.0
1	A	175	LEU	2.7
1	B	20	VAL	2.4
1	B	181	VAL	2.4
1	B	41	VAL	2.1
1	B	18	GLN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NA	F	15	1/1	0.96	0.14	39,39,39,39	0
4	NA	B	260	1/1	0.97	0.16	33,33,33,33	0
4	NA	A	260	1/1	0.97	0.18	30,30,30,30	0
3	CA	B	259	1/1	0.99	0.11	30,30,30,30	0
3	CA	A	259	1/1	0.99	0.08	25,25,25,25	0

## 6.5 Other polymers

There are no such residues in this entry.