



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 2, 2019 – 03:49 PM EDT

PDB ID : 6E4Z  
Title : Anti-PCSK9 fab 6E2 bound to the modified N-terminal peptide from PCSK9  
Authors : Ultsch, M.H.; Kirchhofer, D.K.  
Deposited on : 2018-07-18  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031633  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031633


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4343 (2.20-2.20)
Clashscore	122126	5027 (2.20-2.20)
Ramachandran outliers	120053	4952 (2.20-2.20)
Sidechain outliers	120020	4953 (2.20-2.20)
RSRZ outliers	108989	4245 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	225	<div> <div>5%</div> <div>84%</div> <div>10%</div> <div>7%</div> </div>
2	L	219	<div> <div>5%</div> <div>87%</div> <div>12%</div> </div>
3	P	22	<div> <div>9%</div> <div>64%</div> <div>5%</div> <div>32%</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3573 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 6E2 heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	H	209	Total	C	N	O	P	S	0	2	0
			1620	1021	272	317	2	8			

- Molecule 2 is a protein called 6E2 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	218	Total	C	N	O	S	0	0	0
			1682	1057	285	333	7			


- Molecule 3 is a protein called Proprotein convertase subtilisin/kexin type 9.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	P	15	Total	C	N	O	P	S	0	1	0
			139	77	19	40	2	1			

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	1	Total	Zn	0	0
			1	1		
4	L	4	Total	Zn	0	0
			4	4		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	L	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	64	Total	O	0	0
			64	64		
6	L	50	Total	O	0	0
			50	50		
6	P	8	Total	O	0	0
			8	8		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 6E2 heavy chain



- Molecule 2: 6E2 light chain



- Molecule 3: Proprotein convertase subtilisin/kexin type 9



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.93Å 99.78Å 86.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.42 – 2.20 33.42 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (33.42-2.20) 99.8 (33.42-2.20)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.11 (at 2.20Å)	Xtriage
Refinement program	PHENIX (1.10.1-2155_2155: ???)	Depositor
R, $R_{free}$	0.195 , 0.236 0.195 , 0.236	Depositor DCC
$R_{free}$ test set	1060 reflections (3.35%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.7	Xtriage
Anisotropy	0.105	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 46.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3573	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HIP, ZN, SEP, SO4, NEP, TYS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	H	0.41	0/1627	0.60	0/2210
2	L	0.38	0/1720	0.57	0/2335
3	P	0.42	0/100	0.54	0/130
All	All	0.39	0/3447	0.58	0/4675

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1620	0	1586	12	0
2	L	1682	0	1641	16	0
3	P	139	0	108	1	0
4	H	1	0	0	0	0
4	L	4	0	0	0	0
5	L	5	0	0	0	0
6	H	64	0	0	1	0
6	L	50	0	0	3	0
6	P	8	0	0	0	0
All	All	3573	0	3335	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:27(D):HIS:O	2:L:28:ASN:N	2.20	0.72
2:L:27(D):HIS:CE1	3:P:49:GLU:OE2	2.44	0.69
1:H:181:VAL:HG11	2:L:135:LEU:HD22	1.76	0.68
1:H:148:GLU:OE1	6:H:401:HOH:O	2.13	0.67
1:H:119:PRO:HB3	1:H:145:TYR:HB3	1.77	0.66
1:H:63:VAL:HG13	1:H:67:PHE:HB2	1.81	0.62
2:L:163:VAL:HG22	2:L:175:LEU:HD12	1.87	0.56
1:H:22:CYS:SG	1:H:92[B]:CYS:SG	3.01	0.52
2:L:27:LYS:NZ	6:L:402:HOH:O	2.43	0.51
2:L:151:ASP:HA	2:L:191:VAL:HG13	1.96	0.48
1:H:1:GLU:O	1:H:26:ARG:NH1	2.46	0.47
2:L:91:HIS:HD2	6:L:444:HOH:O	1.96	0.47
2:L:120:PRO:HG2	2:L:130:ALA:HB1	1.97	0.46
2:L:155:GLN:HG3	2:L:158:ASN:HD21	1.79	0.46
2:L:89:MET:CE	2:L:91:HIS:HB3	2.47	0.45
2:L:108:ARG:HH12	2:L:111:ALA:HB2	1.81	0.44
1:H:170:LEU:HD13	1:H:176:TYR:CZ	2.52	0.44
2:L:91:HIS:HE1	6:L:422:HOH:O	1.99	0.44
2:L:122:ASP:O	2:L:126:LYS:HG2	2.18	0.43
1:H:6:GLU:CD	1:H:106:GLY:H	2.21	0.43
2:L:105:GLU:OE2	2:L:173:TYR:OH	2.30	0.43
1:H:193:THR:HB	1:H:210:LYS:HE3	2.01	0.42
2:L:145:LYS:HB2	2:L:197:THR:HB	2.02	0.42
1:H:145:TYR:CE1	1:H:150:VAL:HG13	2.55	0.41
2:L:108:ARG:NH1	2:L:111:ALA:HB2	2.35	0.41
1:H:1:GLU:HG2	1:H:26:ARG:HD2	2.02	0.41
1:H:14:PRO:HD3	1:H:112:SER:C	2.41	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	205/225 (91%)	197 (96%)	7 (3%)	1 (0%)	31	33
2	L	216/219 (99%)	209 (97%)	5 (2%)	2 (1%)	19	17
3	P	11/22 (50%)	11 (100%)	0	0	100	100
All	All	432/466 (93%)	417 (96%)	12 (3%)	3 (1%)	24	23

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	L	27(E)	SER
1	H	161	SER
2	L	212	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	180/192 (94%)	178 (99%)	2 (1%)	76	87
2	L	191/192 (100%)	187 (98%)	4 (2%)	56	70
3	P	11/16 (69%)	11 (100%)	0	100	100
All	All	382/400 (96%)	376 (98%)	6 (2%)	65	79

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	28	THR
1	H	135	THR
2	L	27(C)	LEU
2	L	105	GLU
2	L	114	SER
2	L	185	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no

such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	HIP	H	58[B]	1	11,14,15	5.45	4 (36%)	8,20,22	2.02	5 (62%)
1	NEP	H	58[A]	1	11,14,15	1.76	3 (27%)	6,20,22	1.80	1 (16%)
3	TYS	P	38	3	15,16,17	1.13	1 (6%)	18,22,24	1.02	1 (5%)
3	SEP	P	47[A]	3	9,9,10	1.55	1 (11%)	9,12,14	1.94	2 (22%)
3	SEP	P	47[B]	3	9,9,10	1.58	1 (11%)	9,12,14	2.00	2 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HIP	H	58[B]	1	-	0/4/12/14	0/1/1/1
1	NEP	H	58[A]	1	-	0/4/12/14	0/1/1/1
3	TYS	P	38	3	-	0/9/11/13	0/1/1/1
3	SEP	P	47[A]	3	-	0/5/8/10	0/0/0/0
3	SEP	P	47[B]	3	-	0/5/8/10	0/0/0/0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	58[B]	HIP	P-O3P	-3.19	1.48	1.54
3	P	38	TYS	OH-CZ	-2.79	1.38	1.42

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	58[B]	HIP	CG-ND1	-2.36	1.34	1.37
1	H	58[A]	NEP	P-O1P	2.52	1.59	1.54
1	H	58[A]	NEP	P-O2P	2.79	1.60	1.54
3	P	47[B]	SEP	P-O1P	3.36	1.61	1.50
3	P	47[A]	SEP	P-O1P	3.40	1.61	1.50
1	H	58[B]	HIP	P-O2P	3.45	1.61	1.54
1	H	58[A]	NEP	CD2-CG	3.63	1.41	1.36
1	H	58[B]	HIP	P-O1P	17.20	1.62	1.47

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	58[B]	HIP	O2P-P-O1P	-3.27	106.37	113.44
1	H	58[A]	NEP	O2P-P-O1P	-2.79	96.18	106.82
1	H	58[B]	HIP	CG-CB-CA	-2.76	108.68	114.02
3	P	47[A]	SEP	P-OG-CB	-2.31	111.93	118.30
1	H	58[B]	HIP	O-C-CA	-2.23	119.01	124.98
3	P	47[B]	SEP	P-OG-CB	-2.08	112.58	118.30
1	H	58[B]	HIP	O3P-P-O1P	-2.07	108.96	113.44
1	H	58[B]	HIP	CD2-NE2-CE1	2.02	108.93	105.78
3	P	38	TYS	OH-CZ-CE2	2.41	123.25	118.64
3	P	47[A]	SEP	OG-CB-CA	4.63	112.65	108.14
3	P	47[B]	SEP	OG-CB-CA	4.71	112.73	108.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 5 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	SO4	L	305	4	4,4,4	0.20	0	6,6,6	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SO4	L	305	4	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	H	208/225 (92%)	0.07	11 (5%) 26 25	32, 56, 119, 144	0
2	L	218/219 (99%)	0.16	10 (4%) 32 31	35, 64, 105, 137	0
3	P	13/22 (59%)	0.21	2 (15%) 2 1	41, 49, 99, 128	0
All	All	439/466 (94%)	0.12	23 (5%) 27 26	32, 61, 107, 144	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	P	35	ASP	5.8
2	L	184	ALA	4.6
2	L	27(E)	SER	3.8
2	L	145	LYS	3.3
1	H	184	VAL	3.3
1	H	190	GLY	3.0
3	P	36	GLY	3.0
2	L	29	GLY	2.9
1	H	161	SER	2.8
1	H	192	GLN	2.8
1	H	193	THR	2.7
2	L	183	LYS	2.5
2	L	213	GLU	2.5
2	L	28	ASN	2.5
1	H	213	PRO	2.3
2	L	154	LEU	2.1
2	L	143	GLU	2.1
1	H	154	TRP	2.1
1	H	185	PRO	2.1
1	H	189	LEU	2.0
1	H	138	LEU	2.0
1	H	214	LYS	2.0
2	L	144	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SEP	P	47[B]	10/11	0.75	0.16	54,80,102,106	10
3	SEP	P	47[A]	10/11	0.75	0.16	54,80,101,103	10
1	NEP	H	58[A]	14/15	0.92	0.19	31,41,53,68	14
1	HIP	H	58[B]	14/15	0.96	0.17	32,39,43,49	14
3	TYS	P	38	16/17	0.98	0.11	31,35,47,48	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	ZN	L	301	1/1	0.88	0.04	102,102,102,102	0
4	ZN	L	302	1/1	0.94	0.06	108,108,108,108	0
4	ZN	L	303	1/1	0.94	0.04	118,118,118,118	0
5	SO4	L	305	5/5	0.94	0.16	75,76,91,99	0
4	ZN	L	304	1/1	0.95	0.05	103,103,103,103	0
4	ZN	H	301	1/1	0.99	0.15	44,44,44,44	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.