



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 08:35 pm GMT

PDB ID : 3E5U  
Title : OCPA complexed CprK (C200S)  
Authors : Levy, C.  
Deposited on : 2008-08-14  
Resolution : 1.83 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	3313 (1.86-1.82)
Clashscore	122126	3530 (1.86-1.82)
Ramachandran outliers	120053	3495 (1.86-1.82)
Sidechain outliers	120020	3496 (1.86-1.82)
RSRZ outliers	108989	3265 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	<div> <div>8%</div> <div> <div></div> <div>74%</div> <div>11%</div> <div>•</div> <div>14%</div> </div> </div>
1	B	250	<div> <div>7%</div> <div> <div></div> <div>76%</div> <div>10%</div> <div>•</div> <div>13%</div> </div> </div>
1	C	250	<div> <div>8%</div> <div> <div></div> <div>74%</div> <div>11%</div> <div>•</div> <div>12%</div> </div> </div>
1	D	250	<div> <div>6%</div> <div> <div></div> <div>77%</div> <div>10%</div> <div>•</div> <div>12%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	3C4	A	505	-	X	-	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7860 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclic nucleotide-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	219	Total	C	N	O	S	0	2	0
			1757	1132	289	325	11			
1	A	215	Total	C	N	O	S	0	2	0
			1720	1110	282	317	11			
1	D	220	Total	C	N	O	S	0	2	0
			1749	1128	285	325	11			
1	B	218	Total	C	N	O	S	0	2	0
			1754	1130	289	326	9			

There are 76 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	200	SER	CYS	ENGINEERED	UNP Q18R04
C	233	SER	-	EXPRESSION TAG	UNP Q18R04
C	234	ASP	-	EXPRESSION TAG	UNP Q18R04
C	235	PRO	-	EXPRESSION TAG	UNP Q18R04
C	236	ASN	-	EXPRESSION TAG	UNP Q18R04
C	237	SER	-	EXPRESSION TAG	UNP Q18R04
C	238	SER	-	EXPRESSION TAG	UNP Q18R04
C	239	SER	-	EXPRESSION TAG	UNP Q18R04
C	240	VAL	-	EXPRESSION TAG	UNP Q18R04
C	241	ASP	-	EXPRESSION TAG	UNP Q18R04
C	242	LYS	-	EXPRESSION TAG	UNP Q18R04
C	243	LEU	-	EXPRESSION TAG	UNP Q18R04
C	244	ALA	-	EXPRESSION TAG	UNP Q18R04
C	245	ALA	-	EXPRESSION TAG	UNP Q18R04
C	246	ALA	-	EXPRESSION TAG	UNP Q18R04
C	247	LEU	-	EXPRESSION TAG	UNP Q18R04
C	248	ASP	-	EXPRESSION TAG	UNP Q18R04
C	249	HIS	-	EXPRESSION TAG	UNP Q18R04
C	250	HIS	-	EXPRESSION TAG	UNP Q18R04
A	200	SER	CYS	ENGINEERED	UNP Q18R04
A	233	SER	-	EXPRESSION TAG	UNP Q18R04

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	234	ASP	-	EXPRESSION TAG	UNP Q18R04
A	235	PRO	-	EXPRESSION TAG	UNP Q18R04
A	236	ASN	-	EXPRESSION TAG	UNP Q18R04
A	237	SER	-	EXPRESSION TAG	UNP Q18R04
A	238	SER	-	EXPRESSION TAG	UNP Q18R04
A	239	SER	-	EXPRESSION TAG	UNP Q18R04
A	240	VAL	-	EXPRESSION TAG	UNP Q18R04
A	241	ASP	-	EXPRESSION TAG	UNP Q18R04
A	242	LYS	-	EXPRESSION TAG	UNP Q18R04
A	243	LEU	-	EXPRESSION TAG	UNP Q18R04
A	244	ALA	-	EXPRESSION TAG	UNP Q18R04
A	245	ALA	-	EXPRESSION TAG	UNP Q18R04
A	246	ALA	-	EXPRESSION TAG	UNP Q18R04
A	247	LEU	-	EXPRESSION TAG	UNP Q18R04
A	248	ASP	-	EXPRESSION TAG	UNP Q18R04
A	249	HIS	-	EXPRESSION TAG	UNP Q18R04
A	250	HIS	-	EXPRESSION TAG	UNP Q18R04
D	200	SER	CYS	ENGINEERED	UNP Q18R04
D	233	SER	-	EXPRESSION TAG	UNP Q18R04
D	234	ASP	-	EXPRESSION TAG	UNP Q18R04
D	235	PRO	-	EXPRESSION TAG	UNP Q18R04
D	236	ASN	-	EXPRESSION TAG	UNP Q18R04
D	237	SER	-	EXPRESSION TAG	UNP Q18R04
D	238	SER	-	EXPRESSION TAG	UNP Q18R04
D	239	SER	-	EXPRESSION TAG	UNP Q18R04
D	240	VAL	-	EXPRESSION TAG	UNP Q18R04
D	241	ASP	-	EXPRESSION TAG	UNP Q18R04
D	242	LYS	-	EXPRESSION TAG	UNP Q18R04
D	243	LEU	-	EXPRESSION TAG	UNP Q18R04
D	244	ALA	-	EXPRESSION TAG	UNP Q18R04
D	245	ALA	-	EXPRESSION TAG	UNP Q18R04
D	246	ALA	-	EXPRESSION TAG	UNP Q18R04
D	247	LEU	-	EXPRESSION TAG	UNP Q18R04
D	248	ASP	-	EXPRESSION TAG	UNP Q18R04
D	249	HIS	-	EXPRESSION TAG	UNP Q18R04
D	250	HIS	-	EXPRESSION TAG	UNP Q18R04
B	200	SER	CYS	ENGINEERED	UNP Q18R04
B	233	SER	-	EXPRESSION TAG	UNP Q18R04
B	234	ASP	-	EXPRESSION TAG	UNP Q18R04
B	235	PRO	-	EXPRESSION TAG	UNP Q18R04
B	236	ASN	-	EXPRESSION TAG	UNP Q18R04
B	237	SER	-	EXPRESSION TAG	UNP Q18R04

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	238	SER	-	EXPRESSION TAG	UNP Q18R04
B	239	SER	-	EXPRESSION TAG	UNP Q18R04
B	240	VAL	-	EXPRESSION TAG	UNP Q18R04
B	241	ASP	-	EXPRESSION TAG	UNP Q18R04
B	242	LYS	-	EXPRESSION TAG	UNP Q18R04
B	243	LEU	-	EXPRESSION TAG	UNP Q18R04
B	244	ALA	-	EXPRESSION TAG	UNP Q18R04
B	245	ALA	-	EXPRESSION TAG	UNP Q18R04
B	246	ALA	-	EXPRESSION TAG	UNP Q18R04
B	247	LEU	-	EXPRESSION TAG	UNP Q18R04
B	248	ASP	-	EXPRESSION TAG	UNP Q18R04
B	249	HIS	-	EXPRESSION TAG	UNP Q18R04
B	250	HIS	-	EXPRESSION TAG	UNP Q18R04

- Molecule 2 is (3-CHLORO-4-HYDROXYPHENYL)ACETIC ACID (three-letter code: 3C4) (formula:  $C_8H_7ClO_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	Cl	O	0	0
			12	8	1	3		
2	A	1	Total	C	Cl	O	0	0
			12	8	1	3		
2	A	1	Total	C	Cl	O	0	0
			12	8	1	3		
2	D	1	Total	C	Cl	O	0	0
			12	8	1	3		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	Cl	O	0	0
			12	8	1	3		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	O	P	0	0
			5	4	1		
3	D	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	219	Total	O	0	0
			219	219		
5	A	197	Total	O	0	0
			197	197		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	178	Total 178	O 178	0	0
5	B	210	Total 210	O 210	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyclic nucleotide-binding protein




- Molecule 1: Cyclic nucleotide-binding protein




- Molecule 1: Cyclic nucleotide-binding protein



- Molecule 1: Cyclic nucleotide-binding protein





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.14Å 117.17Å 84.88Å 90.00° 94.65° 90.00°	Depositor
Resolution (Å)	19.21 – 1.83 19.21 – 1.83	Depositor EDS
% Data completeness (in resolution range)	97.1 (19.21-1.83) 97.1 (19.21-1.83)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.31 (at 1.84Å)	Xtriage
Refinement program	REFMAC 5.2	Depositor
R, $R_{free}$	0.182 , 0.227 0.187 , 0.230	Depositor DCC
$R_{free}$ test set	4666 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.2	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 47.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7860	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NA, PO4, 3C4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.06	3/1761 (0.2%)	0.93	5/2375 (0.2%)
1	B	1.10	3/1795 (0.2%)	0.97	7/2421 (0.3%)
1	C	1.05	2/1799 (0.1%)	0.96	7/2426 (0.3%)
1	D	1.03	0/1789	0.92	2/2413 (0.1%)
All	All	1.06	8/7144 (0.1%)	0.95	21/9635 (0.2%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	68	GLU	CG-CD	7.86	1.63	1.51
1	A	58	GLU	CG-CD	6.80	1.62	1.51
1	A	168	VAL	CB-CG2	5.88	1.65	1.52
1	C	68	GLU	CD-OE2	5.85	1.32	1.25
1	B	68	GLU	CG-CD	5.63	1.60	1.51

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	26	ARG	NE-CZ-NH2	-9.24	115.68	120.30
1	B	117	ARG	NE-CZ-NH2	-8.22	116.19	120.30
1	B	64	ASP	CB-CG-OD1	8.09	125.58	118.30
1	D	56	LEU	CB-CG-CD1	-8.00	97.40	111.00
1	A	26	ARG	NE-CZ-NH2	-7.97	116.32	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1720	0	1736	16	0
1	B	1754	0	1775	22	0
1	C	1757	0	1774	29	0
1	D	1749	0	1762	20	0
2	A	24	0	12	0	0
2	B	12	0	6	0	0
2	C	12	0	5	0	0
2	D	12	0	6	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	1	0
4	A	1	0	0	0	0
5	A	197	0	0	5	0
5	B	210	0	0	10	0
5	C	219	0	0	8	0
5	D	178	0	0	4	0
All	All	7860	0	7076	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 84 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:222:LEU:HD23	5:C:1977:HOH:O	1.62	0.99
1:D:153:ILE:HD13	1:D:187:THR:HG21	1.49	0.93
1:B:49[B]:GLU:HG2	5:B:1313:HOH:O	1.73	0.89
1:B:122:MET:SD	5:B:1975:HOH:O	2.32	0.87
1:C:153:ILE:HD11	1:C:189:VAL:HG21	1.64	0.77

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/250 (85%)	207 (97%)	5 (2%)	1 (0%)	31	16
1	B	218/250 (87%)	214 (98%)	4 (2%)	0	100	100
1	C	219/250 (88%)	214 (98%)	4 (2%)	1 (0%)	31	16
1	D	220/250 (88%)	214 (97%)	5 (2%)	1 (0%)	31	16
All	All	870/1000 (87%)	849 (98%)	18 (2%)	3 (0%)	43	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	20	PHE
1	C	20	PHE
1	A	20	PHE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/220 (86%)	187 (99%)	2 (1%)	76	67
1	B	193/220 (88%)	189 (98%)	4 (2%)	56	41
1	C	193/220 (88%)	191 (99%)	2 (1%)	78	71
1	D	190/220 (86%)	188 (99%)	2 (1%)	76	67
All	All	765/880 (87%)	755 (99%)	10 (1%)	71	61

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	170	ASP
1	D	209	LYS
1	B	189	VAL
1	A	219	LEU
1	B	149	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	81	ASN
1	C	180	GLN
1	D	92	ASN
1	D	228	GLN
1	B	148	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	3C4	A	501	-	9,12,12	4.34	4 (44%)	13,16,16	2.02	6 (46%)
2	3C4	A	505	-	9,12,12	4.22	7 (77%)	13,16,16	3.50	7 (53%)
2	3C4	B	503	-	9,12,12	4.94	4 (44%)	13,16,16	2.35	6 (46%)
3	PO4	B	601	-	4,4,4	1.10	0	6,6,6	0.94	0
2	3C4	C	504	-	9,12,12	3.68	5 (55%)	13,16,16	2.46	5 (38%)
3	PO4	C	603	-	4,4,4	0.71	0	6,6,6	0.46	0
2	3C4	D	502	-	9,12,12	4.34	5 (55%)	13,16,16	2.25	5 (38%)
3	PO4	D	602	-	4,4,4	0.92	0	6,6,6	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3C4	A	501	-	-	0/2/4/4	0/1/1/1
2	3C4	A	505	-	-	0/2/4/4	0/1/1/1
2	3C4	B	503	-	-	0/2/4/4	0/1/1/1
3	PO4	B	601	-	-	0/0/0/0	0/0/0/0
2	3C4	C	504	-	-	0/2/4/4	0/1/1/1
3	PO4	C	603	-	-	0/0/0/0	0/0/0/0
2	3C4	D	502	-	-	0/2/4/4	0/1/1/1
3	PO4	D	602	-	-	0/0/0/0	0/0/0/0

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	503	3C4	CE2-CD2	-7.60	1.25	1.38
2	A	505	3C4	CE2-CD2	-7.07	1.26	1.38
2	A	501	3C4	CE2-CD2	-6.01	1.28	1.38
2	D	502	3C4	CE2-CD2	-5.01	1.30	1.38
2	C	504	3C4	CE2-CD2	-4.83	1.30	1.38

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	505	3C4	CE2-CZ-CE1	-8.42	110.31	118.55
2	C	504	3C4	CE2-CZ-CE1	-6.04	112.64	118.55
2	B	503	3C4	CE2-CZ-CE1	-5.58	113.08	118.55
2	D	502	3C4	CE2-CZ-CE1	-5.20	113.46	118.55
2	A	505	3C4	CE1-CD1-CA	-4.85	117.18	120.44



There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	602	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	215/250 (86%)	0.63	19 (8%) 10 9	18, 25, 44, 52	0
1	B	218/250 (87%)	0.60	18 (8%) 11 10	18, 25, 33, 49	0
1	C	219/250 (87%)	0.64	19 (8%) 10 9	19, 27, 40, 47	0
1	D	220/250 (88%)	0.60	14 (6%) 19 17	18, 27, 42, 48	0
All	All	872/1000 (87%)	0.62	70 (8%) 12 12	18, 26, 42, 52	0

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	177	PRO	6.9
1	A	224	HIS	5.4
1	B	192	VAL	4.6
1	D	169	GLY	4.5
1	B	190	HIS	4.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	3C4	A	505	12/12	0.86	0.16	34,36,41,45	0
4	NA	A	701	1/1	0.96	0.08	32,32,32,32	0
3	PO4	D	602	5/5	0.96	0.26	49,50,51,52	0
3	PO4	C	603	5/5	0.97	0.27	49,51,52,54	0
2	3C4	B	503	12/12	0.98	0.07	14,16,19,19	0
3	PO4	B	601	5/5	0.98	0.19	36,37,40,41	0
2	3C4	D	502	12/12	0.98	0.08	13,18,19,20	0
2	3C4	A	501	12/12	0.98	0.08	14,16,19,19	0
2	3C4	C	504	12/12	0.99	0.06	13,18,19,20	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.