



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 14, 2018 – 11:11 PM EDT

PDB ID : 5E7F
Title : Complex between lactococcal phage Tuc2009 RBP head domain and a nanobody (L06)
Authors : Legrand, P.; Collins, B.; Blangy, S.; Murphy, J.; Spinelli, S.; Gutierrez, C.; Richet, N.; Kellenberger, C.; Desmyter, A.; Mahony, J.; van Sinderen, D.; Cambillau, C.
Deposited on : 2015-10-12
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	rb-20031633
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20031633

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	111664	2449 (2.70-2.70)
Clashscore	122126	2756 (2.70-2.70)
Ramachandran outliers	120053	2716 (2.70-2.70)
Sidechain outliers	120020	2716 (2.70-2.70)
RSRZ outliers	108989	2376 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	131	<div><div></div><div></div><div></div><div></div><div></div></div> <div>94%</div> <div>...</div>
1	B	131	<div><div></div><div></div><div></div><div></div><div></div></div> <div>15%</div> <div>91%</div> <div>.. 5%</div>
1	C	131	<div><div></div><div></div><div></div><div></div><div></div></div> <div>5%</div> <div>89%</div> <div>9% .</div>
2	G	174	<div><div></div><div></div><div></div><div></div><div></div></div> <div>3%</div> <div>63%</div> <div>10%</div> <div>26%</div>
2	H	174	<div><div></div><div></div><div></div><div></div><div></div></div> <div>7%</div> <div>64%</div> <div>9%</div> <div>26%</div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	I	174	 <p>7% 61% 11% 26%</p>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5883 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called nanobody L06.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	126	Total	C	N	O	S	0	0	0
			943	582	161	193	7			
1	B	125	Total	C	N	O	S	0	0	0
			933	576	158	192	7			
1	C	129	Total	C	N	O	S	0	0	0
			973	600	170	196	7			

- Molecule 2 is a protein called Major structural protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	128	Total	C	N	O	S	0	1	0
			998	639	163	191	5			
2	H	128	Total	C	N	O	S	0	1	0
			998	639	163	191	5			
2	I	128	Total	C	N	O	S	0	1	0
			998	639	163	191	5			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	174	ALA	-	expression tag	UNP Q38610
H	174	ALA	-	expression tag	UNP Q38610
I	174	ALA	-	expression tag	UNP Q38610

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	10	Total	O	0	0
			10	10		
3	B	3	Total	O	0	0
			3	3		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	11	Total 11	O 11	0	0
3	G	7	Total 7	O 7	0	0
3	H	8	Total 8	O 8	0	0
3	I	1	Total 1	O 1	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: nanobody L06




- Molecule 1: nanobody L06




- Molecule 1: nanobody L06




- Molecule 2: Major structural protein 1





- Molecule 2: Major structural protein 1





● Molecule 2: Major structural protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	84.73Å 88.04Å 147.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.80 – 2.70 34.53 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (34.80-2.70) 99.9 (34.53-2.70)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 2.68Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.194 , 0.208 0.207 , 0.217	Depositor DCC
R_{free} test set	1550 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	60.6	Xtriage
Anisotropy	0.744	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 55.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5883	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.39	0/964	0.62	0/1311
1	B	0.38	0/953	0.62	0/1296
1	C	0.41	0/997	0.64	0/1356
2	G	0.42	0/1030	0.66	0/1396
2	H	0.41	0/1030	0.65	0/1396
2	I	0.41	0/1030	0.66	0/1396
All	All	0.41	0/6004	0.64	0/8151

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	943	0	872	2	0
1	B	933	0	865	4	0
1	C	973	0	893	11	0
2	G	998	0	941	10	0
2	H	998	0	941	13	0
2	I	998	0	941	17	0
3	A	10	0	0	0	0
3	B	3	0	0	0	0
3	C	11	0	0	0	0
3	G	7	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	H	8	0	0	0	0
3	I	1	0	0	0	0
All	All	5883	0	5453	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:63[B]:TYR:HB3	2:I:63[B]:TYR:CZ	2.31	0.65
2:G:68:ALA:HB3	2:H:116:TYR:CD1	2.36	0.61
2:H:63[B]:TYR:HB3	2:I:63[B]:TYR:OH	2.06	0.56
2:G:125:HIS:HD2	2:G:128:GLN:O	1.90	0.55
2:I:125:HIS:HD2	2:I:128:GLN:O	1.90	0.54
2:H:125:HIS:HD2	2:H:128:GLN:O	1.91	0.54
2:H:68:ALA:HB3	2:I:116:TYR:CD1	2.43	0.53
2:G:63[B]:TYR:HB3	2:H:63[B]:TYR:CZ	2.43	0.53
1:A:106:SER:HB3	2:G:49:SER:HB2	1.91	0.52
1:C:54:PHE:HB3	2:I:58:LYS:HE3	1.93	0.51
2:G:63[B]:TYR:HB3	2:H:63[B]:TYR:OH	2.08	0.51
1:C:102:THR:OG1	2:I:70:LYS:HG3	2.13	0.48
2:I:142:LYS:HB2	2:I:145:LEU:HD12	1.97	0.47
1:C:103:THR:HG23	2:I:70:LYS:CB	2.45	0.47
2:I:76:VAL:HG23	2:I:166:MET:HB3	1.98	0.45
1:C:102:THR:HG21	2:I:68:ALA:O	2.16	0.45
1:B:106:SER:HB3	2:H:49:SER:HB2	1.99	0.45
2:G:142:LYS:HB2	2:G:145:LEU:HD12	1.99	0.45
1:C:103:THR:HG23	2:I:70:LYS:HB3	1.99	0.44
2:G:76:VAL:HG23	2:G:166:MET:HB3	1.98	0.44
1:B:49:LEU:HD23	2:H:57:LYS:HG3	1.98	0.44
1:C:36:MET:HB3	1:C:80:VAL:HG21	2.00	0.43
2:H:142:LYS:HB2	2:H:145:LEU:HD12	2.00	0.43
2:H:93:HIS:CE1	2:I:121:TYR:H	2.37	0.43
2:G:61:ASP:O	2:H:174:ALA:HB3	2.19	0.43
2:I:69:ASN:OD1	2:I:73:GLY:HA2	2.19	0.42
1:B:36:MET:HB3	1:B:80:VAL:HG21	2.01	0.42
1:C:24:CYS:O	1:C:79:THR:HA	2.20	0.42
2:H:76:VAL:HG23	2:H:166:MET:HB3	2.01	0.42
1:C:108:PRO:HA	2:I:49:SER:HA	2.01	0.42
2:G:116:TYR:CD1	2:I:68:ALA:HB3	2.54	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:63:ASP:HA	1:C:66:LYS:HG3	2.03	0.41
2:I:89:GLU:HG2	2:I:135:LYS:HG3	2.03	0.41
1:A:36:MET:HB3	1:A:80:VAL:HG21	2.02	0.41
2:G:89:GLU:HG2	2:G:135:LYS:HG3	2.03	0.41
1:B:83:GLN:HE21	1:B:85:ASN:ND2	2.19	0.40
1:C:102:THR:HB	2:I:69:ASN:HA	2.04	0.40
1:C:7:VAL:O	1:C:24:CYS:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	124/131 (95%)	122 (98%)	2 (2%)	0	100	100
1	B	123/131 (94%)	120 (98%)	3 (2%)	0	100	100
1	C	127/131 (97%)	124 (98%)	3 (2%)	0	100	100
2	G	127/174 (73%)	125 (98%)	2 (2%)	0	100	100
2	H	127/174 (73%)	124 (98%)	3 (2%)	0	100	100
2	I	127/174 (73%)	125 (98%)	2 (2%)	0	100	100
All	All	755/915 (82%)	740 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	102/107 (95%)	101 (99%)	1 (1%)	78	92
1	B	101/107 (94%)	100 (99%)	1 (1%)	78	92
1	C	105/107 (98%)	104 (99%)	1 (1%)	78	92
2	G	102/139 (73%)	97 (95%)	5 (5%)	27	56
2	H	102/139 (73%)	99 (97%)	3 (3%)	45	75
2	I	102/139 (73%)	97 (95%)	5 (5%)	27	56
All	All	614/738 (83%)	598 (97%)	16 (3%)	49	79

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	106	SER
1	B	106	SER
1	C	106	SER
2	G	70	LYS
2	G	82	TRP
2	G	122	THR
2	G	151	ILE
2	G	162	LYS
2	H	82	TRP
2	H	122	THR
2	H	151	ILE
2	I	61	ASP
2	I	69	ASN
2	I	82	TRP
2	I	122	THR
2	I	151	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	85	ASN
1	B	85	ASN
1	C	85	ASN
1	C	130	HIS
2	G	93	HIS
2	G	125	HIS
2	H	93	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	125	HIS
2	I	93	HIS
2	I	125	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	126/131 (96%)	-0.02	1 (0%) 86 87	56, 67, 87, 122	0
1	B	125/131 (95%)	0.47	19 (15%) 2 1	64, 94, 126, 170	0
1	C	129/131 (98%)	-0.03	6 (4%) 31 29	51, 66, 101, 125	0
2	G	128/174 (73%)	0.19	5 (3%) 39 38	54, 69, 102, 111	0
2	H	128/174 (73%)	0.47	12 (9%) 8 6	54, 73, 109, 133	0
2	I	128/174 (73%)	0.40	13 (10%) 7 5	64, 82, 110, 127	0
All	All	764/915 (83%)	0.25	56 (7%) 15 13	51, 75, 113, 170	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	97	TYR	5.4
2	I	63[A]	TYR	5.0
2	H	97	TYR	5.0
1	B	126	SER	4.9
1	B	3	GLN	4.7
1	C	3	GLN	4.7
2	G	63[A]	TYR	4.5
1	B	77	LYS	4.5
2	I	157	ARG	4.4
2	H	157	ARG	4.3
1	B	76	ALA	4.2
2	H	63[A]	TYR	4.2
1	C	43	PRO	4.2
2	H	98	SER	4.1
1	B	78	ASN	3.5
2	I	155	ASN	3.5
2	I	127	ASN	3.3
1	B	17	GLY	3.3
2	H	127	ASN	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	67	GLY	3.3
1	B	127	SER	3.2
2	I	159	GLY	3.2
2	H	159	GLY	3.2
1	B	33	ASP	3.2
2	H	155	ASN	3.1
2	G	97	TYR	3.0
1	B	19	SER	3.0
2	I	110	SER	3.0
1	C	44	GLY	3.0
2	H	174	ALA	2.9
2	I	156	GLY	2.9
2	I	98	SER	2.9
2	G	98	SER	2.7
1	B	32	ASP	2.7
1	C	130	HIS	2.5
1	B	75	ASN	2.5
2	G	155	ASN	2.5
2	I	111	GLY	2.5
1	A	128	HIS	2.5
1	C	42	ALA	2.4
1	B	16	ALA	2.4
1	B	86	SER	2.4
2	H	154	ALA	2.3
2	I	72	SER	2.3
2	H	70	LYS	2.3
2	I	109	PRO	2.2
1	B	68	ARG	2.2
1	B	74	ASP	2.2
1	B	89	PRO	2.2
1	B	66	LYS	2.1
1	B	29	PHE	2.1
2	H	156	GLY	2.1
2	G	100	GLY	2.0
1	C	131	HIS	2.0
2	H	96	GLY	2.0
2	I	158	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.