



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 21, 2019 – 12:16 PM EST

PDB ID : 6E7X
Title : Heterodimer of the GluN1b-GluN2B NMDA receptor amino-terminal domains bound to allosteric inhibitor 93-97
Authors : Regan, M.C.; Furukawa, H.
Deposited on : 2018-07-27
Resolution : 2.58 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : rb-20031633
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20031633


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	3182 (2.60-2.56)
Clashscore	122126	3541 (2.60-2.56)
Ramachandran outliers	120053	3489 (2.60-2.56)
Sidechain outliers	120020	3489 (2.60-2.56)
RSRZ outliers	108989	3120 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	385	<div> <div>2%</div> <div>77%</div> <div>15%</div> <div>7%</div> </div>
1	C	385	<div> <div>4%</div> <div>76%</div> <div>17%</div> <div>7%</div> </div>
2	B	363	<div> <div>5%</div> <div>85%</div> <div>13%</div> <div>•</div> </div>
2	D	363	<div> <div>9%</div> <div>85%</div> <div>14%</div> <div>•</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	C	502	-	-	-	X
3	NAG	D	502	-	-	-	X
7	CL	A	510	-	-	X	-
7	CL	B	504	-	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 11406 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate receptor ionotropic, NMDA 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	358	Total	C	N	O	S	0	0	0
			2750	1756	478	505	11			
1	C	357	Total	C	N	O	S	0	0	0
			2692	1718	461	502	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	GLN	ASN	engineered mutation	UNP A0A1L8F5J9
A	371	GLN	ASN	engineered mutation	UNP A0A1L8F5J9
C	61	GLN	ASN	engineered mutation	UNP A0A1L8F5J9
C	371	GLN	ASN	engineered mutation	UNP A0A1L8F5J9


- Molecule 2 is a protein called Glutamate receptor ionotropic, NMDA 2B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	363	Total	C	N	O	S	0	0	0
			2774	1788	438	533	15			
2	D	363	Total	C	N	O	S	0	0	0
			2732	1758	439	519	16			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	348	ASP	ASN	engineered mutation	UNP Q00960
D	348	ASP	ASN	engineered mutation	UNP Q00960

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			13	7	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is BETA-D-MANNOSE (three-letter code: BMA) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		


- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Na 1 1	0	0
6	C	1	Total Na 1 1	0	0

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	3	Total Cl 3 3	0	0
7	A	2	Total Cl 2 2	0	0
7	D	2	Total Cl 2 2	0	0
7	C	2	Total Cl 2 2	0	0

- Molecule 8 is N-{4-[(2S)-3-{[2-(3,4-dichlorophenyl)ethyl](2-methylpropyl)amino}-2-hydroxypropoxy]phenyl}methanesulfonamide (three-letter code: T97) (formula: C₂₂H₃₀Cl₂N₂O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total C Cl N O S 31 22 2 2 4 1	0	0
8	D	1	Total C Cl N O S 31 22 2 2 4 1	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	97	Total 97	O 97	0	0
9	B	58	Total 58	O 58	0	0
9	C	52	Total 52	O 52	0	0
9	D	34	Total 34	O 34	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamate receptor ionotropic, NMDA 1




- Molecule 1: Glutamate receptor ionotropic, NMDA 1



- Molecule 2: Glutamate receptor ionotropic, NMDA 2B



● Molecule 2: Glutamate receptor ionotropic, NMDA 2B



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	268.42Å 59.54Å 146.11Å 90.00° 116.69° 90.00°	Depositor
Resolution (Å)	25.00 – 2.58 34.39 – 2.58	Depositor EDS
% Data completeness (in resolution range)	85.6 (25.00-2.58) 85.8 (34.39-2.58)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.97 (at 2.57Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.185 , 0.216 0.190 , 0.218	Depositor DCC
R_{free} test set	2782 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	42.6	Xtriage
Anisotropy	0.208	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11406	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, CL, NA, T97, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/2808	0.35	0/3818
1	C	0.21	0/2748	0.35	0/3746
2	B	0.21	0/2836	0.36	0/3871
2	D	0.21	0/2796	0.36	0/3823
All	All	0.21	0/11188	0.35	0/15258

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	D	0	3
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	57	ASP	Peptide
2	D	333	ASN	Peptide
2	D	55	GLU	Peptide
2	D	61	HIS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2750	0	2711	39	1
1	C	2692	0	2613	53	0
2	B	2774	0	2612	49	0
2	D	2732	0	2551	50	1
3	A	42	0	37	0	0
3	B	28	0	26	0	0
3	C	27	0	22	4	0
3	D	14	0	13	0	0
4	A	11	0	8	0	0
5	A	22	0	20	0	0
6	A	1	0	0	0	0
6	C	1	0	0	0	0
7	A	2	0	0	4	0
7	B	3	0	0	3	0
7	C	2	0	0	0	0
7	D	2	0	0	1	0
8	B	31	0	0	3	0
8	D	31	0	0	3	0
9	A	97	0	0	1	0
9	B	58	0	0	3	0
9	C	52	0	0	4	0
9	D	34	0	0	1	0
All	All	11406	0	10613	195	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 195 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:501:NAG:C8	3:C:501:NAG:N2	1.97	1.28
2:B:235:GLU:OE1	2:B:235:GLU:OE2	1.74	1.05
3:C:501:NAG:O7	3:C:501:NAG:N2	1.95	1.00
2:D:54:HIS:O	2:D:57:ASP:CB	2.23	0.87
2:D:110:GLN:HG2	8:D:503:T97:C07	2.08	0.83

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:380:ARG:O	2:D:393:ARG:NH1[2_557]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/385 (91%)	338 (96%)	13 (4%)	1 (0%)	43	65
1	C	351/385 (91%)	340 (97%)	11 (3%)	0	100	100
2	B	361/363 (99%)	346 (96%)	11 (3%)	4 (1%)	16	31
2	D	361/363 (99%)	344 (95%)	14 (4%)	3 (1%)	21	40
All	All	1425/1496 (95%)	1368 (96%)	49 (3%)	8 (1%)	27	49

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	185	GLU
2	B	58	ASP
2	B	92	ARG
2	D	92	ARG
2	D	284	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/331 (87%)	281 (98%)	6 (2%)	56	78
1	C	278/331 (84%)	276 (99%)	2 (1%)	85	94
2	B	288/326 (88%)	283 (98%)	5 (2%)	63	82
2	D	280/326 (86%)	276 (99%)	4 (1%)	69	86
All	All	1133/1314 (86%)	1116 (98%)	17 (2%)	67	85

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	188	SER
2	B	287	TYR
2	D	110	GLN
2	B	134	MET
2	D	133	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	321	ASN
1	C	384	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 11 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	501	1	14,14,15	0.29	0	17,19,21	0.69	0
3	NAG	A	502	1,3	14,14,15	0.27	0	17,19,21	0.74	0
3	NAG	A	503	3,4	14,14,15	0.30	0	17,19,21	0.85	1 (5%)
4	BMA	A	504	3,5	11,11,12	0.24	0	15,15,17	0.59	0
5	MAN	A	505	4	11,11,12	0.24	0	15,15,17	0.61	0
5	MAN	A	506	4	11,11,12	0.23	0	15,15,17	0.61	0
3	NAG	B	501	2	14,14,15	0.26	0	17,19,21	0.69	0
3	NAG	B	502	2	14,14,15	0.38	0	17,19,21	1.01	1 (5%)
8	T97	B	503	-	32,32,32	1.44	4 (12%)	44,44,44	2.02	4 (9%)
3	NAG	C	501	1	11,11,15	0.35	0	13,15,21	0.50	0
3	NAG	C	502	1	14,14,15	0.27	0	17,19,21	1.31	1 (5%)
3	NAG	D	502	2	14,14,15	0.33	0	17,19,21	0.66	0
8	T97	D	503	-	32,32,32	1.44	4 (12%)	44,44,44	2.00	4 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	501	1	-	0/6/23/26	0/1/1/1
3	NAG	A	502	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	503	3,4	-	0/6/23/26	0/1/1/1
4	BMA	A	504	3,5	-	0/2/19/22	0/1/1/1
5	MAN	A	505	4	-	0/2/19/22	0/1/1/1
5	MAN	A	506	4	-	0/2/19/22	0/1/1/1
3	NAG	B	501	2	-	0/6/23/26	0/1/1/1
3	NAG	B	502	2	-	0/6/23/26	0/1/1/1
8	T97	B	503	-	-	0/23/23/23	0/2/2/2
3	NAG	C	501	1	-	0/2/19/26	0/1/1/1
3	NAG	C	502	1	-	0/6/23/26	0/1/1/1
3	NAG	D	502	2	-	0/6/23/26	0/1/1/1
8	T97	D	503	-	-	0/23/23/23	0/2/2/2

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	B	503	T97	C01-CL2	2.08	1.78	1.73
8	D	503	T97	C01-CL2	2.18	1.78	1.73
8	B	503	T97	C02-CL1	2.18	1.78	1.73
8	D	503	T97	C02-CL1	2.26	1.79	1.73
8	D	503	T97	C26-S25	2.64	1.82	1.75

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	503	T97	O28-S25-O27	-11.61	101.41	118.79
8	D	503	T97	O28-S25-O27	-11.50	101.58	118.79
8	B	503	T97	C11-C10-N09	-2.52	110.53	116.00
8	D	503	T97	C11-C10-N09	-2.46	110.65	116.00
3	A	503	NAG	O5-C1-C2	-2.05	108.70	111.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	503	T97	3	0
3	C	501	NAG	4	0
8	D	503	T97	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	358/385 (92%)	-0.19	7 (1%) 65 61	21, 37, 69, 104	0
1	C	357/385 (92%)	0.08	14 (3%) 39 34	31, 53, 86, 129	0
2	B	363/363 (100%)	0.25	19 (5%) 27 23	22, 48, 86, 128	0
2	D	363/363 (100%)	0.31	33 (9%) 9 7	30, 53, 98, 141	0
All	All	1441/1496 (96%)	0.11	73 (5%) 28 24	21, 48, 86, 141	0

The worst 5 of 73 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	344	PHE	6.1
2	B	58	ASP	5.8
2	B	48	VAL	5.6
2	B	55	GLU	5.3
2	D	57	ASP	5.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	B	501	14/15	0.70	0.31	76,91,100,101	0
3	NAG	D	502	14/15	0.76	0.41	97,103,106,109	0
5	MAN	A	506	11/12	0.77	0.30	110,119,124,127	0
3	NAG	C	502	14/15	0.78	0.44	99,105,114,116	0
3	NAG	B	502	14/15	0.78	0.39	98,116,129,133	0
7	CL	D	504	1/1	0.83	0.33	72,72,72,72	0
7	CL	D	505	1/1	0.86	0.15	78,78,78,78	0
3	NAG	C	501	13/15	0.90	0.19	51,60,66,67	0
8	T97	B	503	31/31	0.91	0.22	29,50,64,70	0
7	CL	B	504	1/1	0.92	0.15	59,59,59,59	0
3	NAG	A	503	14/15	0.93	0.18	61,68,73,77	0
4	BMA	A	504	11/12	0.93	0.25	73,81,93,115	0
3	NAG	A	502	14/15	0.94	0.17	50,56,57,62	0
3	NAG	A	501	14/15	0.94	0.18	42,53,65,70	0
5	MAN	A	505	11/12	0.95	0.18	57,64,69,71	0
6	NA	A	508	1/1	0.95	0.16	32,32,32,32	0
7	CL	B	505	1/1	0.95	0.18	74,74,74,74	0
8	T97	D	503	31/31	0.95	0.14	38,50,65,70	0
7	CL	B	506	1/1	0.95	0.11	64,64,64,64	0
7	CL	C	504	1/1	0.95	0.20	76,76,76,76	0
7	CL	C	505	1/1	0.96	0.11	64,64,64,64	0
7	CL	A	510	1/1	0.96	0.09	48,48,48,48	0
7	CL	A	509	1/1	0.97	0.09	66,66,66,66	0
6	NA	C	503	1/1	0.97	0.06	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.