



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 11:08 am GMT

PDB ID : 4E8F
Title : Structural Basis for the Activity of a Cytoplasmic RNA Terminal U-transferase
Authors : Yates, L.A.; Fleurdepine, S.; Rissland, O.S.; DeColibus, L.; Harlos, K.; Norbury, C.J.; Gilbert, R.J.C.
Deposited on : 2012-03-20
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2767 (2.60-2.60)
Clashscore	122126	3110 (2.60-2.60)
Ramachandran outliers	120053	3062 (2.60-2.60)
Sidechain outliers	120020	3062 (2.60-2.60)
RSRZ outliers	108989	2706 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	405	<div> <div></div> <div> <div></div> <div>73%</div> <div>7%</div> <div>19%</div> </div> </div>
1	B	405	<div> <div>5%</div> <div> <div></div> <div>65%</div> <div>11%</div> <div>21%</div> </div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5297 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Poly(A) RNA polymerase protein cid1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	327	Total	C	N	O	S	0	0	0
			2638	1709	438	482	9			
1	B	319	Total	C	N	O	S	0	0	0
			2564	1665	424	466	9			

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	49	Total	O	0	0
			49	49		
4	B	32	Total	O	0	0
			32	32		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Poly(A) RNA polymerase protein cid1



- Molecule 1: Poly(A) RNA polymerase protein cid1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.15Å 101.21Å 113.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.79 – 2.60 56.78 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.5 (56.79-2.60) 99.4 (56.78-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.64 (at 2.58Å)	Xtriage
Refinement program	BUSTER 2.9.2	Depositor
R, R_{free}	0.192 , 0.239 0.196 , 0.243	Depositor DCC
R_{free} test set	1494 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	57.4	Xtriage
Anisotropy	0.643	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 49.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5297	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/2701	0.59	0/3645
1	B	0.47	0/2625	0.63	0/3545
All	All	0.46	0/5326	0.61	0/7190

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2638	0	2642	12	0
1	B	2564	0	2555	21	0
2	A	4	0	3	0	0
2	B	4	0	3	0	0
3	B	6	0	8	0	0
4	A	49	0	0	1	0
4	B	32	0	0	0	0
All	All	5297	0	5211	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:ILE:HD11	1:A:124:GLU:HB2	1.80	0.63
1:A:266:SER:HB2	1:A:271:LEU:HD13	1.82	0.62
1:B:73:ARG:HD2	1:B:85:LEU:HB3	1.81	0.61
1:B:266:SER:HB2	1:B:271:LEU:HD13	1.84	0.59
1:A:228:PRO:HB2	1:A:263:GLN:HG3	1.85	0.58
1:B:98:LYS:HE3	1:B:102:MET:HG2	1.86	0.57
1:A:293:ARG:HE	1:A:334:ILE:HD11	1.70	0.56
1:B:137:ARG:HB2	1:B:205:TYR:HD2	1.72	0.55
1:B:293:ARG:HE	1:B:334:ILE:HD11	1.72	0.53
1:B:100:SER:HB2	1:B:205:TYR:HE1	1.74	0.53
1:B:88:PHE:HB3	1:B:95:LEU:HB3	1.90	0.53
1:B:79:ILE:HD11	1:B:121:PHE:HA	1.92	0.52
1:A:263:GLN:HG2	4:A:636:HOH:O	2.10	0.51
1:B:100:SER:HB2	1:B:205:TYR:CE1	2.49	0.48
1:A:372:SER:HA	1:A:375:GLU:HG3	1.97	0.46
1:B:99:ASN:O	1:B:204:PRO:HG2	2.15	0.46
1:B:92:GLU:C	1:B:94:GLY:H	2.18	0.46
1:A:256:LEU:HA	1:A:259:ILE:HD12	1.99	0.45
1:B:79:ILE:CD1	1:B:121:PHE:HA	2.47	0.44
1:B:140:ILE:HD11	1:B:341:THR:HG21	1.98	0.44
1:B:193:LYS:HE2	1:B:211:SER:HB3	1.99	0.44
1:A:101:ASP:OD1	1:A:158:GLN:HB2	2.18	0.43
1:B:75:CYS:SG	1:B:128:GLU:HG3	2.59	0.43
1:A:79:ILE:H	1:A:79:ILE:HG12	1.74	0.42
1:B:256:LEU:HA	1:B:259:ILE:HD12	2.01	0.42
1:B:90:SER:HB2	1:B:94:GLY:HA3	2.01	0.42
1:B:97:LEU:HA	1:B:212:TYR:HB2	2.02	0.42
1:A:73:ARG:HG2	1:A:85:LEU:HB3	2.02	0.41
1:B:347:LEU:O	1:B:351:ARG:HB2	2.21	0.41
1:A:193:LYS:HE2	1:A:211:SER:HB3	2.04	0.40
1:B:337:ASN:O	1:B:340:ARG:HG2	2.21	0.40
1:A:79:ILE:HD11	1:A:124:GLU:CB	2.49	0.40
1:B:91:LEU:HA	1:B:92:GLU:HA	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/405 (79%)	314 (98%)	6 (2%)	1 (0%)	43	68
1	B	309/405 (76%)	295 (96%)	12 (4%)	2 (1%)	27	51
All	All	630/810 (78%)	609 (97%)	18 (3%)	3 (0%)	31	56

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	115	ASP
1	B	149	THR
1	B	94	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	290/363 (80%)	274 (94%)	16 (6%)	24	47
1	B	280/363 (77%)	254 (91%)	26 (9%)	10	19
All	All	570/726 (78%)	528 (93%)	42 (7%)	15	30

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	SER
1	A	57	ILE
1	A	59	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	79	ILE
1	A	100	SER
1	A	103	ASP
1	A	139	ARG
1	A	140	ILE
1	A	153	PHE
1	A	181	LEU
1	A	190	LEU
1	A	197	LYS
1	A	271	LEU
1	A	303	GLU
1	A	334	ILE
1	A	371	ASP
1	B	57	ILE
1	B	72	LEU
1	B	74	LEU
1	B	79	ILE
1	B	91	LEU
1	B	92	GLU
1	B	95	LEU
1	B	97	LEU
1	B	99	ASN
1	B	107	LEU
1	B	108	MET
1	B	116	THR
1	B	142	ILE
1	B	148	ASP
1	B	150	LYS
1	B	158	GLN
1	B	164	ASN
1	B	175	LEU
1	B	181	LEU
1	B	190	LEU
1	B	263	GLN
1	B	271	LEU
1	B	287	GLU
1	B	303	GLU
1	B	334	ILE
1	B	371	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	171	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	A	500	-	1,3,3	5.01	1 (100%)	0,3,3	0.00	-
3	GOL	B	500	-	5,5,5	0.10	0	5,5,5	0.18	0
2	ACT	B	501	-	1,3,3	4.84	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACT	A	500	-	-	0/0/0/0	0/0/0/0
3	GOL	B	500	-	-	0/4/4/4	0/0/0/0
2	ACT	B	501	-	-	0/0/0/0	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	ACT	CH3-C	4.84	1.54	1.48
2	A	500	ACT	CH3-C	5.01	1.55	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	327/405 (80%)	0.08	5 (1%) 73 69	40, 66, 111, 142	0
1	B	319/405 (78%)	0.48	22 (6%) 17 12	46, 75, 123, 144	0
All	All	646/810 (79%)	0.28	27 (4%) 36 28	40, 71, 120, 144	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	98	LYS	7.0
1	B	96	ALA	6.6
1	B	97	LEU	6.5
1	B	91	LEU	5.8
1	B	100	SER	4.8
1	B	93	SER	4.6
1	A	147	SER	4.2
1	B	92	GLU	4.0
1	B	101	ASP	3.9
1	B	95	LEU	3.7
1	B	137	ARG	3.5
1	B	205	TYR	3.5
1	A	157	PHE	3.5
1	B	99	ASN	3.3
1	B	152	GLY	3.2
1	B	248	PHE	3.0
1	A	146	THR	3.0
1	B	245	VAL	3.0
1	B	380	PRO	2.8
1	B	169	ILE	2.7
1	B	88	PHE	2.6
1	B	90	SER	2.5
1	B	145	LEU	2.5
1	A	135	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	308	SER	2.1
1	A	156	SER	2.1
1	B	149	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	500	6/6	0.82	0.19	89,91,94,95	0
2	ACT	B	501	4/4	0.83	0.85	102,105,105,105	0
2	ACT	A	500	4/4	0.84	0.33	86,86,86,87	0

6.5 Other polymers [i](#)

There are no such residues in this entry.