



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2020 – 06:46 AM EST

PDB ID : 3EDK  
Title : Structural base for cyclodextrin hydrolysis  
Authors : Buedenbender, S.; Schulz, G.E.  
Deposited on : 2008-09-03  
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : 2.8  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.8


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	7886 (1.80-1.76)
Clashscore	122126	8858 (1.80-1.76)
Ramachandran outliers	120053	8764 (1.80-1.76)
Sidechain outliers	120020	8763 (1.80-1.76)
RSRZ outliers	108989	7741 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	601	<div> <div>5%</div> <div> <div></div> <div>92%</div> <div>7%</div> <div>.</div> </div> </div>
1	B	601	<div> <div>6%</div> <div> <div></div> <div>88%</div> <div>11%</div> <div>..</div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11035 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclomaltoextrinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	597	Total	C	N	O	S	0	21	0
			4852	3063	875	890	24			
1	B	597	Total	C	N	O	S	0	25	0
			4874	3076	883	890	25			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	49	PRO	THR	ENGINEERED	UNP Q8KKG0
A	340	GLN	GLU	ENGINEERED	UNP Q8KKG0
B	49	PRO	THR	ENGINEERED	UNP Q8KKG0
B	340	GLN	GLU	ENGINEERED	UNP Q8KKG0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Ca	0	0
			2	2		
2	A	2	Total	Ca	0	0
			2	2		

- Molecule 3 is gamma-cyclodextrin (three-letter code: RCD) (formula: C<sub>48</sub>H<sub>80</sub>O<sub>40</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 88	C 48	O 40	0	0
3	B	1	Total 88	C 48	O 40	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $\text{C}_3\text{H}_8\text{O}_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	584	Total	O	0	0
			584	584		
5	B	521	Total	O	0	0
			521	521		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Cyclomaltodextrinase



#### • Molecule 1: Cyclomaltodextrinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.14Å 110.79Å 106.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.71 – 1.77 19.49 – 1.77	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.71-1.77) 99.9 (19.49-1.77)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.88 (at 1.77Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.175 , 0.208 0.172 , 0.205	Depositor DCC
$R_{free}$ test set	2445 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.6	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 50.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.012 for -h,l,k 0.012 for -k,-h,l 0.011 for l,-k,h 0.001 for l,h,k 0.001 for k,l,h	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	11035	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RCD, GOL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.68	0/5084	0.69	0/6890
1	B	0.68	6/5119 (0.1%)	0.79	12/6933 (0.2%)
All	All	0.68	6/10203 (0.1%)	0.74	12/13823 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	341	GLU	CD-OE1	-8.07	1.16	1.25
1	B	406[A]	GLU	CG-CD	6.14	1.61	1.51
1	B	406[B]	GLU	CG-CD	6.14	1.61	1.51
1	B	253	ARG	CZ-NH1	-5.72	1.25	1.33
1	B	406[A]	GLU	CB-CG	5.03	1.61	1.52
1	B	406[B]	GLU	CB-CG	5.03	1.61	1.52

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	470[A]	ARG	NE-CZ-NH2	14.57	127.59	120.30
1	B	470[B]	ARG	NE-CZ-NH2	14.57	127.59	120.30
1	B	470[A]	ARG	NE-CZ-NH1	-10.96	114.82	120.30
1	B	470[B]	ARG	NE-CZ-NH1	-10.96	114.82	120.30
1	B	194	ARG	NE-CZ-NH1	-9.27	115.67	120.30
1	B	253	ARG	NE-CZ-NH2	8.85	124.73	120.30
1	B	93	ARG	NE-CZ-NH1	-7.67	116.47	120.30
1	B	93	ARG	NE-CZ-NH2	6.40	123.50	120.30
1	B	470[A]	ARG	CD-NE-CZ	6.31	132.44	123.60
1	B	470[B]	ARG	CD-NE-CZ	6.31	132.44	123.60
1	B	194	ARG	NE-CZ-NH2	5.93	123.27	120.30
1	B	253	ARG	NE-CZ-NH1	-5.46	117.57	120.30



There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4852	0	4655	28	0
1	B	4874	0	4675	61	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	88	0	80	2	0
3	B	88	0	80	2	0
4	A	12	0	16	0	0
4	B	12	0	16	0	0
5	A	584	0	0	7	0
5	B	521	0	0	10	0
All	All	11035	0	9522	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:100:ARG:NH1	1:B:300:ILE:HG22	1.69	1.07
1:B:100:ARG:CZ	5:B:1268:HOH:O	2.05	1.04
1:B:113:MET:SD	1:B:470[B]:ARG:CD	2.61	0.88
1:B:100:ARG:HH11	1:B:300:ILE:HG22	1.35	0.86
1:B:138:HIS:HD2	1:B:140:GLY:H	1.23	0.86
1:B:434[B]:ARG:NH2	5:B:1163:HOH:O	2.08	0.85
1:B:74:ASP:OD1	1:B:87[A]:ARG:HG2	1.79	0.82
1:B:113:MET:SD	1:B:470[B]:ARG:HD2	2.20	0.81
1:B:417:HIS:O	1:B:470[A]:ARG:HD2	1.82	0.80
1:B:253:ARG:NH2	5:B:1096:HOH:O	2.16	0.79
1:B:181:HIS:HD2	1:B:183:TYR:H	1.29	0.78
1:A:181:HIS:HD2	1:A:183:TYR:H	1.31	0.77
1:B:100:ARG:HH11	1:B:300:ILE:CG2	1.97	0.77

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:47:VAL:O	1:A:48:SER:HB3	1.85	0.75
1:B:113:MET:SD	1:B:470[A]:ARG:HG2	2.26	0.75
1:B:341:GLU:OE2	5:B:1096:HOH:O	2.04	0.74
1:B:93:ARG:HD2	1:B:301:GLU:OE2	1.89	0.73
1:B:100:ARG:NH1	5:B:1268:HOH:O	2.14	0.71
1:A:353:GLN:HE21	1:A:405:PRO:HD2	1.57	0.70
1:B:8:HIS:HD2	1:B:28:HIS:ND1	1.90	0.69
1:A:8:HIS:HD2	1:A:28:HIS:ND1	1.91	0.68
1:B:353:GLN:HE21	1:B:405:PRO:HD2	1.58	0.68
1:B:100:ARG:HH12	1:B:300:ILE:HG22	1.60	0.64
1:A:251:HIS:HD2	1:A:315:TYR:OH	1.81	0.64
1:B:251:HIS:HD2	1:B:315:TYR:OH	1.80	0.63
1:A:252:HIS:HE1	1:B:525:GLU:OE2	1.82	0.63
1:A:525:GLU:OE2	1:B:252:HIS:HE1	1.83	0.62
1:A:47:VAL:O	1:A:48:SER:CB	2.47	0.62
1:B:113:MET:SD	1:B:470[B]:ARG:HD3	2.39	0.61
1:B:155:ASP:OD2	1:B:211:ARG:HD3	2.00	0.61
1:A:419:MET:HE3	5:A:1280:HOH:O	2.01	0.60
1:B:192:ASP:OD2	1:B:194:ARG:HD2	2.01	0.60
1:B:138:HIS:CD2	1:B:140:GLY:H	2.12	0.60
1:A:495[A]:ARG:NH2	5:A:1075:HOH:O	2.35	0.59
1:B:93:ARG:CD	1:B:301:GLU:OE2	2.50	0.59
1:B:192:ASP:OD2	1:B:194:ARG:CD	2.51	0.58
1:B:461:VAL:HG23	1:B:462[B]:LYS:HG2	1.84	0.58
1:B:138:HIS:HD2	1:B:140:GLY:N	1.98	0.58
1:B:3:PRO:HB3	1:B:7:GLU:HG3	1.85	0.58
1:A:78:LYS:HG2	1:A:83:SER:HB2	1.86	0.58
1:A:181:HIS:CD2	1:A:183:TYR:H	2.17	0.57
1:A:252:HIS:HD2	5:B:855:HOH:O	1.88	0.56
1:A:263[B]:GLN:NE2	5:A:1231:HOH:O	2.38	0.56
5:A:849:HOH:O	1:B:252:HIS:HD2	1.90	0.55
1:B:200:ASP:OD1	1:B:203[A]:ARG:NH2	2.36	0.55
1:B:384:LYS:HB3	1:B:387:GLU:HG3	1.87	0.54
1:B:181:HIS:CD2	1:B:183:TYR:H	2.18	0.54
1:B:431:ASP:OD1	1:B:434[B]:ARG:NH1	2.41	0.54
1:B:253:ARG:NH1	3:B:700:RCD:O2E	2.41	0.54
1:A:495[A]:ARG:NE	5:A:892:HOH:O	2.40	0.53
1:B:100:ARG:NH1	1:B:300:ILE:CG2	2.53	0.53
1:A:202:VAL:O	1:A:206[B]:THR:HG23	2.09	0.52
1:B:173:ASN:O	1:B:179:SER:HB2	2.09	0.52
1:B:181:HIS:HE1	5:B:1067:HOH:O	1.91	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:353:GLN:NE2	1:A:406:GLU:H	2.07	0.52
1:A:511:GLN:HE22	1:A:538:LYS:HZ2	1.58	0.51
1:A:24:GLN:NE2	1:A:290:ASN:HD22	2.09	0.51
1:B:417:HIS:O	1:B:470[B]:ARG:NH1	2.44	0.50
1:B:24:GLN:HE22	1:B:290:ASN:HD22	1.60	0.50
1:B:466:ASP:O	1:B:470[B]:ARG:HG3	2.11	0.50
1:A:463:GLY:HA2	3:A:700:RCD:O3E	2.13	0.49
1:B:466:ASP:O	1:B:470[A]:ARG:HD3	2.12	0.49
1:B:538:LYS:HZ1	1:B:594[B]:GLU:CD	2.16	0.49
1:B:398:LEU:HD11	1:B:443:MET:SD	2.53	0.49
3:B:700:RCD:H6EA	3:B:700:RCD:H3D	1.96	0.48
1:B:19:GLN:O	1:B:21:LYS:HE2	2.14	0.47
1:A:173:ASN:O	1:A:179:SER:HB2	2.14	0.47
1:A:569:VAL:HG22	1:A:576[A]:THR:HG22	1.97	0.47
1:A:393:GLU:HG3	5:A:1009:HOH:O	2.15	0.46
1:B:24:GLN:NE2	1:B:290:ASN:HD22	2.14	0.46
1:B:194:ARG:NH1	5:B:961:HOH:O	2.39	0.45
1:B:538:LYS:HD3	1:B:594[B]:GLU:HG3	1.97	0.45
1:A:511:GLN:HE22	1:A:538:LYS:NZ	2.14	0.44
1:B:112:ILE:HG22	1:B:451:SER:HB2	1.99	0.44
1:A:507:TRP:O	1:A:511:GLN:HG2	2.17	0.44
1:B:192:ASP:OD2	1:B:194:ARG:HD3	2.18	0.42
1:B:538:LYS:NZ	1:B:594[B]:GLU:CD	2.74	0.41
1:A:344[B]:THR:HG22	5:A:1151:HOH:O	2.20	0.41
1:B:47:VAL:HB	1:B:61:ASP:HB2	2.03	0.41
1:B:576[A]:THR:HG23	5:B:1104:HOH:O	2.20	0.41
3:A:700:RCD:H1D	3:A:700:RCD:O3E	2.20	0.41
1:B:242:TYR:O	1:B:245:LYS:HG2	2.21	0.40
1:B:426:ALA:HB2	5:B:931:HOH:O	2.22	0.40
1:A:99:GLN:CD	1:A:334:ARG:HE	2.25	0.40
1:A:246:PHE:CD1	1:A:276:GLU:HA	2.56	0.40
1:B:207:GLU:OE1	1:B:211:ARG:NH2	2.54	0.40
1:B:234:LEU:HA	1:B:235:PRO:HD3	1.98	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	616/601 (102%)	599 (97%)	16 (3%)	1 (0%)	49	32
1	B	620/601 (103%)	607 (98%)	13 (2%)	0	100	100
All	All	1236/1202 (103%)	1206 (98%)	29 (2%)	1 (0%)	53	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	48	SER

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	513/493 (104%)	504 (98%)	9 (2%)	62	48
1	B	517/493 (105%)	503 (97%)	14 (3%)	48	31
All	All	1030/986 (104%)	1007 (98%)	23 (2%)	56	39

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	THR
1	A	51	ARG
1	A	137	ARG
1	A	253[A]	ARG
1	A	253[B]	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	436	ASN
1	A	448	GLN
1	A	521	HIS
1	A	581	ARG
1	B	4	THR
1	B	46	LEU
1	B	51	ARG
1	B	87[A]	ARG
1	B	87[B]	ARG
1	B	90	LEU
1	B	194	ARG
1	B	369	LEU
1	B	436	ASN
1	B	448	GLN
1	B	455	ILE
1	B	521	HIS
1	B	584	ARG
1	B	598	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (32) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	HIS
1	A	24	GLN
1	A	111	GLN
1	A	138	HIS
1	A	181	HIS
1	A	217	GLN
1	A	241	ASN
1	A	251	HIS
1	A	252	HIS
1	A	257	GLN
1	A	296	ASN
1	A	353	GLN
1	A	358	ASN
1	A	448	GLN
1	A	498	GLN
1	A	511	GLN
1	B	8	HIS
1	B	24	GLN
1	B	111	GLN
1	B	138	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	181	HIS
1	B	217	GLN
1	B	241	ASN
1	B	251	HIS
1	B	252	HIS
1	B	257	GLN
1	B	353	GLN
1	B	358	ASN
1	B	436	ASN
1	B	448	GLN
1	B	498	GLN
1	B	511	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	RCD	A	700	-	96,96,96	0.60	0	144,144,144	1.13	12 (8%)
4	GOL	A	800	-	5,5,5	0.59	0	5,5,5	0.64	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	A	801	-	5,5,5	0.43	0	5,5,5	0.41	0
3	RCD	B	700	-	96,96,96	0.53	0	144,144,144	0.92	6 (4%)
4	GOL	B	800	-	5,5,5	0.52	0	5,5,5	0.29	0
4	GOL	B	801	-	5,5,5	0.31	0	5,5,5	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RCD	A	700	-	-	12/48/208/208	-
4	GOL	A	800	-	-	0/4/4/4	-
4	GOL	A	801	-	-	2/4/4/4	-
3	RCD	B	700	-	-	7/48/208/208	-
4	GOL	B	800	-	-	0/4/4/4	-
4	GOL	B	801	-	-	1/4/4/4	-

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	700	RCD	O5D-C5D-C4D	3.26	116.66	109.76
3	B	700	RCD	O4F-C4F-C3F	2.96	115.25	107.28
3	A	700	RCD	C1A-O5A-C5M	2.78	119.17	113.70
3	A	700	RCD	O4D-C1C-C2C	2.78	115.41	108.10
3	A	700	RCD	C1D-O5D-C5D	2.66	118.93	113.70
3	A	700	RCD	C1B-O4C-C4C	-2.63	111.40	117.97
3	A	700	RCD	O4B-C4B-C3B	2.50	114.01	107.28
3	B	700	RCD	C1C-O4D-C4D	-2.45	111.85	117.97
3	B	700	RCD	C1D-O5D-C5D	2.33	118.28	113.70
3	A	700	RCD	C1E-O4F-C4F	-2.28	112.28	117.97
3	A	700	RCD	O5A-C5M-C4A	2.24	114.51	109.76
3	B	700	RCD	C1G-O4H-C4H	-2.23	112.42	117.97
3	B	700	RCD	C1E-O4F-C4F	2.17	123.39	117.97
3	A	700	RCD	O5C-C5C-C6C	2.16	111.81	106.43
3	A	700	RCD	C2E-C3E-C4E	2.12	114.51	109.67
3	B	700	RCD	O4G-C4G-C5G	-2.09	103.68	109.43
3	A	700	RCD	O3E-C3E-C2E	-2.08	105.51	110.34
3	A	700	RCD	O4D-C4D-C3D	2.02	112.71	107.28

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	801	GOL	O1-C1-C2-C3
3	B	700	RCD	O5D-C1D-O4E-C4E
3	A	700	RCD	O5A-C1A-O4B-C4B
3	A	700	RCD	O5D-C5D-C6D-O6D
3	A	700	RCD	O5F-C5F-C6F-O6F
3	A	700	RCD	C4D-C5D-C6D-O6D
3	B	700	RCD	O5F-C5F-C6F-O6F
3	A	700	RCD	C5D-C4D-O4D-C1C
3	B	700	RCD	C3F-C4F-O4F-C1E
4	A	801	GOL	O1-C1-C2-C3
3	A	700	RCD	C5E-C4E-O4E-C1D
3	A	700	RCD	C3E-C4E-O4E-C1D
3	A	700	RCD	C4F-C5F-C6F-O6F
3	B	700	RCD	C4F-C5F-C6F-O6F
3	B	700	RCD	C5F-C4F-O4F-C1E
3	A	700	RCD	C3D-C4D-O4D-C1C
4	A	801	GOL	O1-C1-C2-O2
3	B	700	RCD	C3G-C4G-O4G-C1F
3	B	700	RCD	C5G-C4G-O4G-C1F
3	A	700	RCD	C4B-C5B-C6B-O6B
3	A	700	RCD	O5F-C1F-O4G-C4G
3	A	700	RCD	C2F-C1F-O4G-C4G

There are no ring outliers.


2 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	700	RCD	2	0
3	B	700	RCD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.



The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	597/601 (99%)	0.12	28 (4%) 31 30	10, 17, 35, 54	0
1	B	597/601 (99%)	0.22	34 (5%) 24 23	9, 19, 39, 58	0
All	All	1194/1202 (99%)	0.17	62 (5%) 27 26	9, 18, 37, 58	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	THR	17.8
1	B	3	PRO	16.3
1	A	3	PRO	15.6
1	B	4	THR	8.6
1	B	50	THR	7.7
1	A	50	THR	6.9
1	B	53	PRO	6.4
1	A	79	GLY	6.3
1	B	81	GLY	6.3
1	A	51	ARG	6.1
1	B	51	ARG	5.7
1	A	81	GLY	5.7
1	A	49	PRO	5.0
1	B	49	PRO	4.5
1	B	581	ARG	4.5
1	B	599	PRO	4.5
1	A	52	VAL	4.3
1	A	53	PRO	4.2
1	A	48	SER	4.2
1	A	80	ASP	4.1
1	A	30	ARG	4.1
1	B	386	GLY	4.1
1	B	52	VAL	3.8
1	B	80	ASP	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	385	THR	3.8
1	B	48	SER	3.6
1	B	95	GLN	3.5
1	B	5	ALA	3.5
1	A	95	GLN	3.4
1	B	137[A]	ARG	3.3
1	A	463	GLY	3.3
1	B	30[A]	ARG	3.3
1	A	34	ARG	3.1
1	A	581	ARG	3.1
1	B	576[A]	THR	3.0
1	B	387	GLU	2.9
1	B	55	ALA	2.9
1	A	137	ARG	2.8
1	A	462	LYS	2.7
1	B	598	LEU	2.7
1	B	388[A]	GLU	2.7
1	A	85	ARG	2.6
1	A	5	ALA	2.6
1	B	40	ASP	2.5
1	B	510	ASN	2.5
1	B	82	ARG	2.5
1	B	85	ARG	2.5
1	B	64	ILE	2.4
1	A	576[A]	THR	2.4
1	B	548[A]	ASP	2.3
1	A	128	ALA	2.3
1	A	47	VAL	2.2
1	A	83	SER	2.2
1	A	94	GLU	2.2
1	B	148	ARG	2.1
1	A	245[A]	LYS	2.1
1	B	536[A]	LYS	2.1
1	B	232	LYS	2.1
1	B	78	LYS	2.1
1	A	78	LYS	2.0
1	A	82	ARG	2.0
1	B	36	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	RCD	A	700	88/88	0.74	0.24	8,23,41,42	88
3	RCD	B	700	88/88	0.75	0.20	10,22,29,33	88
4	GOL	A	801	6/6	0.89	0.13	37,38,39,40	0
4	GOL	B	800	6/6	0.96	0.08	20,21,22,23	0
4	GOL	A	800	6/6	0.96	0.09	13,15,16,16	0
4	GOL	B	801	6/6	0.97	0.11	29,32,34,35	0
2	CA	B	603	1/1	0.99	0.07	20,20,20,20	0
2	CA	A	602	1/1	0.99	0.04	15,15,15,15	0
2	CA	B	602	1/1	0.99	0.06	16,16,16,16	0
2	CA	A	603	1/1	0.99	0.06	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around RCD A 700:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around RCD B 700:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.