



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 06:52 am GMT

PDB ID : 2F53
Title : Directed Evolution of Human T-cell Receptor CDR2 residues by phage display dramatically enhances affinity for cognate peptide-MHC without apparent cross-reactivity
Authors : Rizkallah, P.J.; Jakobsen, B.K.; Dunn, S.M.; Sami, M.
Deposited on : 2005-11-25
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4608 (2.10-2.10)
Clashscore	122126	5109 (2.10-2.10)
Ramachandran outliers	120053	5059 (2.10-2.10)
Sidechain outliers	120020	5060 (2.10-2.10)
RSRZ outliers	108989	4497 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	<div> <div>4%</div> <div> <div></div> <div>88%</div> <div>12%</div> </div> </div>
2	B	100	<div> <div>%</div> <div> <div></div> <div>91%</div> <div>8%</div> </div> </div>
3	C	9	<div> <div>11%</div> <div> <div></div> <div>100%</div> </div> </div>
4	D	193	<div> <div>9%</div> <div> <div></div> <div>77%</div> <div>21%</div> </div> </div>
5	E	243	<div> <div>2%</div> <div> <div></div> <div>81%</div> <div>16%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	GOL	E	805	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 7335 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	2	0
			2259	1413	409	428	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	1	0
			835	530	140	160	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	CLONING ARTIFACT	UNP P61769
B	91	CYS	LYS	ENGINEERED	UNP P61769

- Molecule 3 is a protein called Cancer/testis antigen 1B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	S	0	0	0
			75	49	11	13	2			

- Molecule 4 is a protein called T-cell Receptor, alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	193	Total	C	N	O	S	0	3	0
			1493	931	253	300	9			


- Molecule 5 is a protein called T-cell receptor, beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	243	Total	C	N	O	S	0	1	0
			1920	1205	330	374	11			

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Na	0	0
			1	1		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		
7	E	1	Total	C	O	0	0
			6	3	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	252	Total	O	0	0
			252	252		
8	B	112	Total	O	0	0
			112	112		
8	C	9	Total	O	0	0
			9	9		
8	D	134	Total	O	0	0
			134	134		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	E	221	Total 221	O 221	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	76.23Å 53.97Å 119.91Å 90.00° 96.83° 90.00°	Depositor
Resolution (Å)	37.85 – 2.10 37.85 – 1.99	Depositor EDS
% Data completeness (in resolution range)	98.7 (37.85-2.10) 97.5 (37.85-1.99)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 2.00Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.166 , 0.231 0.176 , 0.237	Depositor DCC
R_{free} test set	3285 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.515	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7335	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/2333	0.71	0/3167
2	B	0.71	0/863	0.81	0/1167
3	C	0.76	0/76	0.80	0/101
4	D	0.63	1/1541 (0.1%)	0.76	1/2093 (0.0%)
5	E	0.69	0/1976	0.72	0/2690
All	All	0.67	1/6789 (0.0%)	0.74	1/9218 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	36	ARG	CB-CG	-6.75	1.34	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	D	36	ARG	NE-CZ-NH2	-6.75	116.93	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	84	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2259	0	2107	28	0
2	B	835	0	796	9	0
3	C	75	0	79	0	0
4	D	1493	0	1438	47	0
5	E	1920	0	1814	36	0
6	A	1	0	0	0	0
7	A	6	0	8	0	0
7	B	12	0	16	0	0
7	E	6	0	8	4	0
8	A	252	0	0	7	0
8	B	112	0	0	1	0
8	C	9	0	0	0	0
8	D	134	0	0	1	0
8	E	221	0	0	5	0
All	All	7335	0	6266	112	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (112) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:VAL:HB	8:B:825:HOH:O	1.31	1.23
4:D:130:SER:OG	4:D:131:SER:HA	1.60	1.02
4:D:1:GLN:HE22	4:D:26:ASP:H	1.14	0.95
4:D:37:GLN:HE22	5:E:35:GLN:HE22	1.15	0.89
2:B:21:ASN:HD22	2:B:22:PHE:H	1.25	0.85
4:D:31:ASN:HD22	4:D:31:ASN:C	1.81	0.82
4:D:130:SER:OG	4:D:131:SER:CA	2.31	0.79
4:D:130:SER:CB	4:D:131:SER:HA	2.13	0.78
1:A:220:ASP:O	8:A:921:HOH:O	2.06	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:ASP:O	1:A:197:HIS:CD2	2.44	0.71
4:D:18:LEU:HD11	4:D:76:ILE:HD12	1.73	0.71
1:A:234:ARG:HE	1:A:242:GLN:HE21	1.38	0.70
1:A:224:GLN:O	1:A:228:THR:HG23	1.94	0.68
4:D:130:SER:HG	4:D:131:SER:HA	1.60	0.66
1:A:196:ASP:O	1:A:197:HIS:CG	2.50	0.65
4:D:17:ASN:ND2	4:D:78:ALA:H	1.94	0.65
5:E:175:LYS:NZ	8:E:967:HOH:O	2.30	0.63
2:B:13:HIS:H	2:B:21:ASN:HD21	1.46	0.63
4:D:136:CYS:HG	4:D:186:CYS:HG	0.65	0.63
2:B:21:ASN:ND2	2:B:22:PHE:H	1.97	0.62
1:A:188:HIS:HD2	8:A:834:HOH:O	1.82	0.62
4:D:1:GLN:HE21	4:D:2:GLU:H	1.48	0.62
4:D:17:ASN:HD22	4:D:78:ALA:H	1.48	0.61
1:A:65:ARG:HG3	5:E:52:MET:HE3	1.83	0.60
4:D:1:GLN:HE22	4:D:26:ASP:N	1.94	0.60
4:D:62:ASN:ND2	8:D:242:HOH:O	2.35	0.60
4:D:130:SER:OG	4:D:131:SER:CB	2.50	0.59
1:A:111:ARG:HD2	1:A:113[B]:TYR:OH	2.03	0.58
4:D:153:SER:O	4:D:154:ASP:HB2	2.03	0.58
1:A:115:GLN:NE2	8:A:1038:HOH:O	2.27	0.58
2:B:21:ASN:HD22	2:B:22:PHE:N	1.99	0.58
4:D:50:PRO:HD2	4:D:53[A]:GLN:CD	2.24	0.57
1:A:194:VAL:O	1:A:195:SER:OG	2.21	0.57
4:D:130:SER:CB	4:D:131:SER:CA	2.82	0.57
1:A:268:LYS:HB2	1:A:268:LYS:NZ	2.19	0.57
5:E:136:GLN:HB3	7:E:805:GOL:H31	1.84	0.57
5:E:39:MET:HE2	5:E:39:MET:HA	1.85	0.57
4:D:149:GLN:O	4:D:151:LYS:N	2.38	0.57
5:E:221:THR:HG22	5:E:221:THR:O	2.03	0.57
5:E:136:GLN:HE21	7:E:805:GOL:C3	2.18	0.56
5:E:62:TYR:C	5:E:63:ASN:HD22	2.09	0.55
5:E:55:GLN:NE2	8:E:897:HOH:O	2.39	0.55
4:D:131:SER:O	4:D:132:ASP:HB2	2.07	0.54
5:E:23:GLN:HE22	5:E:27:HIS:H	1.54	0.54
1:A:253:GLN:NE2	8:A:976:HOH:O	2.39	0.54
4:D:151:LYS:HD2	4:D:151:LYS:O	2.07	0.54
1:A:226:GLN:HA	1:A:226:GLN:HE21	1.73	0.53
5:E:136:GLN:CB	7:E:805:GOL:H31	2.39	0.53
1:A:96:GLN:OE1	2:B:31:HIS:HE1	1.92	0.52
4:D:38:ASP:HB2	4:D:41:LYS:HD2	1.91	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.45	0.52
1:A:93:HIS:HD2	1:A:119:ASP:OD1	1.93	0.52
1:A:197:HIS:HB2	8:A:975:HOH:O	2.09	0.52
1:A:6:ARG:NE	1:A:113[A]:TYR:OH	2.34	0.52
5:E:58:VAL:O	5:E:58:VAL:HG23	2.10	0.51
5:E:129:GLU:HG2	8:E:953:HOH:O	2.12	0.50
5:E:-1:ASN:HD22	5:E:-1:ASN:C	2.14	0.50
8:A:847:HOH:O	2:B:31:HIS:HD2	1.94	0.50
4:D:190:PHE:O	4:D:191:ASN:CB	2.58	0.50
4:D:36:ARG:HB2	4:D:46:LEU:HD11	1.94	0.50
1:A:65:ARG:HG2	4:D:97:GLY:O	2.12	0.49
1:A:226:GLN:HA	1:A:226:GLN:NE2	2.27	0.49
4:D:19:VAL:HG12	4:D:75:TYR:CD2	2.47	0.49
5:E:11:LEU:HD11	5:E:17:MET:HB2	1.95	0.49
1:A:74:HIS:CE1	1:A:97:ARG:HE	2.31	0.48
4:D:37:GLN:HE22	5:E:35:GLN:NE2	1.96	0.48
4:D:31:ASN:ND2	4:D:31:ASN:C	2.55	0.48
4:D:163:LEU:HB3	5:E:168:CYS:HB3	1.95	0.48
4:D:11:SER:OG	4:D:112:HIS:HE1	1.97	0.48
5:E:124:VAL:HG23	5:E:234:ALA:HB3	1.95	0.47
5:E:77:LEU:N	5:E:77:LEU:HD22	2.29	0.47
4:D:190:PHE:O	4:D:191:ASN:HB2	2.14	0.47
1:A:81:LEU:HD13	1:A:118:TYR:CD1	2.50	0.47
5:E:206:ARG:NH2	5:E:208:GLN:OE1	2.47	0.47
5:E:13:THR:HG22	5:E:81:PRO:HD3	1.96	0.46
4:D:31:ASN:HD21	4:D:92:ARG:HG2	1.79	0.46
5:E:228:VAL:O	5:E:230:GLN:HG2	2.15	0.46
4:D:163:LEU:HD12	4:D:163:LEU:C	2.37	0.45
4:D:5:GLN:HE21	4:D:104:GLY:HA3	1.81	0.45
4:D:1:GLN:HA	4:D:1:GLN:NE2	2.31	0.45
5:E:8:PHE:CE1	5:E:107:ARG:HG3	2.51	0.45
5:E:136:GLN:NE2	7:E:805:GOL:H32	2.32	0.45
4:D:48:LEU:C	4:D:48:LEU:HD13	2.37	0.45
1:A:131:ARG:HD2	8:A:1048:HOH:O	2.17	0.44
5:E:21:CYS:HB2	5:E:32:TRP:CZ2	2.53	0.44
4:D:156:TYR:O	4:D:177:ALA:HA	2.17	0.44
5:E:17:MET:HE3	5:E:19:LEU:HD23	2.00	0.43
5:E:221:THR:CG2	5:E:221:THR:O	2.66	0.43
4:D:1:GLN:HA	4:D:1:GLN:HE21	1.83	0.43
1:A:197:HIS:ND1	1:A:198:GLU:HG3	2.34	0.43
4:D:100:ILE:HD11	5:E:46:TYR:CE2	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:18:LEU:HD12	4:D:18:LEU:C	2.39	0.43
5:E:13:THR:HG21	8:E:1010:HOH:O	2.18	0.43
5:E:2:VAL:CG1	5:E:89[B]:CYS:SG	3.07	0.42
4:D:163:LEU:HD12	4:D:163:LEU:O	2.19	0.42
2:B:31:HIS:CD2	2:B:32:PRO:HA	2.54	0.42
5:E:174:LEU:HD12	5:E:174:LEU:C	2.40	0.42
5:E:144:ALA:O	5:E:186:ALA:HA	2.19	0.42
1:A:231:VAL:CG1	1:A:244:TRP:CZ2	3.03	0.42
2:B:22:PHE:CE2	2:B:69:GLU:HG2	2.54	0.42
4:D:131:SER:O	4:D:132:ASP:CB	2.67	0.42
5:E:175:LYS:HE2	8:E:989:HOH:O	2.19	0.42
4:D:11:SER:OG	4:D:112:HIS:CE1	2.73	0.41
5:E:149:PRO:HG2	5:E:151:HIS:CD2	2.56	0.41
4:D:20:LEU:HD22	4:D:107:THR:HG21	2.03	0.41
1:A:74:HIS:HE1	1:A:97:ARG:HE	1.69	0.41
4:D:1:GLN:HE21	4:D:2:GLU:N	2.17	0.41
5:E:44:ILE:HG22	5:E:45:HIS:CD2	2.55	0.41
4:D:62:ASN:ND2	4:D:75:TYR:HB2	2.36	0.41
5:E:154:LEU:HD23	5:E:155:SER:N	2.35	0.41
1:A:254:GLU:H	1:A:254:GLU:CD	2.24	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/275 (100%)	266 (97%)	9 (3%)	0	100	100
2	B	99/100 (99%)	97 (98%)	2 (2%)	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
4	D	194/193 (100%)	179 (92%)	11 (6%)	4 (2%)	8	3

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	E	242/243 (100%)	233 (96%)	9 (4%)	0	100	100
All	All	817/820 (100%)	782 (96%)	31 (4%)	4 (0%)	31	27

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	130	SER
4	D	150	SER
4	D	152	ASP
4	D	131	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/231 (101%)	232 (100%)	1 (0%)	92	95
2	B	96/95 (101%)	93 (97%)	3 (3%)	43	45
3	C	9/9 (100%)	9 (100%)	0	100	100
4	D	172/169 (102%)	169 (98%)	3 (2%)	63	70
5	E	212/211 (100%)	205 (97%)	7 (3%)	41	43
All	All	722/715 (101%)	708 (98%)	14 (2%)	58	66

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	268	LYS
2	B	21	ASN
2	B	40	LEU
2	B	83	ASN
4	D	0	LYS
4	D	31	ASN
4	D	62	ASN
5	E	-1	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
5	E	23	GLN
5	E	52	MET
5	E	96	ASN
5	E	107	ARG
5	E	168	CYS
5	E	190	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (46) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	GLN
1	A	70	HIS
1	A	74	HIS
1	A	86	ASN
1	A	93	HIS
1	A	141	GLN
1	A	174	ASN
1	A	188	HIS
1	A	191	HIS
1	A	226	GLN
1	A	242	GLN
1	A	253	GLN
1	A	255	GLN
2	B	2	GLN
2	B	21	ASN
2	B	24	ASN
2	B	31	HIS
2	B	83	ASN
2	B	89	GLN
4	D	1	GLN
4	D	5	GLN
4	D	17	ASN
4	D	21	ASN
4	D	31	ASN
4	D	62	ASN
4	D	80	GLN
4	D	112	HIS
4	D	116	GLN
4	D	124	GLN
4	D	144	GLN
4	D	146	ASN
4	D	188	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	191	ASN
5	E	-1	ASN
5	E	9	GLN
5	E	15	GLN
5	E	23	GLN
5	E	26	ASN
5	E	35	GLN
5	E	55	GLN
5	E	60	ASN
5	E	63	ASN
5	E	96	ASN
5	E	116	ASN
5	E	136	GLN
5	E	151	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GOL	A	803	-	5,5,5	0.42	0	5,5,5	0.14	0
7	GOL	B	802	-	5,5,5	0.41	0	5,5,5	0.45	0
7	GOL	B	804	-	5,5,5	0.42	0	5,5,5	0.18	0
7	GOL	E	805	-	5,5,5	0.78	0	5,5,5	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	803	-	-	0/4/4/4	0/0/0/0
7	GOL	B	802	-	-	0/4/4/4	0/0/0/0
7	GOL	B	804	-	-	0/4/4/4	0/0/0/0
7	GOL	E	805	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	E	805	GOL	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	-0.03	11 (4%) 38 45	3, 12, 35, 63	0
2	B	100/100 (100%)	-0.29	1 (1%) 82 85	3, 11, 32, 66	0
3	C	9/9 (100%)	1.29	1 (11%) 5 7	2, 3, 6, 10	0
4	D	193/193 (100%)	0.35	17 (8%) 10 13	8, 20, 45, 59	0
5	E	243/243 (100%)	0.03	4 (1%) 72 76	2, 10, 26, 57	0
All	All	820/820 (100%)	0.06	34 (4%) 37 44	2, 13, 38, 66	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	ILE	6.1
1	A	194	VAL	5.0
4	D	183	ASP	5.0
4	D	150	SER	4.9
4	D	188	ASN	4.6
1	A	197	HIS	4.5
4	D	187	ALA	4.3
1	A	196	ASP	4.3
1	A	195	SER	4.2
5	E	-1	ASN	4.0
4	D	182	SER	4.0
4	D	-1	MET	3.5
4	D	154	ASP	3.3
1	A	17	ARG	3.1
5	E	0	ALA	3.1
4	D	191	ASN	3.0
4	D	146	ASN	2.9
4	D	151	LYS	2.9
1	A	267	PRO	2.8
4	D	148	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	193	ALA	2.7
4	D	132	ASP	2.7
4	D	114	TYR	2.7
4	D	153	SER	2.6
1	A	18	GLY	2.6
4	D	149	GLN	2.6
1	A	192	HIS	2.6
5	E	223	ASP	2.5
1	A	16	GLY	2.4
4	D	144	GLN	2.4
4	D	152	ASP	2.2
3	C	3	LEU	2.1
1	A	225	THR	2.1
5	E	216	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	GOL	E	805	6/6	0.83	0.31	15,36,40,45	0
7	GOL	B	804	6/6	0.88	0.33	28,47,56,58	0
7	GOL	A	803	6/6	0.91	0.13	30,38,42,42	0
6	NA	A	801	1/1	0.95	0.20	27,27,27,27	0
7	GOL	B	802	6/6	0.95	0.32	9,10,25,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.