



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2018 – 04:57 am GMT

PDB ID : 2F55
Title : Two hepatitis c virus ns3 helicase domains complexed with the same strand of dna
Authors : Lu, J.Z.; Jordan, J.B.; Sakon, J.
Deposited on : 2005-11-25
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30686


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122078	1021 (3.34-3.26)
Ramachandran outliers	120005	1003 (3.34-3.26)
Sidechain outliers	119972	1002 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	D	13	<div> <div>23%</div> <div>69%</div> <div>8%</div> </div>
2	E	3	<div> <div>100%</div> </div>
3	A	435	<div> <div>38%</div> <div>52%</div> <div>9%</div> <div>..</div> </div>
3	B	435	<div> <div>28%</div> <div>57%</div> <div>14%</div> <div>.</div> </div>
3	C	435	<div> <div>37%</div> <div>51%</div> <div>10%</div> <div>..</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10129 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(P*(DU)P*(DU)P*(DU)P*(DU)P*(DU)P*(DU)P*(DU)P*(DU)P*(DU)P*(DU)P*(DU))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	13	Total	C	N	O	P	0	0	0
			247	117	26	91	13			

- Molecule 2 is a DNA chain called 5'-D(P*(DU)P*(DU)P*(DU))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	3	Total	C	N	O	P	0	0	0
			57	27	6	21	3			

- Molecule 3 is a protein called polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	432	Total	C	N	O	S	0	0	0
			3251	2063	553	616	19			
3	B	432	Total	C	N	O	S	0	0	0
			3251	2063	553	616	19			
3	C	432	Total	C	N	O	S	0	0	0
			3251	2063	553	616	19			

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	3	Total	O	0	0
			3	3		
5	A	20	Total	O	0	0
			20	20		
5	B	14	Total	O	0	0
			14	14		
5	C	25	Total	O	0	0
			25	25		

C622
R623
S624


• Molecule 3: polyprotein

Chain B: 28% 57% 14%



• Molecule 3: polyprotein

Chain C: 37% 51% 10%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	108.20Å 109.80Å 183.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.30	Depositor
% Data completeness (in resolution range)	83.2 (50.00-3.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.247 , 0.273	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	10129	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	D	0.70	0/272	0.98	1/412 (0.2%)
2	E	0.70	0/62	0.99	0/92
3	A	0.54	3/3331 (0.1%)	0.73	2/4548 (0.0%)
3	B	0.59	4/3331 (0.1%)	0.74	1/4548 (0.0%)
3	C	0.53	5/3331 (0.2%)	0.72	1/4548 (0.0%)
All	All	0.56	12/10327 (0.1%)	0.74	5/14148 (0.0%)

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	469	ARG	NE-CZ	7.51	1.42	1.33
3	B	469	ARG	NE-CZ	7.47	1.42	1.33
3	A	469	ARG	NE-CZ	7.45	1.42	1.33
3	B	469	ARG	CZ-NH2	7.42	1.42	1.33
3	C	469	ARG	CZ-NH2	7.38	1.42	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	402	PRO	CA-N-CD	-12.76	93.64	111.50
3	B	402	PRO	CA-N-CD	-12.74	93.67	111.50
3	A	402	PRO	CA-N-CD	-11.23	95.78	111.50
3	A	401	PRO	C-N-CD	7.38	143.90	128.40
1	D	17	DU	O4'-C1'-C2'	5.03	109.92	105.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	247	0	131	18	0
2	E	57	0	31	6	0
3	A	3251	0	3216	332	0
3	B	3251	0	3215	434	0
3	C	3251	0	3215	354	0
4	A	5	0	0	0	0
4	C	5	0	0	1	0
5	A	20	0	0	2	0
5	B	14	0	0	0	0
5	C	25	0	0	3	0
5	D	3	0	0	0	0
All	All	10129	0	9808	1118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 56.

The worst 5 of 1118 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:469:ARG:HB2	3:A:469:ARG:NH1	1.31	1.43
3:C:513:LEU:O	3:C:517:LEU:CD2	1.65	1.42
3:A:360:LYS:NZ	3:A:386:LEU:HD21	1.06	1.36
3:A:360:LYS:NZ	3:A:386:LEU:CD2	1.88	1.35
3:A:339:ILE:CG2	3:A:474:ARG:HG2	1.54	1.35

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	428/435 (98%)	318 (74%)	86 (20%)	24 (6%)	2	13
3	B	428/435 (98%)	282 (66%)	104 (24%)	42 (10%)	1	4
3	C	428/435 (98%)	296 (69%)	92 (22%)	40 (9%)	1	4
All	All	1284/1305 (98%)	896 (70%)	282 (22%)	106 (8%)	1	6

5 of 106 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	246	HIS
3	A	349	PHE
3	A	413	ALA
3	A	583	LYS
3	B	245	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	352/354 (99%)	323 (92%)	29 (8%)	12	40
3	B	352/354 (99%)	311 (88%)	41 (12%)	6	25
3	C	352/354 (99%)	320 (91%)	32 (9%)	10	35
All	All	1056/1062 (99%)	954 (90%)	102 (10%)	9	32

5 of 102 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	B	393	ARG
3	B	524	VAL
3	C	526	GLN
3	B	402	PRO
3	B	460	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
3	B	564	GLN
3	C	198	GLN
3	C	564	GLN
3	B	580	GLN
3	C	201	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	101	-	4,4,4	0.33	0	6,6,6	0.20	0
4	SO4	C	102	-	4,4,4	0.28	0	6,6,6	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SO4	A	101	-	-	0/0/0/0	0/0/0/0
4	SO4	C	102	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	102	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.