



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 08:27 pm GMT

PDB ID : 3F5U
Title : Crystal structure of the death associated protein kinase in complex with AMPPNP and Mg²⁺
Authors : McNamara, L.K.; Watterson, D.M.; Brunzelle, J.S.
Deposited on : 2008-11-04
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	trunk30967
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	7193 (2.00-2.00)
Clashscore	122126	8267 (2.00-2.00)
Ramachandran outliers	120053	8166 (2.00-2.00)
Sidechain outliers	120020	8165 (2.00-2.00)
RSRZ outliers	108989	7011 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	<div> <div>9%</div> <div>76%</div> <div>15%</div> <div>• 5%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2589 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Death-associated protein kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	280	2453	1573	409	467	4	12	21	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	286	SER	-	EXPRESSION TAG	UNP P53355
A	287	ALA	-	EXPRESSION TAG	UNP P53355
A	288	TRP	-	EXPRESSION TAG	UNP P53355
A	289	SER	-	EXPRESSION TAG	UNP P53355
A	290	HIS	-	EXPRESSION TAG	UNP P53355
A	291	PRO	-	EXPRESSION TAG	UNP P53355
A	292	GLN	-	EXPRESSION TAG	UNP P53355
A	293	PHE	-	EXPRESSION TAG	UNP P53355
A	294	GLU	-	EXPRESSION TAG	UNP P53355
A	295	LYS	-	EXPRESSION TAG	UNP P53355

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	99	Total	O	0	0
			99	99		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Death-associated protein kinase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.67Å 62.46Å 88.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 28.57 – 1.98	Depositor EDS
% Data completeness (in resolution range)	99.0 (30.00-2.00) 99.1 (28.57-1.98)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.14 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.177 , 0.245 0.181 , 0.243	Depositor DCC
R_{free} test set	941 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	27.5	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 53.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2589	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	1/2501 (0.0%)	0.88	6/3378 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	108	LYS	C-N	5.97	1.47	1.34

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	176	ASN	N-CA-C	8.48	133.90	111.00
1	A	109[A]	GLU	N-CA-C	6.13	127.54	111.00
1	A	109[B]	GLU	N-CA-C	6.13	127.54	111.00
1	A	24[A]	PHE	CB-CA-C	-5.57	99.27	110.40
1	A	24[B]	PHE	CB-CA-C	-5.57	99.27	110.40
1	A	108	LYS	C-N-CA	5.07	134.38	121.70

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	109[A]	GLU	CA

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	107	GLU	Mainchain,Peptide
1	A	108	LYS	Mainchain,Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2453	0	2433	60	1
2	A	31	0	13	1	0
3	A	1	0	0	0	0
4	A	5	0	0	1	0
5	A	99	0	0	4	1
All	All	2589	0	2446	60	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:71[A]:ILE:HG23	1:A:76[A]:VAL:HG21	1.40	1.03
1:A:77:ILE:HD11	1:A:96:VAL:CG2	1.90	1.01
1:A:71[A]:ILE:CG2	1:A:76[A]:VAL:HG21	1.91	1.01
1:A:108:LYS:CG	1:A:109[B]:GLU:H	1.60	0.98
1:A:71[A]:ILE:CG2	1:A:76[A]:VAL:CG2	2.43	0.97
1:A:50:LYS:H	1:A:50:LYS:HD2	1.30	0.94
1:A:71[A]:ILE:HG21	1:A:76[A]:VAL:CG2	2.07	0.83
1:A:108:LYS:CG	1:A:109[B]:GLU:N	2.44	0.78
1:A:71[A]:ILE:HG21	1:A:76[A]:VAL:HG23	1.66	0.77
1:A:108:LYS:HG3	1:A:109[B]:GLU:H	1.50	0.76
1:A:71[A]:ILE:CG2	1:A:76[A]:VAL:HG23	2.16	0.74
1:A:77:ILE:HD11	1:A:96:VAL:HG23	1.70	0.73
1:A:23[A]:GLN:O	1:A:24[A]:PHE:CB	2.33	0.73
1:A:23[A]:GLN:O	1:A:24[A]:PHE:HB2	1.90	0.71
1:A:152[A]:VAL:HG12	5:A:379:HOH:O	1.89	0.71

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:LYS:H	1:A:50:LYS:CD	2.04	0.70
1:A:77:ILE:CD1	1:A:96:VAL:CG2	2.69	0.68
1:A:71[A]:ILE:HG23	1:A:76[A]:VAL:CG2	2.18	0.64
1:A:108:LYS:HG2	1:A:109[B]:GLU:H	1.54	0.63
1:A:48:ARG:HG2	1:A:48:ARG:HH21	1.64	0.62
1:A:131:HIS:HE1	1:A:199:ASP:OD2	1.84	0.61
1:A:50:LYS:HD2	1:A:50:LYS:N	2.11	0.60
1:A:71[B]:ILE:HG13	1:A:129:TYR:HE2	1.67	0.60
1:A:77:ILE:CD1	1:A:96:VAL:HG21	2.32	0.59
1:A:23[A]:GLN:O	1:A:24[A]:PHE:CG	2.55	0.59
1:A:182[B]:GLU:OE2	1:A:215:SER:OG	2.20	0.58
1:A:141:LYS:HE3	1:A:143:GLU:HG2	1.86	0.58
1:A:152[A]:VAL:CG1	1:A:153:PRO:HD2	2.33	0.58
1:A:96:VAL:HG11	1:A:146:MET:HB2	1.84	0.58
1:A:175:LYS:HD2	4:A:298:SO4:O3	2.04	0.57
1:A:152[A]:VAL:HG13	1:A:153:PRO:HD2	1.88	0.56
1:A:70:GLU:OE2	1:A:167:LYS:HE2	2.05	0.56
1:A:120[C]:LEU:HD11	1:A:207:THR:HG23	1.89	0.54
1:A:48:ARG:HG2	1:A:48:ARG:NH2	2.22	0.54
1:A:164:LEU:O	1:A:166:HIS:HD2	1.91	0.53
1:A:173:GLU:OE1	1:A:175:LYS:HB2	2.10	0.52
1:A:222:LYS:HB2	1:A:294:GLU:HG2	1.91	0.51
1:A:77:ILE:HD13	1:A:160:ILE:HG22	1.93	0.51
1:A:176:ASN:C	5:A:386:HOH:O	2.48	0.51
1:A:223[A]:GLN:HG2	1:A:224:GLU:N	2.25	0.50
1:A:265:THR:OG1	1:A:267[B]:GLN:HG2	2.11	0.50
1:A:23[A]:GLN:HG3	2:A:296:ANP:O1G	2.11	0.50
1:A:48:ARG:NH1	1:A:87[A]:THR:HG22	2.26	0.49
1:A:176:ASN:HB3	5:A:386:HOH:O	2.12	0.49
1:A:254:ARG:HB3	1:A:264:MET:HG3	1.95	0.48
1:A:45:LYS:HB2	1:A:45:LYS:HE2	1.54	0.48
1:A:109[A]:GLU:O	1:A:109[A]:GLU:HG2	2.14	0.47
1:A:223[A]:GLN:HE21	1:A:223[A]:GLN:H	1.62	0.47
1:A:23[A]:GLN:C	1:A:24[A]:PHE:CG	2.89	0.46
1:A:141:LYS:HE3	1:A:143:GLU:CG	2.45	0.46
1:A:223[A]:GLN:NE2	1:A:223[A]:GLN:H	2.15	0.44
1:A:166:HIS:CE1	1:A:175:LYS:HD3	2.53	0.44
1:A:101:LEU:O	1:A:105:LEU:HG	2.18	0.44
1:A:146:MET:HG3	1:A:160:ILE:HG21	2.02	0.42
1:A:157:ILE:HD12	1:A:157:ILE:C	2.40	0.41
1:A:29:LYS:HD3	1:A:38:GLN:NE2	2.36	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:176:ASN:CB	5:A:386:HOH:O	2.67	0.41
1:A:120[B]:LEU:O	1:A:124:LEU:HG	2.21	0.41
1:A:271:GLN:HE22	1:A:276:LYS:NZ	2.19	0.40
1:A:73:HIS:HB3	1:A:76[A]:VAL:HG22	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223[B]:GLN:OE1	5:A:354:HOH:O[3_545]	1.42	0.78

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	297/295 (101%)	288 (97%)	8 (3%)	1 (0%)	43 39

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	110	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	271/263 (103%)	256 (94%)	15 (6%)	24 19

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23[A]	GLN
1	A	23[B]	GLN
1	A	45	LYS
1	A	48	ARG
1	A	50	LYS
1	A	53	ARG
1	A	54	ARG
1	A	77	ILE
1	A	107	GLU
1	A	118	GLU
1	A	131	HIS
1	A	150	ARG
1	A	156	ARG
1	A	223[A]	GLN
1	A	223[B]	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	GLN
1	A	131	HIS
1	A	166	HIS
1	A	228	ASN
1	A	271	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ANP	A	296	3	29,33,33	2.16	7 (24%)	29,52,52	2.09	6 (20%)
4	SO4	A	298	-	4,4,4	0.17	0	6,6,6	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	A	296	3	-	0/13/38/38	0/3/3/3
4	SO4	A	298	-	-	0/0/0/0	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	296	ANP	O4'-C1'	2.20	1.44	1.41
2	A	296	ANP	PB-O1B	2.98	1.49	1.46
2	A	296	ANP	C5-C4	3.27	1.47	1.40
2	A	296	ANP	PB-N3B	3.31	1.72	1.63
2	A	296	ANP	PG-N3B	3.87	1.73	1.63
2	A	296	ANP	PB-O3A	5.43	1.65	1.59
2	A	296	ANP	PG-O1G	6.10	1.53	1.46

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	296	ANP	N3-C2-N1	-7.26	122.65	128.86
2	A	296	ANP	O1G-PG-N3B	-4.98	104.35	111.79
2	A	296	ANP	O3A-PB-N3B	2.38	113.20	106.59
2	A	296	ANP	C2-N1-C6	2.49	122.98	118.75
2	A	296	ANP	C4'-O4'-C1'	2.50	112.43	109.83
2	A	296	ANP	O3G-PG-O2G	2.88	115.44	107.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	296	ANP	1	0
4	A	298	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	280/295 (94%)	0.21	27 (9%) 8 7	17, 26, 52, 100	3 (1%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	293	PHE	10.2
1	A	292	GLN	5.7
1	A	151	ASN	5.3
1	A	291	PRO	5.0
1	A	176	ASN	5.0
1	A	152[A]	VAL	4.4
1	A	153	PRO	4.2
1	A	53	ARG	4.1
1	A	155	PRO	3.7
1	A	109[A]	GLU	3.7
1	A	277	PRO	3.7
1	A	24[A]	PHE	3.7
1	A	106	ALA	3.1
1	A	154	LYS	2.9
1	A	48	ARG	2.9
1	A	175	LYS	2.9
1	A	294	GLU	2.9
1	A	108	LYS	2.6
1	A	150	ARG	2.6
1	A	52	SER	2.5
1	A	54	ARG	2.4
1	A	169	ASP	2.4
1	A	110	SER	2.3
1	A	102	PHE	2.2
1	A	174	PHE	2.2
1	A	50	LYS	2.1
1	A	51	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MG	A	297	1/1	0.67	0.09	51,51,51,51	0
4	SO4	A	298	5/5	0.94	0.21	84,84,84,84	0
2	ANP	A	296	31/31	0.96	0.11	19,28,43,46	3

6.5 Other polymers [i](#)

There are no such residues in this entry.