



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 06:38 am GMT

PDB ID : 1F8F
Title : CRYSTAL STRUCTURE OF BENZYL ALCOHOL DEHYDROGENASE
FROM ACINETOBACTER CALCOACETICUS
Authors : Beauchamp, J.C.; Gillooly, D.; Warwicker, J.; Fewson, C.A.; Laphorn, A.J.
Deposited on : 2000-06-30
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	5027 (2.20-2.20)
Ramachandran outliers	120053	4952 (2.20-2.20)
Sidechain outliers	120020	4953 (2.20-2.20)
RSRZ outliers	108989	4245 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	371	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EOH	A	374	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3040 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BENZYL ALCOHOL DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	362	Total	C	N	O	S	0	1	0
			2672	1702	449	509	12			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	54	LYS	TYR	CONFLICT	UNP Q59096
A	194	PHE	LEU	CONFLICT	UNP Q59096
A	268	SER	ARG	CONFLICT	UNP Q59096
A	359	ARG	HIS	CONFLICT	UNP Q59096

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 5	O 4	S 1	0	0
3	A	1	Total 5	O 4	S 1	0	0

- Molecule 4 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $\text{C}_{21}\text{H}_{27}\text{N}_7\text{O}_{14}\text{P}_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 44	C 21	N 7	O 14	P 2	0	0

- Molecule 5 is ETHANOL (three-letter code: EOH) (formula: C_2H_6O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			3	2	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	309	Total	O	0	0
			309	309		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BENZYL ALCOHOL DEHYDROGENASE



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	82.29Å 96.98Å 103.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.20 19.76 – 2.20	Depositor EDS
% Data completeness (in resolution range)	91.0 (20.00-2.20) 91.6 (19.76-2.20)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 2.19Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.197 , 0.263 0.192 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	27.2	Xtriage
Anisotropy	0.276	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 59.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3040	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, EOH, SO4, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/2722	1.81	51/3693 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (51) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	330	ARG	CD-NE-CZ	35.12	172.76	123.60
1	A	114	ARG	NE-CZ-NH2	14.81	127.71	120.30
1	A	227	ARG	NE-CZ-NH1	10.77	125.68	120.30
1	A	330	ARG	NE-CZ-NH2	10.56	125.58	120.30
1	A	300	ASP	CB-CG-OD1	10.29	127.56	118.30
1	A	299	PHE	CB-CG-CD2	-10.01	113.80	120.80
1	A	140	PHE	CB-CG-CD2	-9.77	113.96	120.80
1	A	27	ARG	CD-NE-CZ	9.20	136.49	123.60
1	A	299	PHE	CB-CG-CD1	8.41	126.69	120.80
1	A	47	ASP	CB-CG-OD1	8.16	125.64	118.30
1	A	114	ARG	NE-CZ-NH1	-8.05	116.28	120.30
1	A	153	GLU	OE1-CD-OE2	-8.05	113.64	123.30
1	A	27	ARG	NE-CZ-NH1	7.61	124.10	120.30
1	A	140	PHE	CB-CG-CD1	7.59	126.11	120.80
1	A	227	ARG	CD-NE-CZ	7.55	134.17	123.60
1	A	110	GLU	OE1-CD-OE2	-7.52	114.27	123.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	211	LYS	N-CA-CB	7.32	123.77	110.60
1	A	253	GLU	OE1-CD-OE2	-7.30	114.54	123.30
1	A	111	PHE	CA-C-N	7.24	133.12	117.20
1	A	222	ASP	CB-CG-OD2	7.16	124.75	118.30
1	A	305	LEU	CA-CB-CG	6.80	130.94	115.30
1	A	186	LEU	O-C-N	-6.79	111.83	122.70
1	A	270	GLU	CB-CG-CD	6.54	131.85	114.20
1	A	54	LYS	C-N-CA	6.46	137.85	121.70
1	A	277	ASP	CB-CG-OD2	6.38	124.04	118.30
1	A	227	ARG	NH1-CZ-NH2	-6.36	112.41	119.40
1	A	340	ASP	CB-CG-OD1	6.07	123.77	118.30
1	A	306	LEU	CA-CB-CG	6.02	129.14	115.30
1	A	162	ASP	CB-CG-OD1	6.01	123.71	118.30
1	A	55	TYR	CB-CG-CD1	5.94	124.56	121.00
1	A	318	SER	CA-C-N	5.91	128.02	116.20
1	A	24	LEU	CA-CB-CG	5.85	128.75	115.30
1	A	186	LEU	C-N-CA	5.74	136.04	121.70
1	A	110	GLU	CG-CD-OE1	5.71	129.73	118.30
1	A	71	ILE	C-N-CA	5.64	135.81	121.70
1	A	66	GLU	OE1-CD-OE2	-5.64	116.54	123.30
1	A	111	PHE	O-C-N	-5.62	113.72	122.70
1	A	305	LEU	CB-CA-C	5.59	120.83	110.20
1	A	349	ASP	CB-CG-OD1	5.54	123.29	118.30
1	A	27	ARG	NE-CZ-NH2	-5.53	117.54	120.30
1	A	318	SER	C-N-CA	-5.50	110.74	122.30
1	A	311	ILE	CA-CB-CG1	5.48	121.41	111.00
1	A	116	PHE	CB-CG-CD2	5.47	124.63	120.80
1	A	327	GLU	OE1-CD-OE2	-5.45	116.77	123.30
1	A	311	ILE	CG1-CB-CG2	-5.41	99.50	111.40
1	A	290	ALA	N-CA-CB	-5.33	102.63	110.10
1	A	20	GLU	CA-CB-CG	5.17	124.78	113.40
1	A	186	LEU	CA-C-O	5.17	130.95	120.10
1	A	80	GLU	CA-CB-CG	5.14	124.72	113.40
1	A	88	VAL	O-C-N	-5.14	114.48	122.70
1	A	120	ASP	CB-CG-OD1	5.07	122.87	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	186	LEU	Mainchain
1	A	367	ILE	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2672	0	2715	34	0
2	A	2	0	0	0	0
3	A	10	0	0	0	0
4	A	44	0	26	1	0
5	A	3	0	6	2	0
6	A	309	0	0	1	0
All	All	3040	0	2747	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:GLY:HA2	1:A:100:CYS:HB3	1.68	0.76
1:A:271:ILE:HD13	1:A:274:GLN:NE2	2.02	0.75
1:A:330:ARG:O	1:A:334:GLN:HG3	1.94	0.68
4:A:375:NAD:H4N	5:A:374:EOH:H11	1.84	0.60
1:A:145:PHE:CE1	1:A:368:ILE:HD13	2.38	0.58
1:A:357:ASP:HB3	1:A:362:ILE:HD11	1.88	0.56
1:A:11:VAL:O	1:A:13:PRO:HD3	2.08	0.53
1:A:26:ILE:HD12	1:A:127:LEU:CD2	2.38	0.53
1:A:84:GLY:O	1:A:161:LYS:NZ	2.43	0.52
1:A:7:ILE:HG21	1:A:148:TYR:HB2	1.92	0.52
1:A:95:GLY:HA2	1:A:100:CYS:CB	2.41	0.51
1:A:299:PHE:HE2	1:A:304:LEU:HD22	1.74	0.51
1:A:71:ILE:HD12	1:A:85:ASP:HB2	1.93	0.50
1:A:61:ALA:HB2	1:A:136:ASN:HB2	1.93	0.49
1:A:87:VAL:HG11	1:A:156:THR:HG22	1.95	0.48
1:A:271:ILE:HD13	1:A:274:GLN:HE22	1.75	0.48
1:A:271:ILE:HD13	1:A:274:GLN:HE21	1.78	0.47
1:A:46:THR:HA	1:A:49:ILE:HG13	1.96	0.46
1:A:14:CYS:O	1:A:15:LYS:C	2.54	0.45
1:A:49:ILE:C	1:A:49:ILE:HD12	2.37	0.45
1:A:28:GLN:HB3	1:A:29:PRO:HD2	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:ARG:HG3	1:A:28:GLN:O	2.17	0.44
1:A:225:GLU:O	1:A:229:GLU:HG2	2.16	0.44
1:A:311:ILE:HG21	1:A:311:ILE:HD13	1.74	0.44
1:A:121:SER:HB2	6:A:580:HOH:O	2.18	0.44
1:A:140:PHE:CZ	5:A:374:EOH:H12	2.53	0.43
1:A:30:GLN:O	1:A:31:GLY:C	2.56	0.43
1:A:299:PHE:CE2	1:A:304:LEU:HD22	2.54	0.42
1:A:186:LEU:HD21	1:A:286:ALA:HB2	2.02	0.41
1:A:229:GLU:O	1:A:233:GLN:HG3	2.19	0.41
1:A:362:ILE:HD12	1:A:363:THR:OG1	2.20	0.41
1:A:49:ILE:HD12	1:A:50:VAL:N	2.35	0.41
1:A:271:ILE:HA	1:A:274:GLN:HE21	1.86	0.41
1:A:244:THR:O	1:A:244:THR:HG22	2.21	0.40
1:A:72:GLU:O	1:A:72:GLU:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	359/371 (97%)	339 (94%)	19 (5%)	1 (0%)	43	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	31	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/294 (98%)	275 (96%)	12 (4%)	32	41

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	LYS
1	A	28	GLN
1	A	34	VAL
1	A	82	GLN
1	A	94	CYS
1	A	128	CYS
1	A	143	SER
1	A	264	GLU
1	A	271	ILE
1	A	304	LEU
1	A	305	LEU
1	A	348	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	ASN
1	A	82	GLN
1	A	99	GLN
1	A	274	GLN
1	A	333	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EOH	A	374	-	2,2,2	0.14	0	1,1,1	0.41	0
4	NAD	A	375	-	40,48,48	1.27	5 (12%)	44,73,73	2.13	10 (22%)
3	SO4	A	376	-	4,4,4	0.75	0	6,6,6	0.29	0
3	SO4	A	377	-	4,4,4	0.80	0	6,6,6	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EOH	A	374	-	-	0/0/0/0	0/0/0/0
4	NAD	A	375	-	-	0/22/62/62	0/5/5/5
3	SO4	A	376	-	-	0/0/0/0	0/0/0/0
3	SO4	A	377	-	-	0/0/0/0	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	375	NAD	O4D-C1D	2.07	1.44	1.41
4	A	375	NAD	C4N-C3N	2.17	1.43	1.39
4	A	375	NAD	C7N-N7N	2.85	1.38	1.33
4	A	375	NAD	C6N-N1N	3.08	1.43	1.35
4	A	375	NAD	C3N-C7N	4.40	1.57	1.50

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	375	NAD	C5N-C4N-C3N	-7.70	111.31	120.35
4	A	375	NAD	C5N-C6N-N1N	-5.08	112.64	120.39
4	A	375	NAD	C4D-O4D-C1D	-4.13	105.53	109.83
4	A	375	NAD	C4B-O4B-C1B	-3.49	106.19	109.83
4	A	375	NAD	C3N-C7N-N7N	-2.34	115.04	117.76
4	A	375	NAD	C3N-C2N-N1N	-2.19	118.22	120.41
4	A	375	NAD	O7N-C7N-N7N	2.16	125.74	122.60
4	A	375	NAD	C2N-C3N-C4N	2.30	120.91	118.26
4	A	375	NAD	C5A-C6A-N6A	2.39	125.34	120.47
4	A	375	NAD	C6N-C5N-C4N	4.11	125.57	119.43

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	374	EOH	2	0
4	A	375	NAD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	362/371 (97%)	-0.58	1 (0%) 93 93	19, 30, 60, 83	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	123	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EOH	A	374	3/3	0.85	0.29	45,45,49,50	0
4	NAD	A	375	44/44	0.92	0.16	38,51,54,55	0
3	SO4	A	377	5/5	0.94	0.25	62,62,63,63	0
3	SO4	A	376	5/5	0.94	0.16	69,70,71,71	0
2	ZN	A	372	1/1	0.97	0.04	42,42,42,42	0
2	ZN	A	373	1/1	1.00	0.01	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.