



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 04:37 pm GMT

PDB ID : 5F9C  
Title : Crystal structure of the G121R mutant of human phosphoglucomutase 1  
Authors : Beamer, L.J.  
Deposited on : 2015-12-09  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4155 (2.50-2.50)
Clashscore	122126	4827 (2.50-2.50)
Ramachandran outliers	120053	4735 (2.50-2.50)
Sidechain outliers	120020	4737 (2.50-2.50)
RSRZ outliers	108989	4058 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	562	<div> <div> <div></div> <div>85%</div> <div>12%</div> <div>••</div> </div> <div>6%</div> </div>
1	B	562	<div> <div> <div></div> <div>81%</div> <div>14%</div> <div>• 5%</div> </div> <div>6%</div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8196 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoglucomutase-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	550	Total	C	N	O	S	0	2	0
			4135	2633	695	791	16			
1	B	536	Total	C	N	O	S	0	0	0
			3922	2494	665	747	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	121	ARG	GLY	engineered mutation	UNP P36871
B	121	ARG	GLY	engineered mutation	UNP P36871

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	57	Total	O	0	0
			57	57		
5	B	23	Total	O	0	0
			23	23		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Phosphoglucomutase-1



#### • Molecule 1: Phosphoglucomutase-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	171.98Å 171.98Å 99.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	60.94 – 2.50 60.80 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (60.94-2.50) 94.2 (60.80-2.20)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.21 (at 2.20Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.192 , 0.243 0.196 , 0.245	Depositor DCC
$R_{free}$ test set	3612 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	45.5	Xtriage
Anisotropy	0.439	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 69.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8196	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.45	0/4217	0.58	0/5719
1	B	0.39	0/3990	0.54	0/5420
All	All	0.42	0/8207	0.57	0/11139

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4135	0	4025	39	0
1	B	3922	0	3720	39	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	15	0	0	2	0
3	B	30	0	0	0	0
4	A	12	0	16	1	0
5	A	57	0	0	6	0
5	B	23	0	0	1	0
All	All	8196	0	7761	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 5.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:329:ARG:NH1	1:B:384:ASP:OD2	2.10	0.84
1:A:329:ARG:NH1	1:A:384:ASP:OD2	2.17	0.78
1:A:18:GLY:HA3	5:A:715:HOH:O	1.92	0.70
1:B:319:SER:HB2	1:B:404:THR:HG21	1.73	0.69
1:A:461:ALA:N	1:A:462:ASN:HA	2.07	0.69
3:A:603:SO4:O1	5:A:702:HOH:O	2.12	0.66
1:A:265:ASN:OD1	5:A:703:HOH:O	2.14	0.66
1:A:17:PRO:HG2	1:A:144:THR:HB	1.79	0.65
1:A:121:ARG:HB2	1:A:256:ASP:HA	1.79	0.64
1:A:234:LYS:HE3	1:A:251:CYS:HB2	1.82	0.60
1:B:458:GLN:HG2	1:B:467:THR:HG22	1.83	0.60
1:B:79:ALA:HB1	1:B:177:LEU:HD11	1.84	0.60
1:B:221:ARG:HH11	1:B:275:THR:HG21	1.69	0.58
1:A:34:ASN:OD1	1:A:157:TYR:OH	2.08	0.57
1:A:26:VAL:HG22	1:A:127:PHE:HB2	1.86	0.56
1:B:385:HIS:CD2	1:B:386:ILE:HG12	2.39	0.56
1:A:256:ASP:N	1:A:256:ASP:OD1	2.38	0.56
1:A:461:ALA:HB3	1:A:462:ASN:O	2.08	0.54
1:A:68:LYS:O	1:A:72:GLN:HG2	2.08	0.54
1:B:505:SER:HB2	1:B:513:THR:HB	1.90	0.53
1:A:322:TYR:O	1:A:326:THR:HG23	2.09	0.53
1:A:240:GLU:OE1	5:A:704:HOH:O	2.18	0.52
1:A:4:ILE:HG23	1:A:80:ALA:HB1	1.92	0.51
1:B:62:ASP:HA	1:B:228:VAL:HB	1.92	0.50
1:A:19:THR:N	5:A:715:HOH:O	2.45	0.50
1:B:44:ILE:HG23	1:B:83:ILE:HD11	1.94	0.49
1:A:524:ASP:O	1:A:528:ILE:HG23	2.12	0.49
1:B:322:TYR:O	1:B:326:THR:HG23	2.13	0.48
1:B:162:ASP:OD2	1:B:162:ASP:N	2.39	0.48
1:B:37:GLU:HG3	1:B:73:LEU:HD11	1.95	0.48
1:B:68:LYS:O	1:B:72:GLN:HG2	2.14	0.47
1:A:369:SER:HA	5:A:719:HOH:O	2.15	0.47
1:B:500:ILE:HD11	1:B:539:LEU:HD22	1.97	0.47
1:A:360:LYS:HG3	1:A:361:PHE:N	2.30	0.46
1:A:248:ALA:HB1	1:A:251:CYS:SG	2.57	0.45
1:B:211:LEU:HD22	1:B:403:ALA:HB2	1.97	0.45
1:B:385:HIS:NE2	1:B:386:ILE:HG12	2.30	0.45
1:B:7:VAL:O	1:B:157:TYR:N	2.49	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:159:VAL:O	1:B:161:PRO:HD3	2.17	0.44
1:B:86:LEU:HG	1:B:184:PHE:HZ	1.83	0.44
1:A:117:SER:HB3	1:A:118:HIS:H	1.56	0.44
1:B:12:TYR:OH	1:B:32:SER:HB3	2.18	0.44
1:A:17:PRO:HB3	1:A:39:PHE:HZ	1.82	0.43
1:A:428:TYR:CD2	1:A:558:PRO:HG3	2.53	0.43
1:A:117:SER:O	1:A:289:GLY:HA3	2.18	0.43
1:B:169:VAL:HA	5:B:713:HOH:O	2.19	0.43
1:A:561:ILE:CD1	4:A:606:GOL:H32	2.48	0.43
1:B:86:LEU:HG	1:B:184:PHE:CZ	2.54	0.43
1:B:79:ALA:HB2	1:B:184:PHE:CD2	2.54	0.43
1:B:226:HIS:HB3	1:B:254:LEU:HG	2.00	0.43
1:A:438:ALA:HB1	1:A:504:LEU:HD21	2.01	0.42
1:B:333:ARG:HD2	1:B:337:THR:HB	2.01	0.42
1:A:45:SER:HA	1:A:52:ARG:NH2	2.35	0.42
1:B:372:SER:O	1:B:383:SER:HA	2.20	0.42
1:A:523:LYS:HA	1:A:523:LYS:HD2	1.86	0.42
1:A:73:LEU:HD23	1:A:76:ARG:NH1	2.35	0.42
1:A:385:HIS:CD2	1:A:386:ILE:HG12	2.55	0.41
1:B:21:GLY:HA3	1:B:130:LYS:HA	2.01	0.41
1:B:498:SER:HB3	1:B:539:LEU:HD11	2.02	0.41
1:B:256:ASP:OD1	1:B:256:ASP:N	2.53	0.41
1:A:175:PHE:HB2	1:A:184:PHE:HB3	2.02	0.41
1:A:427:ARG:HD2	1:A:517:TYR:CE2	2.55	0.41
1:B:531:ASP:O	1:B:534:VAL:HG12	2.20	0.41
1:B:39:PHE:O	1:B:43:ILE:HG13	2.20	0.41
1:A:34:ASN:HA	1:A:37:GLU:OE1	2.21	0.41
1:B:197:THR:HA	1:B:200:ARG:NH1	2.36	0.41
1:B:89:GLY:HA3	1:B:94:LEU:HG	2.03	0.41
1:A:434:GLU:HG3	1:A:434:GLU:O	2.20	0.41
1:A:478:ASP:HA	1:A:479:PRO:HD3	1.83	0.41
1:B:449:MET:HG2	1:B:492:LEU:HD21	2.03	0.41
1:A:22:LEU:O	1:A:128:GLY:HA2	2.20	0.40
1:B:539:LEU:HD23	1:B:539:LEU:HA	1.88	0.40
1:A:48:GLU:HA	1:A:49:PRO:HD3	1.94	0.40
1:B:199:LEU:HD13	1:B:396:LEU:HD22	2.02	0.40
1:B:537:ALA:HB3	1:B:538:PRO:HD3	2.03	0.40
1:A:221:ARG:NH1	3:A:604:SO4:O3	2.55	0.40
1:A:223:ASP:HA	1:A:249:VAL:HG23	2.03	0.40
1:B:478:ASP:HB3	1:B:481:ASP:OD1	2.21	0.40
1:B:336:PRO:HB2	1:B:517:TYR:CE2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	546/562 (97%)	524 (96%)	18 (3%)	4 (1%)	24	42
1	B	524/562 (93%)	501 (96%)	21 (4%)	2 (0%)	36	57
All	All	1070/1124 (95%)	1025 (96%)	39 (4%)	6 (1%)	27	46

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	18	GLY
1	A	215	PRO
1	A	2	VAL
1	B	17	PRO
1	B	29	PHE
1	A	463	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	425/463 (92%)	412 (97%)	13 (3%)	43	70
1	B	385/463 (83%)	368 (96%)	17 (4%)	31	55
All	All	810/926 (88%)	780 (96%)	30 (4%)	37	64

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	SER
1	A	117	SER
1	A	180	LYS
1	A	213	SER
1	A	221	ARG
1	A	246	ASN
1	A	266	LEU
1	A	271	ASP
1	A	326	THR
1	A	342	ASP
1	A	404	THR
1	A	525	VAL
1	A	530	GLN
1	B	4	ILE
1	B	153	THR
1	B	162	ASP
1	B	169	VAL
1	B	201	SER
1	B	213	SER
1	B	221	ARG
1	B	246	ASN
1	B	249	VAL
1	B	271	ASP
1	B	342	ASP
1	B	349	LYS
1	B	381	THR
1	B	389	LYS
1	B	404	THR
1	B	462	ASN
1	B	550	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	602	-	4,4,4	0.15	0	6,6,6	0.18	0
3	SO4	A	603	-	4,4,4	0.21	0	6,6,6	0.08	0
3	SO4	A	604	-	4,4,4	0.16	0	6,6,6	0.08	0
4	GOL	A	605	-	5,5,5	0.33	0	5,5,5	0.48	0
4	GOL	A	606	-	5,5,5	0.37	0	5,5,5	0.57	0
3	SO4	B	601	-	4,4,4	0.14	0	6,6,6	0.20	0
3	SO4	B	602	-	4,4,4	0.19	0	6,6,6	0.15	0
3	SO4	B	603	-	4,4,4	0.16	0	6,6,6	0.27	0
3	SO4	B	604	-	4,4,4	0.17	0	6,6,6	0.17	0
3	SO4	B	605	-	4,4,4	0.19	0	6,6,6	0.19	0
3	SO4	B	606	-	4,4,4	0.20	0	6,6,6	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	602	-	-	0/0/0/0	0/0/0/0
3	SO4	A	603	-	-	0/0/0/0	0/0/0/0
3	SO4	A	604	-	-	0/0/0/0	0/0/0/0
4	GOL	A	605	-	-	0/4/4/4	0/0/0/0
4	GOL	A	606	-	-	0/4/4/4	0/0/0/0
3	SO4	B	601	-	-	0/0/0/0	0/0/0/0
3	SO4	B	602	-	-	0/0/0/0	0/0/0/0
3	SO4	B	603	-	-	0/0/0/0	0/0/0/0
3	SO4	B	604	-	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	B	605	-	-	0/0/0/0	0/0/0/0
3	SO4	B	606	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	603	SO4	1	0
3	A	604	SO4	1	0
4	A	606	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	550/562 (97%)	-0.18	4 (0%) 87 88	31, 63, 100, 120	0
1	B	536/562 (95%)	0.13	32 (5%) 22 23	44, 87, 139, 159	0
All	All	1086/1124 (96%)	-0.03	36 (3%) 46 50	31, 73, 128, 159	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	549	LEU	4.6
1	B	463	ASP	3.9
1	B	158	ALA	3.7
1	B	546	VAL	3.2
1	B	7	VAL	3.2
1	B	49	PRO	3.2
1	B	150	ILE	3.1
1	B	464	LYS	3.0
1	B	127	PHE	3.0
1	B	539	LEU	3.0
1	B	543	ALA	2.9
1	B	480	VAL	2.8
1	B	542	ILE	2.8
1	B	527	LYS	2.7
1	B	548	GLN	2.5
1	A	1	MET	2.4
1	B	460	SER	2.4
1	B	5	VAL	2.3
1	B	455	VAL	2.3
1	A	454	PHE	2.2
1	B	466	TYR	2.2
1	B	484	ILE	2.2
1	B	468	VAL	2.2
1	A	154	ILE	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	461	ALA	2.2
1	B	453	SER	2.2
1	B	29	PHE	2.2
1	B	462	ASN	2.1
1	B	482	GLY	2.1
1	A	267	THR	2.1
1	B	536	LEU	2.1
1	B	553	THR	2.1
1	B	145	ASP	2.1
1	B	157	TYR	2.0
1	B	544	LEU	2.0
1	B	35	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	B	607	1/1	0.64	0.17	105,105,105,105	0
3	SO4	A	603	5/5	0.70	0.15	151,151,153,155	0
3	SO4	B	606	5/5	0.71	0.15	157,159,160,160	0
3	SO4	B	602	5/5	0.73	0.26	94,95,98,99	5
4	GOL	A	606	6/6	0.73	0.27	57,70,72,73	6
3	SO4	A	604	5/5	0.77	0.30	115,115,117,118	5
3	SO4	B	604	5/5	0.81	0.22	107,108,110,111	5
3	SO4	B	603	5/5	0.82	0.16	93,94,96,97	5
3	SO4	B	605	5/5	0.90	0.14	85,85,89,93	5
4	GOL	A	605	6/6	0.92	0.16	68,80,84,87	0
3	SO4	B	601	5/5	0.95	0.13	89,96,97,99	5

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	A	602	5/5	0.98	0.15	69,69,72,75	0
2	MG	A	601	1/1	0.98	0.12	67,67,67,67	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.