



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 24, 2018 – 01:45 PM EDT

PDB ID : 6F9T  
Title : Crystal structure of human testis Angiotensin-1 converting enzyme in complex with Sampatrilat.  
Authors : Cozier, G.E.; Acharya, K.R.  
Deposited on : 2017-12-15  
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031021  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031021


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2957 (1.60-1.60)
Clashscore	122126	3202 (1.60-1.60)
Ramachandran outliers	120053	3117 (1.60-1.60)
Sidechain outliers	120020	3116 (1.60-1.60)
RSRZ outliers	108989	2883 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	591	<div> <div>2%</div> <div>93%</div> <div>6%</div> </div>

## 2 Entry composition [i](#)

There are 14 unique types of molecules in this entry. The entry contains 10715 atoms, of which 4921 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Angiotensin-converting enzyme.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	586	9577	3117	4696	833	907	24	0	20	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	64	GLY	GLU	conflict	UNP P12821
A	90	GLN	ASN	conflict	UNP P12821
A	155	GLN	ASN	conflict	UNP P12821
A	337	GLN	ASN	conflict	UNP P12821
A	586	GLN	ASN	conflict	UNP P12821

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	A	1	26	8	12	1	5	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
2	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	A	1	Total	C	H	N	O	0	0
			25	8	11	1	5		
2	A	1	Total	C	H	N	O	0	0
			26	8	12	1	5		

- Molecule 3 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula:  $C_6H_{12}O_5$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			20	6	10	4		

- Molecule 4 is BETA-D-MANNOSE (three-letter code: BMA) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			19	6	8	5		

- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			20	6	9	5		
5	A	1	Total	C	H	O	0	0
			20	6	9	5		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		

- Molecule 7 is Sampatrilat (three-letter code: D0Z) (formula:  $C_{26}H_{40}N_4O_9S$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
7	A	1	Total	C	H	N	O	S	0	0
			78	26	38	4	9	1		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	2	Total	Cl	0	0
			2	2		

- Molecule 9 is IMIDAZOLE (three-letter code: IMD) (formula:  $C_3H_5N_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	H	N	0	0
			10	3	5	2		
9	A	1	Total	C	H	N	0	0
			10	3	5	2		

- Molecule 10 is BORIC ACID (three-letter code: BO3) (formula: BH<sub>3</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	A	1	Total	B	H	O	0	0
			7	1	3	3		

- Molecule 11 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	A	1	Total	C	H	O	0	0
			10	2	6	2		
11	A	1	Total	C	H	O	0	0
			10	2	6	2		
11	A	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 12 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
12	A	1	Total	C	H	O	0	0
			17	4	10	3		
12	A	1	Total	C	H	O	0	0
			17	4	10	3		
12	A	1	Total	C	H	O	0	0
			17	4	10	3		

- Molecule 13 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
13	A	1	Total	C	H	O	0	0
			24	6	14	4		


- Molecule 14 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	A	678	Total	O	0	9
			686	686		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Angiotensin-converting enzyme



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.10Å 84.76Å 133.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.63 – 1.60 71.63 – 1.60	Depositor EDS
% Data completeness (in resolution range)	98.8 (71.63-1.60) 98.8 (71.63-1.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.69 (at 1.60Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, $R_{free}$	0.165 , 0.187 0.165 , 0.188	Depositor DCC
$R_{free}$ test set	2215 reflections (2.59%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.4	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 49.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	10715	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PGE, NAG, CL, BO3, CSO, D0Z, BMA, FUC, IMD, EDO, PEG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.47	0/5080	0.62	0/6905

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4881	4696	4619	22	0
2	A	84	76	74	0	0
3	A	10	10	10	0	0
4	A	11	8	8	0	0
5	A	22	18	18	0	0
6	A	1	0	0	0	0
7	A	40	38	0	2	0
8	A	2	0	0	0	0
9	A	10	10	10	1	0
10	A	4	3	3	0	0
11	A	12	18	18	0	0
12	A	21	30	30	5	0
13	A	10	14	14	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
14	A	686	0	0	11	1
All	All	5794	4921	4804	27	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 27 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:624:ASN:ND2	14:A:802:HOH:O	2.22	0.70
1:A:239:GLU:OE2	14:A:801:HOH:O	2.10	0.69
7:A:712:D0Z:N11	14:A:805:HOH:O	2.29	0.64
1:A:617:PRO:HD2	14:A:997:HOH:O	2.03	0.58
1:A:496:CSO:OD	1:A:623:PRO:HD3	2.07	0.55

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
14:A:1051:HOH:O	14:A:1062:HOH:O[1_655]	2.15	0.05

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	604/591 (102%)	595 (98%)	8 (1%)	1 (0%)	49 27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	530/514 (103%)	528 (100%)	2 (0%)	92	86

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	390	TYR
1	A	394	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CSO	A	496	1	4,6,7	1.14	0	1,6,8	1.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	496	1	-	0/1/5/7	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	496	CSO	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 3 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	701	1,2	14,14,15	0.31	0	17,19,21	0.79	1 (5%)
2	NAG	A	702	2	14,14,15	0.31	0	17,19,21	0.85	1 (5%)
2	NAG	A	703	5	14,14,15	0.29	0	17,19,21	0.45	0
2	NAG	A	704	5	14,14,15	0.17	0	17,19,21	0.75	1 (5%)
2	NAG	A	705	1,3,2	14,14,15	0.28	0	17,19,21	0.58	0
2	NAG	A	706	2,4	14,14,15	0.74	1 (7%)	17,19,21	0.48	0
3	FUC	A	707	2	9,10,11	0.80	1 (11%)	13,14,16	0.67	0
4	BMA	A	708	2,5	11,11,12	0.90	0	15,15,17	1.00	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	MAN	A	709	2,4	11,11,12	1.06	0	15,15,17	1.09	1 (6%)
5	MAN	A	710	2,4	11,11,12	0.85	1 (9%)	15,15,17	1.89	2 (13%)
7	D0Z	A	712	6	34,41,41	2.36	9 (26%)	40,57,57	1.54	8 (20%)
9	IMD	A	715	-	5,5,5	1.44	2 (40%)	5,5,5	0.34	0
9	IMD	A	716	-	5,5,5	1.47	2 (40%)	5,5,5	0.23	0
10	BO3	A	717	-	3,3,3	0.19	0	3,3,3	0.09	0
11	EDO	A	718	-	3,3,3	0.46	0	2,2,2	0.36	0
11	EDO	A	719	-	3,3,3	0.47	0	2,2,2	0.37	0
11	EDO	A	720	-	3,3,3	0.51	0	2,2,2	0.28	0
12	PEG	A	721	-	6,6,6	0.55	0	5,5,5	0.34	0
12	PEG	A	722	-	6,6,6	0.50	0	5,5,5	0.41	0
12	PEG	A	723	-	6,6,6	0.50	0	5,5,5	0.21	0
13	PGE	A	724	-	9,9,9	0.36	0	8,8,8	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	701	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	702	2	-	0/6/23/26	0/1/1/1
2	NAG	A	703	5	-	0/6/23/26	0/1/1/1
2	NAG	A	704	5	-	0/6/23/26	0/1/1/1
2	NAG	A	705	1,3,2	-	0/6/23/26	0/1/1/1
2	NAG	A	706	2,4	-	0/6/23/26	0/1/1/1
3	FUC	A	707	2	-	0/0/17/20	0/1/1/1
4	BMA	A	708	2,5	-	0/2/19/22	0/1/1/1
5	MAN	A	709	2,4	-	0/2/19/22	0/1/1/1
5	MAN	A	710	2,4	-	0/2/19/22	0/1/1/1
7	D0Z	A	712	6	-	0/38/55/55	0/2/2/2
9	IMD	A	715	-	-	0/0/0/0	0/1/1/1
9	IMD	A	716	-	-	0/0/0/0	0/1/1/1
10	BO3	A	717	-	-	0/0/0/0	0/0/0/0
11	EDO	A	718	-	-	0/1/1/1	0/0/0/0
11	EDO	A	719	-	-	0/1/1/1	0/0/0/0
11	EDO	A	720	-	-	0/1/1/1	0/0/0/0
12	PEG	A	721	-	-	0/4/4/4	0/0/0/0
12	PEG	A	722	-	-	0/4/4/4	0/0/0/0
12	PEG	A	723	-	-	0/4/4/4	0/0/0/0
13	PGE	A	724	-	-	0/7/7/7	0/0/0/0



The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	712	D0Z	C22-C18	-6.16	1.41	1.54
7	A	712	D0Z	C19-C20	-3.48	1.40	1.52
7	A	712	D0Z	O13-C12	-2.74	1.17	1.23
2	A	706	NAG	O5-C1	-2.71	1.39	1.43
3	A	707	FUC	O5-C1	-2.04	1.40	1.43

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	712	D0Z	O03-S02-O04	-4.89	111.47	118.79
5	A	710	MAN	O2-C2-C3	-4.69	101.05	110.19
7	A	712	D0Z	C16-C15-N14	-2.84	108.69	112.53
5	A	709	MAN	O2-C2-C3	-2.57	105.18	110.19
7	A	712	D0Z	C22-C21-C20	-2.48	99.13	105.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	712	D0Z	2	0
9	A	716	IMD	1	0
12	A	721	PEG	1	0
12	A	722	PEG	2	0
12	A	723	PEG	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	585/591 (98%)	-0.30	11 (1%) 66 66	9, 18, 40, 67	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	621	TRP	4.5
1	A	619	TYR	4.0
1	A	623	PRO	3.6
1	A	437	GLY	3.2
1	A	625	SER	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CSO	A	496	7/8	0.99	0.07	9,13,17,19	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	702	14/15	0.62	0.28	59,80,107,120	0
12	PEG	A	721	7/7	0.69	0.31	37,50,61,63	0
5	MAN	A	709	11/12	0.71	0.18	42,61,77,81	0
11	EDO	A	720	4/4	0.73	0.24	45,54,63,76	0
2	NAG	A	704	14/15	0.76	0.27	74,89,99,101	0
12	PEG	A	723	7/7	0.76	0.26	41,50,58,65	0
11	EDO	A	719	4/4	0.76	0.31	38,48,58,62	0
2	NAG	A	701	14/15	0.76	0.14	45,59,73,76	0
4	BMA	A	708	11/12	0.76	0.15	48,59,75,90	0
2	NAG	A	703	14/15	0.79	0.17	27,46,54,60	28
9	IMD	A	716	5/5	0.79	0.26	39,48,60,61	0
5	MAN	A	710	11/12	0.80	0.21	58,83,104,104	0
11	EDO	A	718	4/4	0.82	0.13	37,45,52,58	0
12	PEG	A	722	7/7	0.86	0.21	34,43,51,54	0
10	BO3	A	717	4/4	0.89	0.10	28,32,38,41	0
3	FUC	A	707	10/11	0.89	0.12	28,41,50,52	0
13	PGE	A	724	10/10	0.90	0.21	29,47,59,64	0
2	NAG	A	706	14/15	0.90	0.12	29,41,61,70	0
2	NAG	A	705	14/15	0.92	0.08	25,36,49,49	0
9	IMD	A	715	5/5	0.95	0.10	17,21,26,27	0
7	D0Z	A	712	40/40	0.97	0.09	10,16,38,39	0
8	CL	A	714	1/1	1.00	0.04	12,12,12,12	0
6	ZN	A	711	1/1	1.00	0.01	6,6,6,6	0
8	CL	A	713	1/1	1.00	0.12	14,14,14,14	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.