



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 12:45 am GMT

PDB ID : 5FP2  
Title : Crystal structure of the siderophore receptor PirA from *Pseudomonas aeruginosa*  
Authors : Moynie, L.; Tortajada, A.; Naismith, J.H.  
Deposited on : 2015-11-27  
Resolution : 2.97 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2374 (3.00-2.96)
Clashscore	122126	2712 (3.00-2.96)
Ramachandran outliers	120053	2626 (3.00-2.96)
Sidechain outliers	120020	2629 (3.00-2.96)
RSRZ outliers	108989	2268 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	725	 15% 50% 18% • 31%
1	B	725	 15% 50% 21% • 26%
2	X	10	 70% 30%
2	Z	10	 90% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	OCT	A	801	-	-	-	X
3	OCT	B	801	-	-	-	X
3	OCT	B	802	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8201 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FERRIC ENTEROBACTIN RECEPTOR PIRA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	499	Total	C	N	O	S	0	0	0
			3908	2454	686	760	8			
1	B	538	Total	C	N	O	S	0	0	0
			4180	2611	740	818	11			


There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP Q9I527
A	-1	ASP	-	expression tag	UNP Q9I527
A	0	ILE	-	expression tag	UNP Q9I527
A	715	LEU	-	expression tag	UNP Q9I527
A	716	SER	-	expression tag	UNP Q9I527
A	717	HIS	-	expression tag	UNP Q9I527
A	718	HIS	-	expression tag	UNP Q9I527
A	719	HIS	-	expression tag	UNP Q9I527
A	720	HIS	-	expression tag	UNP Q9I527
A	721	HIS	-	expression tag	UNP Q9I527
A	722	HIS	-	expression tag	UNP Q9I527
B	-2	MET	-	expression tag	UNP Q9I527
B	-1	ASP	-	expression tag	UNP Q9I527
B	0	ILE	-	expression tag	UNP Q9I527
B	715	LEU	-	expression tag	UNP Q9I527
B	716	SER	-	expression tag	UNP Q9I527
B	717	HIS	-	expression tag	UNP Q9I527
B	718	HIS	-	expression tag	UNP Q9I527
B	719	HIS	-	expression tag	UNP Q9I527
B	720	HIS	-	expression tag	UNP Q9I527
B	721	HIS	-	expression tag	UNP Q9I527
B	722	HIS	-	expression tag	UNP Q9I527

- Molecule 2 is a protein called FERRIC ENTEROBACTIN RECEPTOR PIRA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	X	7	Total	C	N	O	0	0	0
			33	19	7	7			
2	Z	10	Total	C	N	O	0	0	0
			46	26	10	10			

- Molecule 3 is N-OCTANE (three-letter code: OCT) (formula:  $C_8H_{18}$ ).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	C	0	0
			8	8		
3	B	1	Total	C	0	0
			8	8		
3	B	1	Total	C	0	0
			8	8		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		







## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	183.12Å 95.28Å 131.83Å 90.00° 127.52° 90.00°	Depositor
Resolution (Å)	52.49 – 2.97 52.49 – 2.97	Depositor EDS
% Data completeness (in resolution range)	98.8 (52.49-2.97) 98.8 (52.49-2.97)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.99 (at 2.96Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.263 , 0.319 0.262 , 0.311	Depositor DCC
$R_{free}$ test set	1756 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	73.4	Xtriage
Anisotropy	0.515	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 97.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	8201	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	128.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, OCT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.42	0/3982	0.61	0/5394
1	B	0.48	0/4257	0.66	0/5767
All	All	0.45	0/8239	0.64	0/11161

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	530	TYR	Sidechain
1	B	142	TYR	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3908	0	3813	91	0
1	B	4180	0	4077	141	0
2	X	33	0	8	0	0
2	Z	46	0	9	1	0
3	A	8	0	18	0	0
3	B	16	0	36	0	0
4	B	10	0	0	0	0
All	All	8201	0	7961	226	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 226 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:295:TYR:OH	1:B:193:THR:HG21	1.48	1.11
1:A:193:THR:HG21	1:B:295:TYR:OH	1.57	1.02
1:B:246:THR:HG22	1:B:249:GLN:HB2	1.41	1.01
1:B:484:ARG:NH1	1:B:495:GLY:O	2.11	0.83
1:B:433:LEU:CD2	1:B:443:TRP:CD1	2.65	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	475/725 (66%)	455 (96%)	19 (4%)	1 (0%)	49	83
1	B	518/725 (71%)	493 (95%)	24 (5%)	1 (0%)	49	83
All	All	993/1450 (68%)	948 (96%)	43 (4%)	2 (0%)	49	83

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	141	ARG
1	B	361	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	424/604 (70%)	387 (91%)	37 (9%)	11	38
1	B	454/604 (75%)	407 (90%)	47 (10%)	8	28
All	All	878/1208 (73%)	794 (90%)	84 (10%)	9	33

5 of 84 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	32	LYS
1	B	208	ASP
1	B	620	GLU
1	B	98	LYS
1	B	130	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	244	GLN
1	A	249	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	OCT	A	801	-	7,7,7	0.23	0	6,6,6	0.39	0
4	SO4	B	1716	-	4,4,4	0.32	0	6,6,6	0.25	0
4	SO4	B	1717	-	4,4,4	0.40	0	6,6,6	0.15	0
3	OCT	B	801	-	7,7,7	0.23	0	6,6,6	0.38	0
3	OCT	B	802	-	7,7,7	0.21	0	6,6,6	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OCT	A	801	-	-	0/5/5/5	0/0/0/0
4	SO4	B	1716	-	-	0/0/0/0	0/0/0/0
4	SO4	B	1717	-	-	0/0/0/0	0/0/0/0
3	OCT	B	801	-	-	0/5/5/5	0/0/0/0
3	OCT	B	802	-	-	0/5/5/5	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	499/725 (68%)	1.29	106 (21%) <b>1</b> <b>0</b>	101, 126, 164, 197	0
1	B	538/725 (74%)	1.44	110 (20%) <b>1</b> <b>0</b>	103, 124, 159, 191	0
2	X	0/10	-	-	-	-
2	Z	0/10	-	-	-	-
All	All	1037/1470 (70%)	1.37	216 (20%) <b>1</b> <b>0</b>	101, 125, 161, 197	0

The worst 5 of 216 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	285	ALA	10.4
1	B	263	ILE	9.3
1	B	210	ASP	7.8
1	A	361	LEU	7.6
1	A	287	THR	7.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	OCT	B	802	8/8	0.61	0.69	160,160,160,160	0
4	SO4	B	1716	5/5	0.75	0.39	160,160,160,160	0
3	OCT	A	801	8/8	0.77	0.45	160,160,160,160	0
3	OCT	B	801	8/8	0.78	0.59	160,160,160,160	0
4	SO4	B	1717	5/5	0.79	0.23	160,160,160,160	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.