



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 12:18 am GMT

PDB ID : 1FT8
Title : CRYSTAL STRUCTURE OF THE RNA-BINDING DOMAIN OF THE
MRNA EXPORT FACTOR TAP
Authors : Liker, E.; Fernandez, E.; Izaurralde, E.; Conti, E.
Deposited on : 2000-09-12
Resolution : 3.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	trunk30967
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	1522 (3.20-3.12)
Ramachandran outliers	120053	1493 (3.20-3.12)
Sidechain outliers	120020	1492 (3.20-3.12)
RSRZ outliers	108989	1344 (3.20-3.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	
1	B	271	
1	C	271	
1	D	271	
1	E	271	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 6913 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TIP ASSOCIATING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	242	Total	C	N	O	S	0	0	0
			1961	1245	343	366	7			
1	B	165	Total	C	N	O	S	0	0	0
			1323	831	232	255	5			
1	C	244	Total	C	N	O	S	0	0	0
			1973	1253	346	367	7			
1	D	162	Total	C	N	O	S	0	0	0
			1302	818	229	250	5			
1	E	44	Total	C	N	O	S	0	0	0
			354	236	54	62	2			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	119	TRP	CYS	ENGINEERED	UNP Q9UBU9
A	226	VAL	ALA	ENGINEERED	UNP Q9UBU9
B	119	TRP	CYS	ENGINEERED	UNP Q9UBU9
B	226	VAL	ALA	ENGINEERED	UNP Q9UBU9
C	119	TRP	CYS	ENGINEERED	UNP Q9UBU9
C	226	VAL	ALA	ENGINEERED	UNP Q9UBU9
D	119	TRP	CYS	ENGINEERED	UNP Q9UBU9
D	226	VAL	ALA	ENGINEERED	UNP Q9UBU9
E	119	TRP	CYS	ENGINEERED	UNP Q9UBU9
E	226	VAL	ALA	ENGINEERED	UNP Q9UBU9

• Molecule 1: TIP ASSOCIATING PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	139.92Å 139.92Å 206.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.15 29.94 – 3.16	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-3.15) 99.6 (29.94-3.16)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.10 (at 3.18Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.303 , 0.303 0.299 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	80.0	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 35.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	6913	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.73	4/1996 (0.2%)	0.91	10/2696 (0.4%)
1	B	0.60	1/1340 (0.1%)	0.79	2/1808 (0.1%)
1	C	0.75	7/2008 (0.3%)	0.98	8/2711 (0.3%)
1	D	0.61	1/1318 (0.1%)	0.92	8/1777 (0.5%)
1	E	0.72	0/363	0.95	2/489 (0.4%)
All	All	0.69	13/7025 (0.2%)	0.92	30/9481 (0.3%)

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	165	ASP	CB-CG	-11.57	1.27	1.51
1	C	228	ASP	CB-CG	-10.18	1.30	1.51
1	A	164	GLU	CB-CG	-10.11	1.32	1.52
1	C	165	ASP	CB-CG	-9.04	1.32	1.51
1	C	164	GLU	CB-CG	-7.82	1.37	1.52

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	228	ASP	CB-CG-OD1	-19.55	100.71	118.30
1	A	165	ASP	CB-CG-OD2	-18.87	101.32	118.30
1	C	165	ASP	CB-CG-OD2	-18.30	101.83	118.30
1	C	155	GLU	OE1-CD-OE2	-8.83	112.70	123.30
1	C	165	ASP	CB-CG-OD1	8.22	125.70	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1961	0	2003	171	0
1	B	1323	0	1363	107	0
1	C	1973	0	2013	135	0
1	D	1302	0	1345	113	0
1	E	354	0	349	50	1
All	All	6913	0	7073	536	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

The worst 5 of 536 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:230:LYS:HD2	1:A:273:SER:OG	1.29	1.32
1:B:323:ASP:HB2	1:C:177:TYR:CE2	1.78	1.19
1:C:119:TRP:HZ3	1:C:150:ILE:CD1	1.56	1.19
1:E:144:SER:O	1:E:145:VAL:HG23	1.39	1.19
1:E:145:VAL:HG22	1:E:146:PRO:CD	1.73	1.17

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:142:LYS:NZ	1:E:142:LYS:NZ[8_665]	1.44	0.76

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	238/271 (88%)	213 (90%)	20 (8%)	5 (2%)	8	37
1	B	163/271 (60%)	149 (91%)	10 (6%)	4 (2%)	6	33
1	C	240/271 (89%)	216 (90%)	17 (7%)	7 (3%)	5	29
1	D	160/271 (59%)	139 (87%)	16 (10%)	5 (3%)	4	27
1	E	38/271 (14%)	35 (92%)	2 (5%)	1 (3%)	6	31
All	All	839/1355 (62%)	752 (90%)	65 (8%)	22 (3%)	6	31

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	236	PRO
1	C	157	THR
1	C	236	PRO
1	C	238	LEU
1	C	308	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/245 (91%)	201 (90%)	21 (10%)	9	33
1	B	152/245 (62%)	143 (94%)	9 (6%)	21	57
1	C	222/245 (91%)	205 (92%)	17 (8%)	14	46
1	D	150/245 (61%)	137 (91%)	13 (9%)	11	38
1	E	39/245 (16%)	34 (87%)	5 (13%)	5	21
All	All	785/1225 (64%)	720 (92%)	65 (8%)	12	42

5 of 65 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	367	PRO
1	C	172	LEU
1	E	121	LYS
1	C	122	ILE
1	C	156	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	263	ASN
1	B	297	ASN
1	D	297	ASN
1	B	271	ASN
1	C	156	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	242/271 (89%)	-0.00	6 (2%) 57 42	55, 66, 89, 109	0
1	B	165/271 (60%)	0.00	6 (3%) 42 26	49, 63, 82, 97	0
1	C	244/271 (90%)	0.15	5 (2%) 65 49	55, 74, 106, 106	0
1	D	162/271 (59%)	0.12	7 (4%) 35 20	52, 65, 85, 100	0
1	E	44/271 (16%)	0.97	8 (18%) 1 0	89, 94, 94, 94	0
All	All	857/1355 (63%)	0.11	32 (3%) 41 26	49, 68, 106, 109	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	239	VAL	4.2
1	E	164	GLU	4.0
1	B	367	PRO	3.3
1	A	242	ASN	3.1
1	D	365	GLU	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.