



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 11:27 pm GMT

PDB ID : 2FTK  
Title : beryllofluoride Spo0F complex with Spo0B  
Authors : Varughese, K.I.  
Deposited on : 2006-01-24  
Resolution : 3.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1497 (3.10-3.02)
Clashscore	122126	1601 (3.10-3.02)
Ramachandran outliers	120053	1548 (3.10-3.02)
Sidechain outliers	120020	1547 (3.10-3.02)
RSRZ outliers	108989	1462 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	192	
1	B	192	
1	C	192	
1	D	192	
2	E	124	
2	F	124	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	124	
2	H	124	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9834 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sporulation initiation phosphotransferase B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	181	Total	C	N	O	S	0	0	0
			1497	955	251	284	7			
1	B	182	Total	C	N	O	S	6	0	0
			1505	959	253	286	7			
1	C	181	Total	C	N	O	S	0	0	0
			1497	955	251	284	7			
1	D	182	Total	C	N	O	S	6	0	0
			1505	959	253	286	7			

- Molecule 2 is a protein called Sporulation initiation phosphotransferase F.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
2	E	119	Total	Be	C	F	N	O	S	0	0	0
			954	1	609	3	156	180	5			
2	F	119	Total	Be	C	F	N	O	S	0	0	0
			954	1	609	3	156	180	5			
2	G	119	Total	Be	C	F	N	O	S	0	0	0
			954	1	609	3	156	180	5			
2	H	119	Total	Be	C	F	N	O	S	0	1	0
			964	1	615	3	159	181	5			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	1213	SER	TYR	ENGINEERED	UNP P06628
E	1254	BFD	ASP	MODIFIED RESIDUE	UNP P06628
F	1013	SER	TYR	ENGINEERED	UNP P06628
F	1054	BFD	ASP	MODIFIED RESIDUE	UNP P06628
G	1613	SER	TYR	ENGINEERED	UNP P06628
G	1654	BFD	ASP	MODIFIED RESIDUE	UNP P06628
H	1413	SER	TYR	ENGINEERED	UNP P06628
H	1454	BFD	ASP	MODIFIED RESIDUE	UNP P06628


- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	1	Total 1	Mg 1	0	0
3	G	1	Total 1	Mg 1	0	0
3	F	1	Total 1	Mg 1	0	0
3	E	1	Total 1	Mg 1	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Sporulation initiation phosphotransferase B




#### • Molecule 1: Sporulation initiation phosphotransferase B




#### • Molecule 1: Sporulation initiation phosphotransferase B




- Molecule 1: Sporulation initiation phosphotransferase B




- Molecule 2: Sporulation initiation phosphotransferase F




- Molecule 2: Sporulation initiation phosphotransferase F



- Molecule 2: Sporulation initiation phosphotransferase F



- Molecule 2: Sporulation initiation phosphotransferase F





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.47Å 118.33Å 168.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.86 – 3.05 19.86 – 3.05	Depositor EDS
% Data completeness (in resolution range)	98.6 (19.86-3.05) 98.7 (19.86-3.05)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.21 (at 3.04Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.232 , 0.282 0.222 , 0.273	Depositor DCC
$R_{free}$ test set	1404 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	74.1	Xtriage
Anisotropy	0.113	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 48.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	9834	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BFD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.57	0/1530	0.67	0/2064
1	B	0.54	0/1538	0.71	0/2075
1	C	0.45	0/1530	0.66	1/2064 (0.0%)
1	D	0.44	0/1538	0.63	0/2075
2	E	0.38	0/952	0.61	0/1279
2	F	0.46	0/952	0.66	0/1279
2	G	0.32	0/952	0.61	0/1279
2	H	0.35	0/963	0.60	0/1294
All	All	0.46	0/9955	0.65	1/13409 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	548	LEU	CA-CB-CG	5.22	127.30	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	489	TYR	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1497	0	1457	86	0
1	B	1505	0	1463	80	0
1	C	1497	0	1457	130	0
1	D	1505	0	1463	94	0
2	E	954	0	986	43	0
2	F	954	0	986	43	0
2	G	954	0	986	136	0
2	H	964	0	992	51	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
All	All	9834	0	9790	618	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 618 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:G:1687:LEU:HA	2:G:1690:ILE:HD13	1.26	1.10
2:E:1251:VAL:HG11	2:E:1269:MET:HE1	1.40	1.02
1:C:565:ILE:H	1:C:565:ILE:HD12	1.25	1.01
2:G:1620:ASN:ND2	2:G:1632:GLN:HE21	1.60	0.98
1:B:381:THR:HG23	1:C:581:THR:HG23	1.48	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	179/192 (93%)	165 (92%)	12 (7%)	2 (1%)	16	46
1	B	180/192 (94%)	166 (92%)	13 (7%)	1 (1%)	27	60
1	C	179/192 (93%)	165 (92%)	12 (7%)	2 (1%)	16	46
1	D	180/192 (94%)	168 (93%)	11 (6%)	1 (1%)	27	60
2	E	116/124 (94%)	100 (86%)	15 (13%)	1 (1%)	19	52
2	F	116/124 (94%)	99 (85%)	14 (12%)	3 (3%)	6	25
2	G	116/124 (94%)	84 (72%)	20 (17%)	12 (10%)	0	3
2	H	117/124 (94%)	100 (86%)	14 (12%)	3 (3%)	6	25
All	All	1183/1264 (94%)	1047 (88%)	111 (9%)	25 (2%)	8	29

5 of 25 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	1320	PRO
2	F	1120	PRO
2	G	1684	TYR
2	G	1700	THR
2	H	1520	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/177 (94%)	155 (93%)	11 (7%)	18	48
1	B	167/177 (94%)	152 (91%)	15 (9%)	10	34
1	C	166/177 (94%)	149 (90%)	17 (10%)	8	27
1	D	167/177 (94%)	151 (90%)	16 (10%)	9	31
2	E	104/109 (95%)	98 (94%)	6 (6%)	22	53
2	F	104/109 (95%)	99 (95%)	5 (5%)	28	61

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	G	104/109 (95%)	97 (93%)	7 (7%)	18	48
2	H	105/109 (96%)	97 (92%)	8 (8%)	14	43
All	All	1083/1144 (95%)	998 (92%)	85 (8%)	14	42

5 of 85 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	564	ASP
1	D	647	LYS
2	H	1466	LEU
1	C	566	ARG
1	D	614	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 51 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	475	HIS
1	D	630	HIS
2	H	1432	GLN
1	C	488	HIS
1	D	741	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BFD	E	1254	3,2	9,11,12	1.57	3 (33%)	5,15,17	1.70	1 (20%)
2	BFD	F	1054	3,2	9,11,12	1.44	1 (11%)	5,15,17	1.80	1 (20%)
2	BFD	G	1654	3,2	9,11,12	1.37	0	5,15,17	1.66	1 (20%)
2	BFD	H	1454	3,2	9,11,12	1.36	1 (11%)	5,15,17	1.75	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BFD	E	1254	3,2	-	0/4/11/13	0/0/0/0
2	BFD	F	1054	3,2	-	0/4/11/13	0/0/0/0
2	BFD	G	1654	3,2	-	0/4/11/13	0/0/0/0
2	BFD	H	1454	3,2	-	0/4/11/13	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1254	BFD	F3-BE	-2.32	1.48	1.54
2	E	1254	BFD	F1-BE	-2.20	1.48	1.54
2	F	1054	BFD	F3-BE	-2.18	1.48	1.54
2	H	1454	BFD	CA-C	2.09	1.53	1.50
2	E	1254	BFD	CA-C	2.57	1.53	1.50

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1054	BFD	OD2-CG-CB	-3.44	116.98	124.70
2	H	1454	BFD	OD2-CG-CB	-3.26	117.38	124.70
2	E	1254	BFD	OD2-CG-CB	-3.15	117.64	124.70
2	G	1654	BFD	OD2-CG-CB	-3.09	117.77	124.70

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	1254	BFD	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	1054	BFD	1	0
2	G	1654	BFD	3	0
2	H	1454	BFD	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	181/192 (94%)	-0.48	2 (1%) 80 61	28, 44, 74, 91	0
1	B	182/192 (94%)	-0.47	1 (0%) 90 78	24, 46, 75, 96	3 (1%)
1	C	181/192 (94%)	-0.23	3 (1%) 70 46	42, 71, 100, 115	0
1	D	182/192 (94%)	-0.25	6 (3%) 46 23	41, 63, 98, 114	3 (1%)
2	E	118/124 (95%)	-0.13	4 (3%) 45 22	56, 78, 96, 106	0
2	F	118/124 (95%)	-0.42	1 (0%) 86 69	36, 55, 77, 87	0
2	G	118/124 (95%)	0.20	5 (4%) 36 17	75, 101, 114, 116	0
2	H	118/124 (95%)	0.02	4 (3%) 45 22	56, 87, 112, 121	1 (0%)
All	All	1198/1264 (94%)	-0.25	26 (2%) 62 37	24, 65, 107, 121	7 (0%)

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	G	1720	PRO	6.7
1	C	413	SER	5.0
1	D	613	SER	4.8
2	H	1521	LEU	4.8
1	C	412	ILE	4.6

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	BFD	H	1454	12/13	0.92	0.29	66,68,71,71	4

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BFD	G	1654	12/13	0.93	0.17	85,87,88,88	4
2	BFD	E	1254	12/13	0.95	0.23	60,62,66,67	4
2	BFD	F	1054	12/13	0.96	0.27	39,41,42,43	4

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MG	G	2004	1/1	0.95	0.05	37,37,37,37	0
3	MG	H	2003	1/1	0.97	0.18	37,37,37,37	0
3	MG	F	2002	1/1	0.99	0.34	37,37,37,37	0
3	MG	E	2001	1/1	1.00	0.28	37,37,37,37	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.