



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2019 – 07:10 AM EDT

PDB ID : 1GO2
Title : Structure of Ferredoxin-NADP+ Reductase with Lys 72 replaced by Glu (K72E)
Authors : Hermoso, J.A.; Mayoral, T.; Medina, M.; Sanz-Aparicio, J.; Gomez-Moreno, C.
Deposited on : 2001-10-15
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.0 (224370), CSD as540be (2019)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20031633


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	4167 (1.70-1.70)
Ramachandran outliers	120053	4100 (1.70-1.70)
Sidechain outliers	120020	4100 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	304	 84% 12% ...

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2702 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FERREDONIN-NADP+ REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	295	Total	C	N	O	S	0	0	0
			2338	1487	398	444	9			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	72	GLU	LYS	engineered mutation	UNP P21890

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	306	Total	O	0	0
			306	306		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: FERREDOXIN-NADP+ REDUCTASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	86.64Å 86.64Å 96.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 1.70	Depositor
% Data completeness (in resolution range)	99.6 (10.00-1.70)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.843	Depositor
R, R_{free}	0.220 , 0.250	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2702	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/2394	0.48	0/3244

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	106	HIS	Peptide
1	A	114	TYR	Sidechain
1	A	79	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2338	0	2307	24	0
2	A	53	0	31	0	0
3	A	5	0	0	0	0
4	A	306	0	0	3	0
All	All	2702	0	2338	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:243:HIS:HD2	1:A:246:GLN:HE22	1.31	0.78
1:A:257:HIS:HD2	1:A:299:HIS:HE1	1.37	0.71
1:A:257:HIS:HD2	1:A:299:HIS:CE1	2.15	0.64
1:A:15:TYR:HD2	1:A:52:LEU:HD23	1.63	0.64
1:A:257:HIS:CD2	1:A:299:HIS:HE1	2.15	0.63
1:A:149:ASN:HD22	1:A:186:PHE:H	1.51	0.58
1:A:243:HIS:HD2	1:A:246:GLN:NE2	2.02	0.55
1:A:243:HIS:CD2	1:A:246:GLN:HE22	2.21	0.55
1:A:243:HIS:HE1	4:A:2245:HOH:O	1.91	0.53
1:A:105:LYS:HD2	1:A:106:HIS:H	1.77	0.49
1:A:284:THR:HG23	4:A:2282:HOH:O	2.13	0.49
1:A:103:GLU:HA	1:A:113:VAL:O	2.16	0.45
1:A:60:ILE:HD13	1:A:95:ILE:HD13	1.98	0.45
1:A:152:MET:HE1	1:A:187:SER:HB2	1.97	0.45
1:A:28:ASN:ND2	1:A:41:GLN:HE21	2.16	0.43
1:A:139:GLU:HB3	4:A:2171:HOH:O	2.19	0.43
1:A:254:GLN:HG3	1:A:255:LYS:HG2	2.02	0.42
1:A:262:GLY:O	1:A:302:THR:HA	2.19	0.42
1:A:145:ASP:HA	1:A:146:PRO:HD3	1.88	0.42
1:A:190:VAL:HG12	1:A:236:ILE:HG13	2.02	0.41
1:A:243:HIS:CD2	1:A:246:GLN:NE2	2.85	0.41
1:A:218:LEU:HD21	1:A:220:TYR:OH	2.21	0.41
1:A:39:ILE:HD12	1:A:101:GLN:NE2	2.36	0.41
1:A:289:GLN:O	1:A:293:LYS:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	293/304 (96%)	286 (98%)	5 (2%)	2 (1%)	24 9

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	106	HIS
1	A	105	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	250/256 (98%)	243 (97%)	7 (3%)	47 27

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	VAL
1	A	28	ASN
1	A	105	LYS
1	A	167	TRP
1	A	207	GLU
1	A	255	LYS
1	A	294	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	ASN
1	A	149	ASN
1	A	182	GLN
1	A	243	HIS
1	A	246	GLN
1	A	254	GLN
1	A	257	HIS
1	A	289	GLN
1	A	299	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	1304	-	50,58,58	1.73	10 (20%)	58,89,89	2.32	13 (22%)
3	SO4	A	1305	-	4,4,4	0.44	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	1304	-	-	0/30/50/50	0/6/6/6
3	SO4	A	1305	-	-	0/0/0/0	0/0/0/0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1304	FAD	C5'-C4'	-3.83	1.46	1.51
2	A	1304	FAD	PA-O5B	-2.18	1.50	1.59
2	A	1304	FAD	C6-C7	-2.07	1.32	1.37
2	A	1304	FAD	P-O1P	-2.05	1.43	1.50
2	A	1304	FAD	O4B-C1B	2.03	1.44	1.41
2	A	1304	FAD	C2A-N3A	2.53	1.36	1.32
2	A	1304	FAD	C4X-C10	3.32	1.42	1.38
2	A	1304	FAD	C4-N3	3.58	1.39	1.33
2	A	1304	FAD	C1'-N10	4.49	1.53	1.48
2	A	1304	FAD	C4-C4X	5.62	1.51	1.41

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1304	FAD	C4X-C4-N3	-5.41	115.93	123.47
2	A	1304	FAD	O2'-C2'-C3'	-3.92	99.53	109.11
2	A	1304	FAD	O3'-C3'-C4'	-3.20	101.07	108.84
2	A	1304	FAD	C9A-N10-C10	-2.49	118.52	121.77
2	A	1304	FAD	C4B-O4B-C1B	-2.19	107.55	109.83
2	A	1304	FAD	C4-C4X-C10	-2.18	118.34	119.95
2	A	1304	FAD	C4-C4X-N5	2.01	120.81	118.59
2	A	1304	FAD	O2A-PA-O1A	2.08	122.64	112.21
2	A	1304	FAD	C1'-C2'-C3'	2.31	116.24	109.79
2	A	1304	FAD	C4X-N5-C5X	4.48	121.37	116.77
2	A	1304	FAD	C5A-C6A-N6A	4.61	127.61	120.38
2	A	1304	FAD	C1'-N10-C10	5.11	123.47	118.46
2	A	1304	FAD	C4-N3-C2	10.29	123.83	115.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.