



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 02:34 am GMT

PDB ID : 3GXR  
Title : The crystal structure of g-type lysozyme from Atlantic cod (*Gadus morhua* L.) in complex with NAG oligomers sheds new light on substrate binding and the catalytic mechanism. Structure with NAG to 1.7  
Authors : Helland, R.; Larsen, R.L.; Finstad, S.; Kyomuhendo, P.; Larsen, A.N.  
Deposited on : 2009-04-02  
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	3793 (1.70-1.70)
Clashscore	122126	4167 (1.70-1.70)
Ramachandran outliers	120053	4100 (1.70-1.70)
Sidechain outliers	120020	4100 (1.70-1.70)
RSRZ outliers	108989	3718 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	187	<div> <div>%</div> <div> <div></div> <div>84%</div> <div>13%</div> <div>...</div> </div> </div>
1	B	187	<div> <div>2%</div> <div> <div></div> <div>81%</div> <div>14%</div> <div>..</div> </div> </div>
1	C	187	<div> <div>3%</div> <div> <div></div> <div>76%</div> <div>14%</div> <div>7%</div> </div> </div>
1	D	187	<div> <div>%</div> <div> <div></div> <div>75%</div> <div>22%</div> <div>..</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6211 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Goose-type lysozyme 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	185	Total	C	N	O	S	0	0	0
			1458	907	261	287	3			
1	B	183	Total	C	N	O	S	0	0	0
			1442	896	259	284	3			
1	C	174	Total	C	N	O	S	0	0	0
			1369	849	247	270	3			
1	D	186	Total	C	N	O	S	0	0	0
			1462	909	262	288	3			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	147	ALA	GLY	SEE REMARK 999	UNP B9TU22
B	147	ALA	GLY	SEE REMARK 999	UNP B9TU22
C	147	ALA	GLY	SEE REMARK 999	UNP B9TU22
D	147	ALA	GLY	SEE REMARK 999	UNP B9TU22

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			15	8	1	6		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			15	8	1	6		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			15	8	1	6		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			15	8	1	6		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			15	8	1	6		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	87	Total 87	O 87	0	0
3	B	78	Total 78	O 78	0	0
3	C	67	Total 67	O 67	0	0
3	D	61	Total 61	O 61	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Goose-type lysozyme 1

Chain A: 




#### • Molecule 1: Goose-type lysozyme 1

Chain B: 





#### • Molecule 1: Goose-type lysozyme 1

Chain C: 



#### • Molecule 1: Goose-type lysozyme 1

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.19Å 75.52Å 78.29Å 90.00° 93.25° 90.00°	Depositor
Resolution (Å)	34.00 – 1.70 34.00 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.2 (34.00-1.70) 99.3 (34.00-1.70)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.57 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.4.0069	Depositor
R, $R_{free}$	0.204 , 0.252 0.202 , 0.251	Depositor DCC
$R_{free}$ test set	3526 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.4	Xtriage
Anisotropy	0.251	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 53.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6211	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.01	1/1489 (0.1%)	0.95	3/2011 (0.1%)
1	B	1.02	2/1472 (0.1%)	0.89	1/1988 (0.1%)
1	C	0.94	2/1394 (0.1%)	0.89	2/1876 (0.1%)
1	D	0.98	0/1493	0.85	0/2016
All	All	0.99	5/5848 (0.1%)	0.90	6/7891 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	106	GLU	CG-CD	6.94	1.62	1.51
1	B	12	GLU	CG-CD	6.40	1.61	1.51
1	A	173	TYR	CD2-CE2	6.30	1.48	1.39
1	B	184	TYR	CD2-CE2	5.34	1.47	1.39
1	C	106	GLU	CD-OE2	5.33	1.31	1.25

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	91	ASN	N-CA-C	7.01	129.93	111.00
1	C	5	TYR	N-CA-C	6.33	128.09	111.00
1	A	103	ARG	NE-CZ-NH2	-6.13	117.23	120.30
1	A	6	GLY	N-CA-C	-5.42	99.54	113.10
1	B	72	ARG	NE-CZ-NH1	5.24	122.92	120.30

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1458	0	1397	19	0
1	B	1442	0	1385	30	0
1	C	1369	0	1319	28	0
1	D	1462	0	1400	43	0
2	A	72	0	66	3	0
2	B	43	0	39	7	0
2	D	72	0	66	5	0
3	A	87	0	0	1	0
3	B	78	0	0	2	0
3	C	67	0	0	4	0
3	D	61	0	0	7	0
All	All	6211	0	5672	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 124 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:91:ASN:HB3	3:D:244:HOH:O	1.59	1.01
1:B:101:ASP:OD1	1:B:103:ARG:HD3	1.74	0.87
1:C:4:GLY:HA3	1:C:5:TYR:HB2	1.56	0.86
1:D:98:MET:H	1:D:116:HIS:HE1	1.21	0.84
1:B:98:MET:H	1:B:116:HIS:HE1	1.30	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	183/187 (98%)	177 (97%)	5 (3%)	1 (0%)	31	14
1	B	181/187 (97%)	177 (98%)	4 (2%)	0	100	100
1	C	170/187 (91%)	163 (96%)	6 (4%)	1 (1%)	27	11
1	D	184/187 (98%)	180 (98%)	4 (2%)	0	100	100
All	All	718/748 (96%)	697 (97%)	19 (3%)	2 (0%)	43	25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	91	ASN
1	C	4	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	153/154 (99%)	148 (97%)	5 (3%)	41	20
1	B	152/154 (99%)	148 (97%)	4 (3%)	49	30
1	C	143/154 (93%)	140 (98%)	3 (2%)	56	38
1	D	153/154 (99%)	147 (96%)	6 (4%)	35	15
All	All	601/616 (98%)	583 (97%)	18 (3%)	44	24

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	159	GLU
1	C	3	VAL
1	D	22	ARG
1	B	106	GLU
1	B	108	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	143	GLN
1	C	39	GLN
1	D	119	GLN
1	B	175	ASN
1	C	23	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	201	2	15,15,15	0.65	0	21,21,21	1.75	5 (23%)
2	NAG	A	202	2	14,14,15	1.01	0	19,19,21	1.86	4 (21%)
2	NAG	A	301	2	15,15,15	1.10	1 (6%)	21,21,21	1.08	2 (9%)
2	NAG	A	302	2	14,14,15	1.28	3 (21%)	19,19,21	1.37	4 (21%)
2	NAG	A	303	2	14,14,15	0.80	0	19,19,21	1.57	4 (21%)
2	NAG	B	301	2	15,15,15	0.73	0	21,21,21	1.82	6 (28%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	302	2	14,14,15	0.87	1 (7%)	19,19,21	1.84	4 (21%)
2	NAG	B	303	2	14,14,15	0.75	1 (7%)	19,19,21	1.42	4 (21%)
2	NAG	D	201	2	15,15,15	0.74	0	21,21,21	2.29	6 (28%)
2	NAG	D	202	2	14,14,15	0.79	0	19,19,21	1.10	1 (5%)
2	NAG	D	301	2	15,15,15	0.81	0	21,21,21	1.54	6 (28%)
2	NAG	D	302	2	14,14,15	0.86	0	19,19,21	1.56	2 (10%)
2	NAG	D	303	2	14,14,15	0.75	1 (7%)	19,19,21	1.70	4 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	201	2	-	0/6/26/26	0/1/1/1
2	NAG	A	202	2	-	0/6/22/26	0/1/1/1
2	NAG	A	301	2	-	0/6/26/26	0/1/1/1
2	NAG	A	302	2	-	0/6/22/26	0/1/1/1
2	NAG	A	303	2	-	0/6/22/26	0/1/1/1
2	NAG	B	301	2	-	0/6/26/26	0/1/1/1
2	NAG	B	302	2	-	0/6/22/26	0/1/1/1
2	NAG	B	303	2	-	0/6/22/26	0/1/1/1
2	NAG	D	201	2	-	0/6/26/26	0/1/1/1
2	NAG	D	202	2	-	0/6/22/26	0/1/1/1
2	NAG	D	301	2	-	0/6/26/26	0/1/1/1
2	NAG	D	302	2	-	0/6/22/26	0/1/1/1
2	NAG	D	303	2	-	0/6/22/26	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	302	NAG	O5-C5	-2.47	1.40	1.44
2	B	302	NAG	O5-C5	-2.32	1.40	1.44
2	B	303	NAG	O5-C5	-2.19	1.40	1.44
2	D	303	NAG	O5-C5	-2.03	1.40	1.44
2	A	302	NAG	C6-C5	2.09	1.56	1.50

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	302	NAG	O5-C1-C2	-5.16	104.33	109.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	201	NAG	C6-C5-C4	-4.30	102.83	112.99
2	B	301	NAG	C3-C2-N2	-4.04	102.82	110.58
2	D	302	NAG	O5-C1-C2	-3.74	105.76	109.52
2	D	302	NAG	C1-C2-N2	-3.71	106.43	110.73

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	NAG	2	0
2	A	302	NAG	2	0
2	B	301	NAG	3	0
2	B	302	NAG	4	0
2	B	303	NAG	4	0
2	D	201	NAG	3	0
2	D	202	NAG	1	0
2	D	301	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	185/187 (98%)	0.08	1 (0%) 90 92	12, 19, 29, 35	0
1	B	183/187 (97%)	0.02	3 (1%) 72 76	12, 19, 31, 37	0
1	C	174/187 (93%)	0.06	6 (3%) 45 50	13, 21, 35, 46	0
1	D	186/187 (99%)	0.08	1 (0%) 90 92	12, 20, 33, 41	0
All	All	728/748 (97%)	0.06	11 (1%) 73 78	12, 20, 33, 46	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	92	TYR	5.3
1	D	4	GLY	4.9
1	C	4	GLY	4.8
1	C	3	VAL	3.6
1	B	104	TYR	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	B	302	14/15	0.81	0.52	8,12,14,16	14
2	NAG	D	302	14/15	0.82	0.19	13,15,19,20	14
2	NAG	B	303	14/15	0.83	0.64	4,12,18,18	14
2	NAG	B	301	15/15	0.84	0.52	5,12,14,14	15
2	NAG	D	301	15/15	0.86	0.20	8,13,17,18	15
2	NAG	D	201	15/15	0.90	0.26	12,16,19,20	15
2	NAG	D	303	14/15	0.91	0.17	12,16,19,22	14
2	NAG	A	303	14/15	0.91	0.12	16,23,31,35	0
2	NAG	D	202	14/15	0.92	0.23	9,12,18,21	14
2	NAG	A	201	15/15	0.92	0.09	22,25,33,33	0
2	NAG	A	301	15/15	0.94	0.10	13,16,20,22	0
2	NAG	A	202	14/15	0.94	0.09	16,21,28,31	0
2	NAG	A	302	14/15	0.94	0.09	15,18,27,28	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.