



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 8, 2019 – 10:50 PM EDT

PDB ID : 6H2D
Title : Structure of the soluble AhlC of the tripartite alpha-pore forming toxin, AHL, from *Aeromonas hydrophila*.
Authors : Churchill-Angus, A.M.; Wilson, J.S.; Baker, P.J.
Deposited on : 2018-07-13
Resolution : 2.62 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.3.2
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.3.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	3285 (2.64-2.60)
Clashscore	122126	3641 (2.64-2.60)
Ramachandran outliers	120053	3586 (2.64-2.60)
Sidechain outliers	120020	3586 (2.64-2.60)
RSRZ outliers	108989	3218 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	P	274	<div> <div>2%</div> <div>58%</div> <div>20%</div> <div>22%</div> </div>
1	Q	274	<div> <div>54%</div> <div>18%</div> <div>28%</div> </div>
1	R	274	<div> <div>3%</div> <div>49%</div> <div>16%</div> <div>35%</div> </div>
1	S	274	<div> <div>2%</div> <div>65%</div> <div>16%</div> <div>18%</div> </div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 6152 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AhlC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	P	215	Total	C	N	O	S	0	0	0
			1642	1022	292	326	2			
1	Q	197	Total	C	N	O	S	0	0	0
			1483	914	265	302	2			
1	R	177	Total	C	N	O	S	0	0	0
			1336	821	240	273	2			
1	S	224	Total	C	N	O	S	0	0	0
			1691	1051	300	338	2			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
P	267	LEU	-	expression tag	UNP A0A1N6TH80
P	268	GLU	-	expression tag	UNP A0A1N6TH80
P	269	HIS	-	expression tag	UNP A0A1N6TH80
P	270	HIS	-	expression tag	UNP A0A1N6TH80
P	271	HIS	-	expression tag	UNP A0A1N6TH80
P	272	HIS	-	expression tag	UNP A0A1N6TH80
P	273	HIS	-	expression tag	UNP A0A1N6TH80
P	274	HIS	-	expression tag	UNP A0A1N6TH80
Q	267	LEU	-	expression tag	UNP A0A1N6TH80
Q	268	GLU	-	expression tag	UNP A0A1N6TH80
Q	269	HIS	-	expression tag	UNP A0A1N6TH80
Q	270	HIS	-	expression tag	UNP A0A1N6TH80
Q	271	HIS	-	expression tag	UNP A0A1N6TH80
Q	272	HIS	-	expression tag	UNP A0A1N6TH80
Q	273	HIS	-	expression tag	UNP A0A1N6TH80
Q	274	HIS	-	expression tag	UNP A0A1N6TH80
R	267	LEU	-	expression tag	UNP A0A1N6TH80
R	268	GLU	-	expression tag	UNP A0A1N6TH80
R	269	HIS	-	expression tag	UNP A0A1N6TH80
R	270	HIS	-	expression tag	UNP A0A1N6TH80
R	271	HIS	-	expression tag	UNP A0A1N6TH80

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
R	272	HIS	-	expression tag	UNP A0A1N6TH80
R	273	HIS	-	expression tag	UNP A0A1N6TH80
R	274	HIS	-	expression tag	UNP A0A1N6TH80
S	267	LEU	-	expression tag	UNP A0A1N6TH80
S	268	GLU	-	expression tag	UNP A0A1N6TH80
S	269	HIS	-	expression tag	UNP A0A1N6TH80
S	270	HIS	-	expression tag	UNP A0A1N6TH80
S	271	HIS	-	expression tag	UNP A0A1N6TH80
S	272	HIS	-	expression tag	UNP A0A1N6TH80
S	273	HIS	-	expression tag	UNP A0A1N6TH80
S	274	HIS	-	expression tag	UNP A0A1N6TH80


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: AhlC




• Molecule 1: AhlC




• Molecule 1: AhlC





● Molecule 1: AhlC



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.07Å 61.66Å 129.77Å 90.00° 92.01° 90.00°	Depositor
Resolution (Å)	46.74 – 2.62 46.74 – 2.62	Depositor EDS
% Data completeness (in resolution range)	99.6 (46.74-2.62) 89.9 (46.74-2.62)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0230	Depositor
R, R_{free}	0.269 , 0.334 0.265 , 0.318	Depositor DCC
R_{free} test set	1448 reflections (4.65%)	wwPDB-VP
Wilson B-factor (Å ²)	50.0	Xtriage
Anisotropy	0.244	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 21.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.076 for h,-k,-l	Xtriage
Reported twinning fraction	0.888 for H, K, L 0.112 for -h,-k,l	Depositor
Outliers	2 of 31139 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6152	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	P	0.63	0/1650	0.76	0/2232
1	Q	0.60	0/1489	0.72	0/2017
1	R	0.55	0/1341	0.70	0/1815
1	S	0.59	0/1698	0.73	0/2297
All	All	0.60	0/6178	0.73	0/8361

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	S	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	S	122	ARG	Sidechain
1	S	196	ARG	Sidechain
1	S	217	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	1642	0	1687	42	0
1	Q	1483	0	1493	32	0
1	R	1336	0	1350	25	0
1	S	1691	0	1729	36	0
All	All	6152	0	6259	125	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 125 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:P:28:GLN:NE2	1:P:32:ASN:OD1	2.18	0.77
1:R:23:PHE:HB3	1:R:26:LEU:HD11	1.68	0.75
1:R:21:GLN:NE2	1:R:34:GLN:O	2.22	0.71
1:S:248:ARG:O	1:S:252:THR:HG23	1.93	0.68
1:S:256:THR:O	1:S:260:THR:HG23	1.95	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	P	207/274 (76%)	194 (94%)	12 (6%)	1 (0%)	31	54
1	Q	189/274 (69%)	176 (93%)	12 (6%)	1 (0%)	31	54
1	R	171/274 (62%)	163 (95%)	7 (4%)	1 (1%)	27	49
1	S	216/274 (79%)	201 (93%)	15 (7%)	0	100	100
All	All	783/1096 (71%)	734 (94%)	46 (6%)	3 (0%)	36	58

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	P	90	GLN
1	Q	134	LEU
1	R	27	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	P	181/223 (81%)	177 (98%)	4 (2%)	55	78
1	Q	162/223 (73%)	156 (96%)	6 (4%)	37	63
1	R	147/223 (66%)	143 (97%)	4 (3%)	48	73
1	S	183/223 (82%)	180 (98%)	3 (2%)	65	83
All	All	673/892 (75%)	656 (98%)	17 (2%)	50	75

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Q	174	THR
1	Q	211	PHE
1	R	216	ASP
1	Q	136	SER
1	S	147	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	R	8	GLN
1	R	29	ASN
1	S	188	ASN
1	Q	89	GLN
1	S	135	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	P	215/274 (78%)	-0.19	4 (1%) 66 62	28, 49, 82, 165	0
1	Q	197/274 (71%)	-0.07	2 (1%) 82 79	31, 60, 88, 101	0
1	R	177/274 (64%)	-0.03	9 (5%) 28 22	35, 59, 89, 104	0
1	S	224/274 (81%)	-0.20	5 (2%) 62 56	30, 52, 95, 131	0
All	All	813/1096 (74%)	-0.13	20 (2%) 57 51	28, 55, 90, 165	0

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	R	176	THR	5.0
1	P	250	PHE	4.5
1	S	232	ILE	3.8
1	Q	172	THR	3.6
1	S	165	GLY	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.