



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 09:21 am GMT

PDB ID : 1H3Y  
Title : Crystal structure of a human IgG1 Fc-fragment,high salt condition  
Authors : Krapp, S.; Mimura, Y.; Jefferis, R.; Huber, R.; Sondermann, P.  
Deposited on : 2002-09-19  
Resolution : 4.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1039 (4.54-3.66)
Clashscore	122126	1052 (4.50-3.70)
Ramachandran outliers	120053	1006 (4.50-3.70)
Sidechain outliers	120020	1005 (4.52-3.68)
RSRZ outliers	108989	1161 (4.60-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	
1	B	223	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	1449	X	-	-	-
2	NAG	A	1452	-	-	X	-
3	FUL	A	1445	-	-	X	-
4	MAN	A	1451	-	-	X	-

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 3015 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IG GAMMA-1 CHAIN C REGION.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	0	0	1
			1373	848	245	274	6			
1	B	206	Total	C	N	O	S	0	0	0
			1422	886	246	283	7			

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is BETA-L-FUCOSE (three-letter code: FUL) (formula:  $C_6H_{12}O_5$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	6	4		
3	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 4 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	B	1	Total	C	O	0	0
			11	6	5		
4	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is ALPHA D-GALACTOSE (three-letter code: GLA) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is BETA-D-MANNOSE (three-letter code: BMA) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			11	6	5		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: IG GAMMA-1 CHAIN C REGION

Chain A: 



#### • Molecule 1: IG GAMMA-1 CHAIN C REGION

Chain B: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	152.94Å 152.94Å 116.96Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 4.10 38.24 – 4.13	Depositor EDS
% Data completeness (in resolution range)	80.0 (50.00-4.10) 79.2 (38.24-4.13)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.58 (at 4.13Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.285 , 0.349 0.268 , 0.323	Depositor DCC
$R_{free}$ test set	285 reflections (5.46%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	59.4	Xtriage
Anisotropy	0.349	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 148.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.81	EDS
Total number of atoms	3015	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLA, FUL, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.51	0/1415	0.83	1/1927 (0.1%)
1	B	0.53	0/1466	0.87	4/1988 (0.2%)
All	All	0.52	0/2881	0.85	5/3915 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	402	GLY	N-CA-C	-6.03	98.02	113.10
1	A	402	GLY	N-CA-C	-5.78	98.65	113.10
1	B	443	LEU	CA-CB-CG	-5.73	102.13	115.30
1	B	297	ASN	N-CA-CB	-5.18	101.27	110.60
1	B	395	PRO	N-CA-C	-5.11	98.81	112.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1373	0	997	88	0
1	B	1422	0	1100	114	1
2	A	56	0	48	18	0
2	B	56	0	48	8	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	10	0	10	7	0
3	B	10	0	10	3	0
4	A	33	0	26	8	0
4	B	22	0	18	4	0
5	A	11	0	10	3	0
5	B	11	0	10	2	0
6	B	11	0	8	1	0
All	All	3015	0	2285	221	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 42.

The worst 5 of 221 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1452:NAG:H83	3:B:1445:FUL:H4	1.25	1.18
1:A:346:PRO:HB3	1:A:372:PHE:HB3	1.24	1.10
1:B:325:ASN:ND2	1:B:326:LYS:H	1.52	1.06
1:A:283:GLN:HG2	1:A:285:HIS:H	1.30	0.96
4:B:1451:MAN:H2	2:B:1452:NAG:N2	1.78	0.96

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:309:LEU:CD1	1:B:309:LEU:CD1[9_765]	1.59	0.61

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/223 (92%)	165 (80%)	22 (11%)	18 (9%)	1	14
1	B	204/223 (92%)	159 (78%)	29 (14%)	16 (8%)	1	17
All	All	409/446 (92%)	324 (79%)	51 (12%)	34 (8%)	1	15

5 of 34 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	339	ALA
1	A	377	ILE
1	A	443	LEU
1	B	297	ASN
1	B	377	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/206 (52%)	101 (94%)	7 (6%)	19	51
1	B	126/206 (61%)	115 (91%)	11 (9%)	11	40
All	All	234/412 (57%)	216 (92%)	18 (8%)	14	46

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	271	PRO
1	B	325	ASN
1	B	411	THR
1	A	443	LEU
1	B	260	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	418	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	434	ASN
1	B	325	ASN
1	A	325	ASN
1	B	312	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	1444	1,3,2	14,14,15	1.14	2 (14%)	17,19,21	1.53	3 (17%)
3	FUL	A	1445	2	9,10,11	0.64	0	13,14,16	0.68	0
2	NAG	A	1446	2,4	14,14,15	0.99	1 (7%)	17,19,21	1.51	3 (17%)
4	MAN	A	1447	2,4	11,11,12	0.70	0	15,15,17	1.72	2 (13%)
4	MAN	A	1448	2,4	11,11,12	1.01	1 (9%)	15,15,17	0.82	0
2	NAG	A	1449	5,4	14,14,15	1.32	3 (21%)	17,19,21	2.11	5 (29%)
5	GLA	A	1450	2	11,11,12	1.36	1 (9%)	15,15,17	0.99	0
4	MAN	A	1451	2,4	11,11,12	0.90	0	15,15,17	1.32	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	1452	4	14,14,15	1.26	2 (14%)	17,19,21	1.16	2 (11%)
2	NAG	B	1444	1,3,2	14,14,15	0.76	0	17,19,21	1.61	3 (17%)
3	FUL	B	1445	2	9,10,11	0.93	0	13,14,16	0.82	0
2	NAG	B	1446	2,6	14,14,15	0.75	0	17,19,21	1.28	2 (11%)
6	BMA	B	1447	2,4	11,11,12	1.15	1 (9%)	15,15,17	0.77	0
4	MAN	B	1448	2,6	11,11,12	1.05	1 (9%)	15,15,17	1.40	2 (13%)
2	NAG	B	1449	5,4	14,14,15	0.96	0	17,19,21	1.77	4 (23%)
5	GLA	B	1450	2	11,11,12	0.82	0	15,15,17	1.11	2 (13%)
4	MAN	B	1451	2,6	11,11,12	1.12	1 (9%)	15,15,17	2.28	3 (20%)
2	NAG	B	1452	4	14,14,15	0.88	0	17,19,21	1.12	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1444	1,3,2	-	1/6/23/26	0/1/1/1
3	FUL	A	1445	2	-	0/0/17/20	0/1/1/1
2	NAG	A	1446	2,4	-	0/6/23/26	0/1/1/1
4	MAN	A	1447	2,4	-	0/2/19/22	0/1/1/1
4	MAN	A	1448	2,4	-	0/2/19/22	0/1/1/1
2	NAG	A	1449	5,4	1/1/5/7	0/6/23/26	0/1/1/1
5	GLA	A	1450	2	-	0/2/19/22	0/1/1/1
4	MAN	A	1451	2,4	-	0/2/19/22	0/1/1/1
2	NAG	A	1452	4	-	0/6/23/26	0/1/1/1
2	NAG	B	1444	1,3,2	-	0/6/23/26	0/1/1/1
3	FUL	B	1445	2	-	0/0/17/20	0/1/1/1
2	NAG	B	1446	2,6	-	0/6/23/26	0/1/1/1
6	BMA	B	1447	2,4	-	0/2/19/22	0/1/1/1
4	MAN	B	1448	2,6	-	0/2/19/22	0/1/1/1
2	NAG	B	1449	5,4	-	0/6/23/26	0/1/1/1
5	GLA	B	1450	2	-	0/2/19/22	0/1/1/1
4	MAN	B	1451	2,6	-	0/2/19/22	0/1/1/1
2	NAG	B	1452	4	-	0/6/23/26	0/1/1/1

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1444	NAG	C1-C2	-2.25	1.49	1.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1448	MAN	O2-C2	2.03	1.47	1.43
2	A	1446	NAG	O4-C4	2.11	1.48	1.43
2	A	1452	NAG	O5-C1	2.18	1.47	1.43
2	A	1449	NAG	O5-C5	2.26	1.48	1.43

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1449	NAG	C4-C3-C2	-5.56	102.86	111.02
2	B	1444	NAG	C2-N2-C7	-3.89	117.27	122.94
2	B	1449	NAG	C4-C3-C2	-3.74	105.53	111.02
2	B	1444	NAG	C4-C3-C2	-3.74	105.53	111.02
4	A	1451	MAN	C1-C2-C3	-3.71	104.97	109.66

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	1449	NAG	C1

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1444	NAG	O7-C7-N2-C2

There are no ring outliers.

14 monomers are involved in 32 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1445	FUL	7	0
2	A	1446	NAG	6	0
4	A	1448	MAN	2	0
2	A	1449	NAG	3	0
5	A	1450	GLA	3	0
4	A	1451	MAN	6	0
2	A	1452	NAG	9	0
3	B	1445	FUL	3	0
2	B	1446	NAG	2	0
6	B	1447	BMA	1	0
2	B	1449	NAG	2	0
5	B	1450	GLA	2	0
4	B	1451	MAN	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1452	NAG	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	207/223 (92%)	-0.52	0 100 100	28, 32, 47, 47	0
1	B	206/223 (92%)	-0.76	0 100 100	14, 17, 22, 23	0
All	All	413/446 (92%)	-0.64	0 100 100	14, 23, 46, 47	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	MAN	A	1447	11/12	0.83	0.19	71,73,76,76	0
4	MAN	A	1451	11/12	0.85	0.23	73,75,76,76	0
2	NAG	A	1444	14/15	0.87	0.25	71,73,76,76	0
4	MAN	A	1448	11/12	0.88	0.23	71,73,76,76	0
2	NAG	A	1452	14/15	0.88	0.26	71,73,75,76	0
2	NAG	A	1449	14/15	0.89	0.25	71,74,76,76	0
4	MAN	B	1451	11/12	0.89	0.26	33,35,38,38	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GLA	B	1450	11/12	0.89	0.23	33,34,38,39	0
2	NAG	A	1446	14/15	0.89	0.29	71,74,75,76	0
4	MAN	B	1448	11/12	0.89	0.23	33,35,38,38	0
2	NAG	B	1452	14/15	0.90	0.21	33,36,39,39	0
2	NAG	B	1449	14/15	0.91	0.18	33,35,38,38	0
2	NAG	B	1444	14/15	0.91	0.19	33,35,39,39	0
3	FUL	A	1445	10/11	0.91	0.20	70,73,75,76	0
5	GLA	A	1450	11/12	0.91	0.26	71,72,76,76	0
3	FUL	B	1445	10/11	0.95	0.19	33,36,38,39	0
6	BMA	B	1447	11/12	0.95	0.12	33,36,38,39	0
2	NAG	B	1446	14/15	0.96	0.16	33,35,38,39	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.