



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 11, 2018 – 09:45 PM EST

PDB ID : 3HM7
Title : Crystal structure of allantoinase from *Bacillus halodurans* C-125
Authors : Patskovsky, Y.; Romero, R.; Rutter, M.; Miller, S.; Wasserman, S.R.; Sauder, J.M.; Raushel, F.M.; Burley, S.K.; Almo, S.C.; New York Structural GenomiX Research Consortium (NYSGXRC); New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2009-05-28
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	rb-20031633
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20031633

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	111664	2767 (2.60-2.60)
Clashscore	122126	3110 (2.60-2.60)
Ramachandran outliers	120053	3062 (2.60-2.60)
Sidechain outliers	120020	3062 (2.60-2.60)
RSRZ outliers	108989	2706 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	448	<div><div>7%</div><div>82%</div><div>14%</div><div>• •</div></div>
1	B	448	<div><div>5%</div><div>82%</div><div>15%</div><div>• •</div></div>
1	C	448	<div><div>27%</div><div>78%</div><div>18%</div><div>• •</div></div>
1	D	448	<div><div>7%</div><div>81%</div><div>15%</div><div>• •</div></div>
1	E	448	<div><div>3%</div><div>87%</div><div>10%</div><div>• •</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	448	<div><div></div><div>2%</div><div>84%</div><div>13%</div><div>••</div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 20901 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Allantoinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	436	Total	C	N	O	S	0	4	0
			3425	2166	593	649	17			
1	B	437	Total	C	N	O	S	0	5	0
			3437	2173	594	653	17			
1	C	435	Total	C	N	O	S	0	3	0
			3410	2154	588	651	17			
1	D	436	Total	C	N	O	S	0	2	0
			3415	2157	591	650	17			
1	E	436	Total	C	N	O	S	0	2	0
			3412	2156	590	649	17			
1	F	436	Total	C	N	O	S	0	4	0
			3425	2165	592	651	17			

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
A	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
A	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
A	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
A	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
A	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
B	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
B	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
B	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
B	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
B	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
C	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
C	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
C	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
C	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
C	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
D	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
D	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
D	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
D	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
D	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
E	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
E	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
E	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
E	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
E	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
F	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
F	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
F	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
F	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total 1	Zn 1	0	0
2	E	1	Total 1	Zn 1	0	0
2	B	1	Total 1	Zn 1	0	0
2	C	1	Total 1	Zn 1	0	0
2	A	1	Total 1	Zn 1	0	0
2	F	1	Total 1	Zn 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	53	Total 53	O 53	0	0
3	B	69	Total 69	O 69	0	0
3	C	39	Total 39	O 39	0	0
3	D	47	Total 47	O 47	0	0
3	E	74	Total 74	O 74	0	0
3	F	89	Total 89	O 89	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Allantoinase




• Molecule 1: Allantoinase




• Molecule 1: Allantoinase




- Molecule 1: Allantoinase



- Molecule 1: Allantoinase



- Molecule 1: Allantoinase



H433	GLU
Y434	GLY
Y435	HIS
Q436	HIS
D437	HIS
H438	HIS
	HIS

4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	157.66Å 157.66Å 418.03Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.60 27.39 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-2.60) 100.0 (27.39-2.60)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.94 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.3.0034	Depositor
R, R_{free}	0.240 , 0.267 0.240 , 0.266	Depositor DCC
R_{free} test set	2853 reflections (3.01%)	wwPDB-VP
Wilson B-factor (Å ²)	41.8	Xtriage
Anisotropy	0.393	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 50.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	20901	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 69.84 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.4571e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/3507	0.60	0/4749
1	B	0.44	0/3523	0.61	1/4770 (0.0%)
1	C	0.43	0/3485	0.62	0/4720
1	D	0.42	0/3488	0.59	0/4724
1	E	0.43	0/3488	0.59	0/4724
1	F	0.41	0/3508	0.61	1/4752 (0.0%)
All	All	0.42	0/20999	0.60	2/28439 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	395	LEU	CA-CB-CG	6.19	129.54	115.30
1	F	262	VAL	CB-CA-C	-5.13	101.65	111.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	159	ASP	Peptide
1	A	160	ASP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3425	0	3404	27	0
1	B	3437	0	3411	25	0
1	C	3410	0	3376	32	0
1	D	3415	0	3379	29	0
1	E	3412	0	3380	16	0
1	F	3425	0	3393	19	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	53	0	0	0	0
3	B	69	0	0	0	0
3	C	39	0	0	0	0
3	D	47	0	0	0	0
3	E	74	0	0	0	0
3	F	89	0	0	0	0
All	All	20901	0	20343	138	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 138 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:67:ASN:ND2	1:D:98:ASN:HB2	1.96	0.80
1:F:75:GLU:HG2	1:F:318:PRO:HG3	1.69	0.73
1:B:75:GLU:HG2	1:B:318:PRO:HG3	1.76	0.68
1:E:67:ASN:ND2	1:E:98:ASN:HB2	2.09	0.67
1:F:67:ASN:ND2	1:F:98:ASN:HB2	2.09	0.67

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/448 (98%)	410 (94%)	25 (6%)	3 (1%)	24	46
1	B	440/448 (98%)	425 (97%)	13 (3%)	2 (0%)	31	56
1	C	436/448 (97%)	397 (91%)	35 (8%)	4 (1%)	19	38
1	D	436/448 (97%)	416 (95%)	18 (4%)	2 (0%)	31	56
1	E	436/448 (97%)	419 (96%)	15 (3%)	2 (0%)	31	56
1	F	438/448 (98%)	424 (97%)	12 (3%)	2 (0%)	31	56
All	All	2624/2688 (98%)	2491 (95%)	118 (4%)	15 (1%)	27	51

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	132	PRO
1	A	242	HIS
1	B	242	HIS
1	C	242	HIS
1	D	242	HIS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	375/382 (98%)	343 (92%)	32 (8%)	12	23
1	B	377/382 (99%)	351 (93%)	26 (7%)	17	34
1	C	373/382 (98%)	336 (90%)	37 (10%)	8	16

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	373/382 (98%)	340 (91%)	33 (9%)	11	21
1	E	373/382 (98%)	350 (94%)	23 (6%)	20	40
1	F	375/382 (98%)	347 (92%)	28 (8%)	15	29
All	All	2246/2292 (98%)	2067 (92%)	179 (8%)	13	26

5 of 179 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	304	LEU
1	D	113	ARG
1	F	242	HIS
1	C	328	THR
1	C	393	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	418	GLN
1	D	352	HIS
1	F	383	GLN
1	D	162	GLN
1	B	67	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	436/448 (97%)	0.30	32 (7%) 15 10	22, 38, 54, 77	0
1	B	437/448 (97%)	0.09	21 (4%) 30 23	28, 37, 52, 66	0
1	C	435/448 (97%)	1.48	122 (28%) 0 0	22, 39, 51, 74	0
1	D	436/448 (97%)	0.22	32 (7%) 15 10	26, 38, 53, 70	0
1	E	436/448 (97%)	-0.02	12 (2%) 53 46	25, 37, 47, 68	0
1	F	436/448 (97%)	-0.06	8 (1%) 68 63	26, 37, 48, 64	0
All	All	2616/2688 (97%)	0.33	227 (8%) 10 7	22, 37, 52, 77	0

The worst 5 of 227 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	158	THR	9.2
1	C	402	ASN	7.6
1	A	157	GLY	7.3
1	C	156	CYS	6.9
1	C	113	ARG	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	D	447	1/1	0.84	0.20	40,40,40,40	1
2	ZN	C	447	1/1	0.93	0.20	50,50,50,50	1
2	ZN	F	447	1/1	0.94	0.22	39,39,39,39	1
2	ZN	B	447	1/1	0.96	0.08	39,39,39,39	1
2	ZN	E	447	1/1	0.96	0.08	39,39,39,39	1
2	ZN	A	447	1/1	0.97	0.06	43,43,43,43	1

6.5 Other polymers [i](#)

There are no such residues in this entry.