



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 2, 2019 – 04:21 PM EST

PDB ID : 6HWK
Title : Glucosamine kinase (crystal form B)
Authors : Manso, J.A.; Pereira, P.J.B.
Deposited on : 2018-10-12
Resolution : 2.69 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.4
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	3333 (2.70-2.66)
Clashscore	122126	3672 (2.70-2.66)
Ramachandran outliers	120053	3620 (2.70-2.66)
Sidechain outliers	120020	3620 (2.70-2.66)
RSRZ outliers	108989	3248 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	<div> <div>3%</div> <div>87%</div> <div>9%</div> </div>
1	B	451	<div> <div>4%</div> <div>85%</div> <div>11%</div> </div>
1	C	451	<div> <div>3%</div> <div>83%</div> <div>7%</div> <div>10%</div> </div>
1	D	451	<div> <div>%</div> <div>85%</div> <div>7%</div> <div>8%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12376 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucosamine kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	412	Total	C	N	O	S	0	0	0
			3026	1922	540	558	6			
1	B	403	Total	C	N	O	S	0	0	0
			2958	1881	523	548	6			
1	C	408	Total	C	N	O	S	0	0	0
			3020	1920	543	551	6			
1	D	416	Total	C	N	O	S	0	0	0
			3109	1970	563	570	6			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	439	LYS	-	expression tag	UNP A0A1H7TQR5
A	440	LEU	-	expression tag	UNP A0A1H7TQR5
A	441	ALA	-	expression tag	UNP A0A1H7TQR5
A	442	ALA	-	expression tag	UNP A0A1H7TQR5
A	443	ALA	-	expression tag	UNP A0A1H7TQR5
A	444	LEU	-	expression tag	UNP A0A1H7TQR5
A	445	GLU	-	expression tag	UNP A0A1H7TQR5
A	446	HIS	-	expression tag	UNP A0A1H7TQR5
A	447	HIS	-	expression tag	UNP A0A1H7TQR5
A	448	HIS	-	expression tag	UNP A0A1H7TQR5
A	449	HIS	-	expression tag	UNP A0A1H7TQR5
A	450	HIS	-	expression tag	UNP A0A1H7TQR5
A	451	HIS	-	expression tag	UNP A0A1H7TQR5
B	439	LYS	-	expression tag	UNP A0A1H7TQR5
B	440	LEU	-	expression tag	UNP A0A1H7TQR5
B	441	ALA	-	expression tag	UNP A0A1H7TQR5
B	442	ALA	-	expression tag	UNP A0A1H7TQR5
B	443	ALA	-	expression tag	UNP A0A1H7TQR5
B	444	LEU	-	expression tag	UNP A0A1H7TQR5
B	445	GLU	-	expression tag	UNP A0A1H7TQR5
B	446	HIS	-	expression tag	UNP A0A1H7TQR5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	447	HIS	-	expression tag	UNP A0A1H7TQR5
B	448	HIS	-	expression tag	UNP A0A1H7TQR5
B	449	HIS	-	expression tag	UNP A0A1H7TQR5
B	450	HIS	-	expression tag	UNP A0A1H7TQR5
B	451	HIS	-	expression tag	UNP A0A1H7TQR5
C	439	LYS	-	expression tag	UNP A0A1H7TQR5
C	440	LEU	-	expression tag	UNP A0A1H7TQR5
C	441	ALA	-	expression tag	UNP A0A1H7TQR5
C	442	ALA	-	expression tag	UNP A0A1H7TQR5
C	443	ALA	-	expression tag	UNP A0A1H7TQR5
C	444	LEU	-	expression tag	UNP A0A1H7TQR5
C	445	GLU	-	expression tag	UNP A0A1H7TQR5
C	446	HIS	-	expression tag	UNP A0A1H7TQR5
C	447	HIS	-	expression tag	UNP A0A1H7TQR5
C	448	HIS	-	expression tag	UNP A0A1H7TQR5
C	449	HIS	-	expression tag	UNP A0A1H7TQR5
C	450	HIS	-	expression tag	UNP A0A1H7TQR5
C	451	HIS	-	expression tag	UNP A0A1H7TQR5
D	439	LYS	-	expression tag	UNP A0A1H7TQR5
D	440	LEU	-	expression tag	UNP A0A1H7TQR5
D	441	ALA	-	expression tag	UNP A0A1H7TQR5
D	442	ALA	-	expression tag	UNP A0A1H7TQR5
D	443	ALA	-	expression tag	UNP A0A1H7TQR5
D	444	LEU	-	expression tag	UNP A0A1H7TQR5
D	445	GLU	-	expression tag	UNP A0A1H7TQR5
D	446	HIS	-	expression tag	UNP A0A1H7TQR5
D	447	HIS	-	expression tag	UNP A0A1H7TQR5
D	448	HIS	-	expression tag	UNP A0A1H7TQR5
D	449	HIS	-	expression tag	UNP A0A1H7TQR5
D	450	HIS	-	expression tag	UNP A0A1H7TQR5
D	451	HIS	-	expression tag	UNP A0A1H7TQR5

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Cl 1 1	0	0
2	A	1	Total Cl 1 1	0	0
2	D	1	Total Cl 1 1	0	0
2	C	1	Total Cl 1 1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Mg 1	0	0
3	D	1	Total 1	Mg 1	0	0
3	C	1	Total 1	Mg 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	60	Total 60	O 60	0	0
4	B	54	Total 54	O 54	0	0
4	C	57	Total 57	O 57	0	0
4	D	85	Total 85	O 85	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glucosamine kinase




- Molecule 1: Glucosamine kinase




- Molecule 1: Glucosamine kinase



- Molecule 1: Glucosamine kinase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.20Å 111.22Å 150.59Å 90.00° 93.21° 90.00°	Depositor
Resolution (Å)	47.27 – 2.69 47.27 – 2.69	Depositor EDS
% Data completeness (in resolution range)	98.9 (47.27-2.69) 98.9 (47.27-2.69)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 2.69Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.205 , 0.242 0.206 , 0.241	Depositor DCC
R_{free} test set	2647 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	41.8	Xtriage
Anisotropy	0.319	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 59.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12376	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/3095	0.40	0/4248
1	B	0.23	0/3026	0.40	0/4154
1	C	0.23	0/3087	0.40	0/4236
1	D	0.23	0/3179	0.40	0/4353
All	All	0.23	0/12387	0.40	0/16991

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3026	0	2941	11	0
1	B	2958	0	2868	12	0
1	C	3020	0	2968	17	0
1	D	3109	0	3085	17	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	60	0	0	0	0
4	B	54	0	0	0	0
4	C	57	0	0	0	0
4	D	85	0	0	0	0
All	All	12376	0	11862	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 57 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:PRO:HB2	1:A:73:SER:HB3	1.75	0.68
1:A:149:ALA:HB2	1:A:180:LEU:HD13	1.79	0.64
1:D:140:ALA:HB3	1:D:143:GLU:HG3	1.79	0.63
1:B:61:PRO:HB2	1:B:73:SER:HB2	1.82	0.61
1:C:149:ALA:HB2	1:C:180:LEU:HD13	1.84	0.60

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/451 (89%)	385 (96%)	17 (4%)	0	100	100
1	B	393/451 (87%)	381 (97%)	12 (3%)	0	100	100
1	C	396/451 (88%)	381 (96%)	15 (4%)	0	100	100
1	D	408/451 (90%)	396 (97%)	11 (3%)	1 (0%)	49	75
All	All	1599/1804 (89%)	1543 (96%)	55 (3%)	1 (0%)	53	79

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	19	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/335 (86%)	287 (100%)	0	100	100
1	B	280/335 (84%)	279 (100%)	1 (0%)	92	97
1	C	291/335 (87%)	291 (100%)	0	100	100
1	D	303/335 (90%)	303 (100%)	0	100	100
All	All	1161/1340 (87%)	1160 (100%)	1 (0%)	94	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	70	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	412/451 (91%)	0.35	13 (3%) 47 46	23, 41, 73, 87	0
1	B	403/451 (89%)	0.42	20 (4%) 29 27	24, 46, 86, 113	0
1	C	408/451 (90%)	0.32	12 (2%) 51 50	27, 44, 72, 90	0
1	D	416/451 (92%)	0.23	5 (1%) 79 79	24, 39, 67, 105	0
All	All	1639/1804 (90%)	0.33	50 (3%) 49 48	23, 42, 77, 113	0

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	67	ALA	5.1
1	B	9	VAL	4.8
1	C	70	VAL	4.8
1	B	78	GLY	4.0
1	B	75	PRO	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MG	D	502	1/1	0.86	0.08	44,44,44,44	0
3	MG	C	502	1/1	0.90	0.08	46,46,46,46	0
2	CL	B	501	1/1	0.97	0.14	26,26,26,26	0
2	CL	A	501	1/1	0.98	0.12	34,34,34,34	0
3	MG	A	502	1/1	0.99	0.13	38,38,38,38	0
2	CL	C	501	1/1	0.99	0.11	33,33,33,33	0
2	CL	D	501	1/1	0.99	0.12	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.