



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 09:43 pm GMT

PDB ID : 2I2O
Title : Crystal Structure of an eIF4G-like Protein from *Danio rerio*
Authors : Bitto, E.; Wesenberg, G.E.; Phillips Jr., G.N.; McCoy, J.G.; Bingman, C.A.;
Center for Eukaryotic Structural Genomics (CESG)
Deposited on : 2006-08-16
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	6904 (1.94-1.90)
Clashscore	122126	7577 (1.94-1.90)
Ramachandran outliers	120053	7491 (1.94-1.90)
Sidechain outliers	120020	7491 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	224	
1	B	224	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called eIF4G-like protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	211	Total	C	N	O	S	Se	0	2	0
			1739	1106	295	327	5	6			
1	B	210	Total	C	N	O	S	Se	0	3	0
			1737	1106	295	325	5	6			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ALA	-	CLONING ARTIFACT	UNP Q5EAQ1
A	0	ILE	-	CLONING ARTIFACT	UNP Q5EAQ1
A	1	ALA	-	CLONING ARTIFACT	UNP Q5EAQ1
B	-1	ALA	-	CLONING ARTIFACT	UNP Q5EAQ1
B	0	ILE	-	CLONING ARTIFACT	UNP Q5EAQ1
B	1	ALA	-	CLONING ARTIFACT	UNP Q5EAQ1

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ni	0	0
			1	1		
2	A	1	Total	Ni	0	0
			1	1		


- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	202	Total	O	0	0
			202	202		
3	B	216	Total	O	0	0
			216	216		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: eIF4G-like protein

Chain A: 



- Molecule 1: eIF4G-like protein

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	317.80Å 40.95Å 40.92Å 90.00° 90.26° 90.00°	Depositor
Resolution (Å)	40.61 – 1.92 40.61 – 1.92	Depositor EDS
% Data completeness (in resolution range)	97.4 (40.61-1.92) 97.4 (40.61-1.92)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.52 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.190 , 0.235 0.252 , 0.282	Depositor DCC
R_{free} test set	1998 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	21.0	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 35.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.032 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3896	wwPDB-VP
Average B, all atoms (Å ²)	8.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/1763	0.73	3/2364 (0.1%)
1	B	0.69	0/1761	0.68	0/2362
All	All	0.67	0/3524	0.71	3/4726 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	192	ARG	NE-CZ-NH2	-9.16	115.72	120.30
1	A	192	ARG	NE-CZ-NH1	8.22	124.41	120.30
1	A	192	ARG	CD-NE-CZ	5.06	130.68	123.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	118	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1739	0	1753	32	0
1	B	1737	0	1755	29	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	202	0	0	4	0
3	B	216	0	0	3	0
All	All	3896	0	3508	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181[B]:PHE:CZ	1:B:181[B]:PHE:HE2	1.70	1.08
1:B:117:VAL:O	1:B:120:MSE:HB3	1.63	0.98
1:B:26:LEU:HD21	1:B:58:ILE:HG13	1.47	0.96
1:A:181[B]:PHE:CZ	1:B:181[B]:PHE:CE2	2.61	0.88
1:A:208:LEU:HD22	3:A:650:HOH:O	1.80	0.81
1:A:117:VAL:O	1:A:120:MSE:HB3	1.82	0.79
1:B:99:GLN:NE2	1:B:146:GLU:OE2	2.19	0.75
1:A:181[B]:PHE:CE1	1:B:181[B]:PHE:HE2	2.09	0.71
1:B:119:ASN:O	1:B:121:PRO:HD3	1.95	0.67
1:B:43:ASP:OD1	1:B:82:ARG:NH2	2.22	0.66
1:A:181[B]:PHE:CE1	1:B:181[B]:PHE:CE2	2.82	0.66
1:A:167:VAL:HB	3:A:531:HOH:O	1.96	0.66
1:A:91:GLU:HG2	3:A:540:HOH:O	1.96	0.65
1:A:99:GLN:NE2	1:A:146:GLU:OE2	2.31	0.64
1:A:119:ASN:O	1:A:121:PRO:HD3	2.00	0.62
1:B:111:ILE:HG23	1:B:115:LEU:HD22	1.82	0.60
1:A:118:ASN:CG	1:A:119:ASN:H	2.04	0.60
1:B:140:ASP:HA	1:B:143:LYS:HD3	1.84	0.59
1:A:123:VAL:HG13	3:A:534:HOH:O	2.02	0.59
1:A:117:VAL:HB	1:A:120:MSE:HE2	1.86	0.57
1:A:42:VAL:HG11	1:A:82:ARG:HG3	1.89	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:181[A]:PHE:HE1	1:B:196:LEU:CA	2.22	0.53
1:A:126:VAL:HG21	1:A:165[B]:MSE:SE	2.60	0.51
1:A:181[B]:PHE:HD2	1:A:199:LEU:HD12	1.75	0.51
1:B:214:LYS:HG3	3:B:658:HOH:O	2.11	0.51
1:B:126:VAL:HG21	1:B:165[A]:MSE:HG2	1.93	0.50
1:A:26:LEU:HD21	1:A:58:ILE:HG13	1.93	0.49
1:A:116:LYS:HE2	1:A:121:PRO:HG3	1.95	0.49
1:A:117:VAL:O	1:A:118:ASN:HB3	2.12	0.49
1:A:181[A]:PHE:HE1	1:B:181[A]:PHE:CZ	2.31	0.48
1:A:117:VAL:HB	1:A:120:MSE:CE	2.44	0.48
1:A:23:LYS:HA	1:A:26:LEU:HD12	1.94	0.48
1:B:118:ASN:HB3	1:B:120:MSE:H	1.77	0.48
1:A:126:VAL:CG2	1:A:165[B]:MSE:HG2	2.44	0.47
1:B:126:VAL:CG1	1:B:165[A]:MSE:HG2	2.44	0.47
1:B:126:VAL:HG11	1:B:165[A]:MSE:HG2	1.96	0.47
1:B:126:VAL:CG2	1:B:165[A]:MSE:HG2	2.44	0.47
1:B:42:VAL:HG11	1:B:82:ARG:HG3	1.98	0.46
1:A:126:VAL:HG21	1:A:165[B]:MSE:HG2	1.98	0.46
1:B:185:GLU:CD	3:B:614:HOH:O	2.54	0.46
1:A:181[B]:PHE:CE1	1:A:192:ARG:HG3	2.51	0.45
1:A:73:SER:OG	1:A:77:ARG:NH1	2.49	0.45
1:B:126:VAL:HG21	1:B:165[A]:MSE:SE	2.67	0.44
1:A:118:ASN:CG	1:A:119:ASN:N	2.71	0.43
1:A:181[B]:PHE:HE2	1:A:196:LEU:CA	2.33	0.42
1:B:117:VAL:CG2	1:B:120:MSE:HE3	2.50	0.41
1:B:161:GLN:O	1:B:165[B]:MSE:HG2	2.20	0.41
1:A:60:TYR:O	1:A:64:GLN:HG2	2.20	0.41
1:B:165[B]:MSE:HE3	3:B:521:HOH:O	2.19	0.41
1:B:181[A]:PHE:HD1	1:B:199:LEU:HD12	1.85	0.41
1:A:181[B]:PHE:CE2	1:A:196:LEU:HB2	2.55	0.41
1:A:69:GLN:HG2	1:A:70:THR:HG23	2.04	0.40
1:B:116:LYS:HE2	1:B:121:PRO:HG3	2.04	0.40
1:B:187:LEU:HB2	1:B:192:ARG:NH1	2.36	0.40
1:B:47:LYS:HA	1:B:47:LYS:HD3	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/224 (94%)	204 (97%)	6 (3%)	1 (0%)	31	19
1	B	211/224 (94%)	206 (98%)	4 (2%)	1 (0%)	31	19
All	All	422/448 (94%)	410 (97%)	10 (2%)	2 (0%)	31	19

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	118	ASN
1	B	118	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/199 (98%)	192 (98%)	3 (2%)	67	64
1	B	195/199 (98%)	191 (98%)	4 (2%)	56	50
All	All	390/398 (98%)	383 (98%)	7 (2%)	65	56

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	ASP
1	A	27	LYS
1	A	192	ARG
1	B	117	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	143	LYS
1	B	181[A]	PHE
1	B	181[B]	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.