



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 16, 2019 – 11:16 PM EST

PDB ID : 6IB5
Title : Mutant of flavin-dependent tryptophan halogenase Thal with altered regioselectivity (Thal-RebH5)
Authors : Moritzer, A.; Prior, T.; Niemann, H.H.
Deposited on : 2018-11-29
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : rb-20031633
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20031633


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	5449 (2.14-2.10)
Clashscore	122126	5972 (2.14-2.10)
Ramachandran outliers	120053	5914 (2.14-2.10)
Sidechain outliers	120020	5915 (2.14-2.10)
RSRZ outliers	108989	5321 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	<div> <div>2%</div> <div>91%</div> <div>8%</div> <div>.</div> </div>
1	B	534	<div> <div>6%</div> <div>89%</div> <div>9%</div> <div>.</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8962 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tryptophan 6-halogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	526	Total	C	N	O	S	0	5	0
			4240	2710	727	781	22			
1	B	524	Total	C	N	O	S	0	4	0
			4216	2694	722	779	21			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP A1E280
A	-1	ALA	-	expression tag	UNP A1E280
A	0	MET	-	expression tag	UNP A1E280
A	1	GLY	-	expression tag	UNP A1E280
A	52	ILE	VAL	engineered mutation	UNP A1E280
A	82	ILE	VAL	engineered mutation	UNP A1E280
A	360	THR	SER	engineered mutation	UNP A1E280
A	469	SER	GLY	engineered mutation	UNP A1E280
A	470	ASN	SER	engineered mutation	UNP A1E280
B	-2	GLY	-	expression tag	UNP A1E280
B	-1	ALA	-	expression tag	UNP A1E280
B	0	MET	-	expression tag	UNP A1E280
B	1	GLY	-	expression tag	UNP A1E280
B	52	ILE	VAL	engineered mutation	UNP A1E280
B	82	ILE	VAL	engineered mutation	UNP A1E280
B	360	THR	SER	engineered mutation	UNP A1E280
B	469	SER	GLY	engineered mutation	UNP A1E280
B	470	ASN	SER	engineered mutation	UNP A1E280

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	279	Total O 279 279	0	0
4	B	141	Total O 141 141	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tryptophan 6-halogenase



• Molecule 1: Tryptophan 6-halogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α , β , γ	139.26Å 139.26Å 144.33Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.11 – 2.12 48.11 – 2.12	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.11-2.12) 100.0 (48.11-2.12)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.12Å)	Xtriage
Refinement program	PHENIX (1.14rc3_3199: ???)	Depositor
R, R_{free}	0.163 , 0.192 0.163 , 0.192	Depositor DCC
R_{free} test set	4494 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	51.5	Xtriage
Anisotropy	0.195	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 56.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.034 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8962	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/4366	0.59	0/5925
1	B	0.37	0/4338	0.54	1/5887 (0.0%)
All	All	0.40	0/8704	0.56	1/11812 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	115	LEU	CA-CB-CG	5.01	126.83	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4240	0	4109	28	0
1	B	4216	0	4083	26	0
2	A	5	0	0	0	0
2	B	15	0	0	0	0
3	A	36	0	48	2	0
3	B	30	0	40	1	0
4	A	279	0	0	1	0
4	B	141	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8962	0	8280	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:521:TYR:CZ	1:A:525:ARG:HD2	2.25	0.72
1:A:448:VAL:HG23	1:A:449:THR:HG23	1.78	0.64
1:B:521:TYR:CZ	1:B:525:ARG:HD3	2.33	0.63
1:A:330:ARG:NH1	1:A:331:PHE:O	2.36	0.59
1:B:58:ALA:HB1	1:B:366[B]:TYR:CE2	2.38	0.58
1:A:401:ARG:HH12	3:A:607:GOL:H12	1.70	0.56
1:A:366[B]:TYR:HA	1:A:369:ILE:HG12	1.88	0.55
1:B:402:ASP:HB3	1:B:431:ILE:HG12	1.89	0.55
1:A:413:ARG:O	1:A:423:LYS:NZ	2.38	0.54
1:B:366[A]:TYR:HA	1:B:369:ILE:HG12	1.90	0.54
1:A:449:THR:HB	1:A:453:THR:HB	1.91	0.53
1:B:172:PHE:HZ	1:B:362:ILE:HD12	1.74	0.53
1:B:171:HIS:CE1	1:B:288:PRO:HD2	2.45	0.52
1:B:275:SER:HB2	1:B:286:LYS:HB3	1.91	0.51
1:B:410:PHE:CZ	1:B:435:VAL:HG21	2.46	0.51
1:A:362:ILE:HG22	1:A:366[B]:TYR:CE2	2.45	0.51
1:A:58:ALA:HB1	1:A:366[A]:TYR:CZ	2.47	0.50
1:B:244:PHE:CE2	1:B:332:ARG:HG2	2.46	0.49
1:B:505:ASP:OD1	1:B:508:ARG:NH2	2.42	0.49
1:B:448:VAL:HG23	1:B:449:THR:HG23	1.94	0.49
1:B:193:HIS:CE1	1:B:195:GLU:HG2	2.48	0.49
1:B:244:PHE:CZ	1:B:332:ARG:HG2	2.50	0.47
1:A:104:HIS:HE1	4:A:782:HOH:O	1.97	0.47
1:A:18:TRP:CZ2	1:A:182:ARG:HG3	2.48	0.47
1:B:203:THR:OG1	3:B:606:GOL:H2	2.15	0.47
1:A:362:ILE:HG22	1:A:366[B]:TYR:HE2	1.78	0.47
1:B:156:ALA:HB2	1:B:520:THR:HA	1.97	0.47
1:A:244:PHE:CE2	1:A:332:ARG:HG2	2.50	0.46
1:B:257:VAL:HG12	1:B:328:HIS:ND1	2.31	0.45
1:B:18:TRP:CZ2	1:B:182:ARG:HG3	2.51	0.45
1:B:23:TYR:CE2	1:B:366[B]:TYR:HE1	2.34	0.45
1:A:244:PHE:CZ	1:A:332:ARG:HG2	2.52	0.45
1:A:402:ASP:HB3	1:A:431:ILE:HG12	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:75[B]:ASN:OD1	1:B:290:LEU:HB3	2.17	0.44
1:A:171:HIS:CE1	1:A:288:PRO:HD2	2.53	0.44
1:A:77:SER:HA	1:A:524:LEU:HD21	1.98	0.44
1:A:521:TYR:CE2	1:A:525:ARG:HD2	2.53	0.43
1:B:366[B]:TYR:HE2	1:B:458:PHE:CE1	2.37	0.43
1:B:254:ASN:O	1:B:330:ARG:HD2	2.18	0.43
1:A:361:GLY:HA3	3:A:604:GOL:H32	2.01	0.43
1:B:439:ARG:HA	1:B:482:PRO:HA	2.00	0.43
1:A:366[A]:TYR:HE2	1:A:458:PHE:CE1	2.37	0.42
1:A:275:SER:HB2	1:A:286:LYS:HB3	2.02	0.42
1:A:58:ALA:HB1	1:A:366[A]:TYR:CE2	2.54	0.42
1:B:141:ASP:OD1	1:B:142:GLU:N	2.53	0.41
1:A:200:GLU:HB2	1:A:213:ARG:HB3	2.03	0.41
1:A:254:ASN:O	1:A:330:ARG:HD2	2.21	0.41
1:B:58:ALA:HB1	1:B:366[B]:TYR:CZ	2.55	0.41
1:A:439:ARG:HA	1:A:482:PRO:HA	2.03	0.41
1:B:45:ILE:HD11	1:B:260:ALA:HB3	2.02	0.41
1:A:68:GLU:O	1:A:72[A]:ARG:HG3	2.21	0.40
1:A:95:PRO:HD3	1:A:316:LEU:HD12	2.04	0.40
1:A:385:LEU:HD21	1:B:487:PRO:HG2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	527/534 (99%)	513 (97%)	14 (3%)	0	100	100
1	B	524/534 (98%)	510 (97%)	14 (3%)	0	100	100
All	All	1051/1068 (98%)	1023 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	448/446 (100%)	445 (99%)	3 (1%)	85	89
1	B	445/446 (100%)	441 (99%)	4 (1%)	81	85
All	All	893/892 (100%)	886 (99%)	7 (1%)	83	87

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	147	ASP
1	A	307	ASP
1	A	427	LEU
1	B	147	ASP
1	B	311	LEU
1	B	427	LEU
1	B	484	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	526	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	601	-	4,4,4	0.85	0	6,6,6	0.54	0
3	GOL	A	602	-	5,5,5	0.97	0	5,5,5	0.97	0
3	GOL	A	603	-	5,5,5	0.89	0	5,5,5	1.04	0
3	GOL	A	604	-	5,5,5	0.97	0	5,5,5	0.85	0
3	GOL	A	605	-	5,5,5	0.66	0	5,5,5	1.12	0
3	GOL	A	606	-	5,5,5	0.96	0	5,5,5	0.91	0
3	GOL	A	607	-	5,5,5	1.00	0	5,5,5	1.01	0
2	PO4	B	601	-	4,4,4	0.76	0	6,6,6	0.37	0
2	PO4	B	602	-	4,4,4	0.73	0	6,6,6	0.37	0
2	PO4	B	603	-	4,4,4	0.75	0	6,6,6	0.40	0
3	GOL	B	604	-	5,5,5	0.93	0	5,5,5	0.96	0
3	GOL	B	605	-	5,5,5	1.10	0	5,5,5	0.83	0
3	GOL	B	606	-	5,5,5	0.86	0	5,5,5	1.18	0
3	GOL	B	607	-	5,5,5	0.98	0	5,5,5	0.90	0
3	GOL	B	608	-	5,5,5	0.85	0	5,5,5	1.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	601	-	-	0/0/0/0	0/0/0/0
3	GOL	A	602	-	-	0/4/4/4	0/0/0/0
3	GOL	A	603	-	-	0/4/4/4	0/0/0/0
3	GOL	A	604	-	-	0/4/4/4	0/0/0/0
3	GOL	A	605	-	-	0/4/4/4	0/0/0/0
3	GOL	A	606	-	-	0/4/4/4	0/0/0/0
3	GOL	A	607	-	-	0/4/4/4	0/0/0/0
2	PO4	B	601	-	-	0/0/0/0	0/0/0/0
2	PO4	B	602	-	-	0/0/0/0	0/0/0/0
2	PO4	B	603	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	604	-	-	0/4/4/4	0/0/0/0
3	GOL	B	605	-	-	0/4/4/4	0/0/0/0
3	GOL	B	606	-	-	0/4/4/4	0/0/0/0
3	GOL	B	607	-	-	0/4/4/4	0/0/0/0
3	GOL	B	608	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	604	GOL	1	0
3	A	607	GOL	1	0
3	B	606	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	526/534 (98%)	-0.14	9 (1%) 70 74	39, 51, 80, 147	0
1	B	524/534 (98%)	0.03	34 (6%) 19 23	44, 67, 110, 173	0
All	All	1050/1068 (98%)	-0.06	43 (4%) 37 44	39, 58, 102, 173	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	319	LEU	6.5
1	B	42	ILE	5.9
1	B	452	GLY	5.6
1	B	318	GLY	5.2
1	B	326	PHE	5.1
1	B	43	PRO	5.0
1	B	44	ARG	4.7
1	B	324	THR	4.7
1	B	311	LEU	4.2
1	B	317	TRP	4.0
1	A	42	ILE	3.9
1	B	316	LEU	3.7
1	B	521	TYR	3.7
1	B	45	ILE	3.4
1	B	267	LYS	3.2
1	B	270	VAL	3.2
1	B	517	LEU	3.1
1	B	313	PHE	2.9
1	A	454	TYR	2.9
1	B	40	PRO	2.9
1	B	322	ASP	2.9
1	B	310	THR	2.8
1	B	103	GLY	2.8
1	B	290	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	150	ILE	2.6
1	B	269	GLY	2.5
1	B	149	ALA	2.5
1	B	266	GLU	2.5
1	A	451	GLU	2.4
1	B	325	GLU	2.4
1	B	153	ALA	2.3
1	B	323	ASN	2.3
1	B	92	GLU	2.3
1	B	156	ALA	2.3
1	A	452	GLY	2.3
1	B	514	VAL	2.2
1	B	314	SER	2.1
1	A	160	LEU	2.1
1	A	449	THR	2.1
1	A	322	ASP	2.1
1	A	321	PRO	2.1
1	A	450	ASP	2.0
1	B	315	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	607	6/6	0.55	0.23	101,102,105,105	0
3	GOL	B	607	6/6	0.75	0.25	71,83,88,89	1
2	PO4	B	601	5/5	0.79	0.26	162,165,166,166	0
3	GOL	B	608	6/6	0.80	0.17	67,81,91,98	2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	606	6/6	0.81	0.31	92,93,94,94	2
3	GOL	A	602	6/6	0.82	0.23	87,102,105,112	1
3	GOL	B	606	6/6	0.82	0.20	64,83,90,95	2
3	GOL	B	604	6/6	0.82	0.54	103,105,107,107	1
2	PO4	B	602	5/5	0.83	0.18	145,146,146,146	2
3	GOL	B	605	6/6	0.83	0.16	72,79,93,97	1
2	PO4	B	603	5/5	0.84	0.36	144,150,152,153	1
3	GOL	A	603	6/6	0.87	0.40	102,108,112,116	0
3	GOL	A	605	6/6	0.91	0.17	77,78,82,86	1
3	GOL	A	604	6/6	0.94	0.14	54,60,64,78	2
2	PO4	A	601	5/5	0.98	0.07	61,62,67,72	2

6.5 Other polymers [i](#)

There are no such residues in this entry.