



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 30, 2019 – 02:50 PM EDT

PDB ID : 6IBM
Title : Crystal structure of human alpha-galactosidase A in complex with alpha-galactose configured cyclosulfate ME776
Authors : Rowland, R.J.; Wu, L.; Davies, G.J.
Deposited on : 2018-11-30
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.0 (224370), CSD as540be (2019)
Xtriage (Phenix) : 1.13
EDS : 2.4
buster-report : 1.1.7 (2018)
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2250 (2.08-2.04)
Clashscore	122126	2358 (2.08-2.04)
Ramachandran outliers	120053	2339 (2.08-2.04)
Sidechain outliers	120020	2339 (2.08-2.04)
RSRZ outliers	108989	2211 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	398	 86% 12% •
1	B	398	 89% 9% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEG	A	509	-	-	X	-
9	MAN	B	504	-	-	-	X

2 Entry composition [i](#)


There are 10 unique types of molecules in this entry. The entry contains 6948 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-galactosidase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	390	Total	C	N	O	S	0	3	0
			3139	1998	537	578	26			
1	B	392	Total	C	N	O	S	0	0	0
			3128	1992	532	577	27			

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is BETA-D-MANNOSE (three-letter code: BMA) (formula: $C_6H_{12}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		
4	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	A	1	Total O S 5 4 1	0	0
6	B	1	Total O S 5 4 1	0	0
6	B	1	Total O S 5 4 1	0	0
6	B	1	Total O S 5 4 1	0	0

- Molecule 7 is [(1 {S},2 {R},3 {S},4 {S},5 {R},6 {S})-2-(hydroxymethyl)-3,4,5,6-tetrakis(oxidanyl)cyclohexyl] hydrogen sulfate (three-letter code: HF8) (formula: C₇H₁₄O₉S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O S 16 7 8 1	0	0
7	B	1	Total C O S 16 7 8 1	0	0

- Molecule 8 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			4	2	2		
8	A	1	Total	C	O	0	0
			4	2	2		
8	A	1	Total	C	O	0	0
			4	2	2		
8	A	1	Total	C	O	0	0
			4	2	2		
8	A	1	Total	C	O	0	0
			4	2	2		
8	A	1	Total	C	O	0	0
			4	2	2		
8	A	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 9 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			11	6	5		
9	B	1	Total	C	O	0	0
			11	6	5		
9	B	1	Total	C	O	0	0
			11	6	5		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	157	Total	O	0	0
			157	157		
10	B	115	Total	O	0	0
			115	115		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alpha-galactosidase A



• Molecule 1: Alpha-galactosidase A



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	90.34Å 90.34Å 216.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	78.36 – 2.07 78.24 – 2.07	Depositor EDS
% Data completeness (in resolution range)	99.9 (78.36-2.07) 100.0 (78.24-2.07)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.07Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.180 , 0.242 0.186 , 0.240	Depositor DCC
R_{free} test set	3073 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	44.1	Xtriage
Anisotropy	0.483	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.025 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6948	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HF8, BMA, NAG, EDO, SO4, ACT, PEG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	1/3226 (0.0%)	0.81	1/4382 (0.0%)
1	B	0.71	1/3215 (0.0%)	0.81	0/4368
All	All	0.70	2/6441 (0.0%)	0.81	1/8750 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	170	ASP	CG-OD2	8.73	1.45	1.25
1	A	170	ASP	CG-OD2	8.33	1.44	1.25

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	220	ARG	CG-CD-NE	-6.57	98.01	111.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3139	0	2979	33	0
1	B	3128	0	2971	25	0
2	A	84	0	73	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	84	0	73	0	0
3	A	22	0	20	0	0
3	B	22	0	18	0	0
4	A	14	0	20	5	0
5	A	12	0	9	1	0
5	B	4	0	3	0	0
6	A	15	0	0	1	0
6	B	15	0	0	1	0
7	A	16	0	0	0	0
7	B	16	0	0	0	0
8	A	36	0	54	0	0
8	B	36	0	54	0	0
9	B	33	0	29	0	0
10	A	157	0	0	2	0
10	B	115	0	0	0	0
All	All	6948	0	6303	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:369:VAL:HG13	1:B:378:CYS:SG	2.25	0.77
1:A:403:LEU:HD21	1:A:415:LEU:CD1	2.22	0.69
1:B:209:TRP:N	1:B:210:PRO:HD2	2.15	0.62
1:A:277:TRP:H	4:A:509:PEG:C3	2.13	0.62
1:A:403:LEU:HD21	1:A:415:LEU:HD11	1.84	0.60
1:A:402:ARG:NH2	6:A:514:SO4:O1	2.31	0.59
1:A:338:GLU:OE2	1:A:356:ARG:NH1	2.37	0.58
1:B:403:LEU:HD21	1:B:415:LEU:CD1	2.35	0.57
1:B:403:LEU:HD21	1:B:415:LEU:HD13	1.86	0.57
1:B:177:LEU:H	1:B:177:LEU:HD12	1.69	0.57
1:B:353:MET:HE1	1:B:415:LEU:HD11	1.87	0.56
1:A:46:HIS:CD2	1:A:92:ASP:H	2.25	0.54
1:A:228:ASN:HB3	1:A:245:TRP:CH2	2.41	0.54
1:A:277:TRP:H	4:A:509:PEG:H31	1.72	0.54
1:A:251:GLU:CD	1:A:251:GLU:H	2.11	0.54
1:B:32:LEU:N	1:B:220:ARG:O	2.42	0.53
1:A:204:TRP:HB3	1:A:205:PRO:HD3	1.90	0.53
1:A:239:ILE:HD11	1:A:275:LEU:HD11	1.90	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:345:SER:HA	5:A:512:ACT:H2	1.91	0.52
1:A:332:ARG:HD2	10:A:724:HOH:O	2.10	0.51
1:B:369:VAL:CG1	1:B:378:CYS:CB	2.88	0.51
1:A:277:TRP:CZ3	1:A:306:GLN:HG2	2.47	0.49
1:B:216:TYR:HB3	1:B:256:VAL:HG11	1.94	0.49
1:B:208:MET:C	1:B:210:PRO:HD2	2.33	0.49
1:A:97:ALA:HB3	1:A:107:GLN:HG3	1.94	0.49
1:B:228:ASN:HB3	1:B:245:TRP:CH2	2.47	0.49
1:B:353:MET:O	1:B:412:THR:HA	2.14	0.47
1:B:369:VAL:CG1	1:B:378:CYS:SG	2.98	0.47
1:A:157:GLN:NE2	10:A:607:HOH:O	2.48	0.46
1:A:366:THR:HA	1:A:403:LEU:O	2.16	0.46
1:A:205:PRO:HG3	1:A:219:ILE:HD13	1.97	0.46
1:A:73:ALA:HB2	1:A:89:LEU:CD2	2.46	0.46
1:A:76:MET:HA	1:A:81:TRP:HB2	1.98	0.45
1:A:118:ARG:H	4:A:510:PEG:H21	1.82	0.45
1:B:136:ASP:O	1:B:148:SER:HB2	2.17	0.45
1:A:386:GLN:HA	1:A:415:LEU:HD23	1.98	0.44
1:B:383:PHE:CE1	1:B:395:GLY:HA2	2.52	0.44
1:A:277:TRP:CB	4:A:509:PEG:H31	2.48	0.44
1:A:174:CYS:HA	2:A:506:NAG:H81	1.99	0.44
1:B:177:LEU:HD12	1:B:177:LEU:N	2.30	0.44
1:A:108:ALA:HB1	1:A:113:PHE:HB2	1.98	0.43
1:B:139:ASN:C	1:B:147:GLY:HA3	2.38	0.43
1:A:145:PHE:HB3	1:A:146:PRO:HD2	2.00	0.43
1:A:239:ILE:CD1	1:A:275:LEU:HD11	2.48	0.43
1:B:177:LEU:H	1:B:177:LEU:CD1	2.30	0.42
1:A:232:ILE:HD13	1:A:267:MET:O	2.19	0.42
1:A:349:TRP:CZ3	1:A:373:GLY:HA2	2.54	0.42
1:B:205:PRO:O	1:B:209:TRP:CD1	2.73	0.42
1:B:382:CYS:HA	1:B:418:GLU:O	2.20	0.42
1:B:236:TRP:NE1	1:B:240:LYS:HD2	2.36	0.41
1:B:209:TRP:N	1:B:210:PRO:CD	2.82	0.41
1:A:277:TRP:HB3	4:A:509:PEG:H31	2.02	0.41
1:A:140:LYS:HB2	1:A:173:TYR:CD2	2.56	0.40
1:A:353:MET:O	1:A:412:THR:HA	2.21	0.40
1:A:118:ARG:HA	1:A:162:TRP:O	2.21	0.40
1:B:258:GLY:HA2	1:B:325:GLY:O	2.21	0.40
1:B:402:ARG:NH2	6:B:513:SO4:O3	2.55	0.40
1:B:206:LEU:HA	1:B:209:TRP:HD1	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	391/398 (98%)	378 (97%)	12 (3%)	1 (0%)	43	34
1	B	390/398 (98%)	372 (95%)	17 (4%)	1 (0%)	43	34
All	All	781/796 (98%)	750 (96%)	29 (4%)	2 (0%)	43	34

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	266	ASP
1	A	266	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	330/339 (97%)	327 (99%)	3 (1%)	81	80
1	B	330/339 (97%)	326 (99%)	4 (1%)	74	72
All	All	660/678 (97%)	653 (99%)	7 (1%)	81	74

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51[A]	MET
1	A	51[B]	MET
1	A	225	HIS
1	B	51	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	53	ASN
1	B	225	HIS
1	B	364	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	46	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

51 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	501	1,2	14,14,15	0.65	0	17,19,21	1.04	1 (5%)
2	NAG	A	502	3,2	14,14,15	0.54	0	17,19,21	1.39	4 (23%)
3	BMA	A	503	2	11,11,12	0.47	0	15,15,17	1.34	1 (6%)
2	NAG	A	504	1,2	14,14,15	0.43	0	17,19,21	1.27	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	505	2	14,14,15	0.40	0	17,19,21	1.60	6 (35%)
2	NAG	A	506	1,2	14,14,15	0.56	0	17,19,21	1.20	2 (11%)
2	NAG	A	507	3,2	14,14,15	0.39	0	17,19,21	1.08	1 (5%)
3	BMA	A	508	2	11,11,12	0.46	0	15,15,17	1.24	1 (6%)
4	PEG	A	509	-	6,6,6	0.16	0	5,5,5	0.29	0
4	PEG	A	510	-	6,6,6	0.52	0	5,5,5	0.40	0
5	ACT	A	511	-	1,3,3	4.36	1 (100%)	0,3,3	0.00	-
5	ACT	A	512	-	1,3,3	3.15	1 (100%)	0,3,3	0.00	-
5	ACT	A	513	-	1,3,3	5.10	1 (100%)	0,3,3	0.00	-
6	SO4	A	514	-	4,4,4	0.44	0	6,6,6	0.13	0
6	SO4	A	515	-	4,4,4	0.23	0	6,6,6	0.16	0
6	SO4	A	516	-	4,4,4	0.28	0	6,6,6	0.08	0
7	HF8	A	517	1	16,16,17	1.10	1 (6%)	18,24,26	1.28	2 (11%)
8	EDO	A	518	-	3,3,3	0.39	0	2,2,2	0.28	0
8	EDO	A	519	-	3,3,3	0.12	0	2,2,2	0.53	0
8	EDO	A	520	-	3,3,3	0.15	0	2,2,2	0.15	0
8	EDO	A	521	-	3,3,3	0.23	0	2,2,2	0.27	0
8	EDO	A	522	-	3,3,3	0.22	0	2,2,2	0.41	0
8	EDO	A	523	-	3,3,3	0.04	0	2,2,2	0.22	0
8	EDO	A	524	-	3,3,3	0.18	0	2,2,2	0.30	0
8	EDO	A	525	-	3,3,3	0.15	0	2,2,2	0.33	0
8	EDO	A	526	-	3,3,3	0.31	0	2,2,2	0.27	0
2	NAG	B	501	1,2	14,14,15	0.45	0	17,19,21	1.21	2 (11%)
2	NAG	B	502	3,2	14,14,15	0.54	0	17,19,21	0.92	0
3	BMA	B	503	9,2	11,11,12	0.44	0	15,15,17	0.74	0
9	MAN	B	504	3	11,11,12	0.70	0	15,15,17	1.13	2 (13%)
9	MAN	B	505	9,3	11,11,12	0.46	0	15,15,17	1.15	1 (6%)
9	MAN	B	506	9	11,11,12	0.27	0	15,15,17	0.78	1 (6%)
2	NAG	B	507	1,2	14,14,15	0.46	0	17,19,21	1.06	2 (11%)
2	NAG	B	508	2	14,14,15	0.30	0	17,19,21	1.36	2 (11%)
2	NAG	B	509	1,2	14,14,15	0.49	0	17,19,21	1.24	2 (11%)
2	NAG	B	510	3,2	14,14,15	0.63	0	17,19,21	1.46	1 (5%)
3	BMA	B	511	2	11,11,12	0.45	0	15,15,17	1.17	1 (6%)
5	ACT	B	512	-	1,3,3	3.51	1 (100%)	0,3,3	0.00	-
6	SO4	B	513	-	4,4,4	0.44	0	6,6,6	0.16	0
6	SO4	B	514	-	4,4,4	0.42	0	6,6,6	0.16	0
6	SO4	B	515	-	4,4,4	0.31	0	6,6,6	0.07	0
7	HF8	B	516	1	16,16,17	0.90	1 (6%)	18,24,26	1.35	3 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	EDO	B	517	-	3,3,3	0.03	0	2,2,2	0.19	0
8	EDO	B	518	-	3,3,3	0.18	0	2,2,2	0.28	0
8	EDO	B	519	-	3,3,3	0.13	0	2,2,2	0.16	0
8	EDO	B	520	-	3,3,3	0.09	0	2,2,2	0.02	0
8	EDO	B	521	-	3,3,3	0.15	0	2,2,2	0.32	0
8	EDO	B	522	-	3,3,3	0.07	0	2,2,2	0.08	0
8	EDO	B	523	-	3,3,3	0.42	0	2,2,2	0.63	0
8	EDO	B	524	-	3,3,3	0.14	0	2,2,2	0.18	0
8	EDO	B	525	-	3,3,3	0.15	0	2,2,2	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	501	1,2	-	2/6/23/26	0/1/1/1
2	NAG	A	502	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	503	2	-	0/2/19/22	0/1/1/1
2	NAG	A	504	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	505	2	-	0/6/23/26	0/1/1/1
2	NAG	A	506	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	507	3,2	-	2/6/23/26	0/1/1/1
3	BMA	A	508	2	-	0/2/19/22	0/1/1/1
4	PEG	A	509	-	-	2/4/4/4	-
4	PEG	A	510	-	-	2/4/4/4	-
7	HF8	A	517	1	-	0/7/27/31	0/1/1/1
8	EDO	A	518	-	-	1/1/1/1	-
8	EDO	A	519	-	-	0/1/1/1	-
8	EDO	A	520	-	-	1/1/1/1	-
8	EDO	A	521	-	-	1/1/1/1	-
8	EDO	A	522	-	-	1/1/1/1	-
8	EDO	A	523	-	-	0/1/1/1	-
8	EDO	A	524	-	-	1/1/1/1	-
8	EDO	A	525	-	-	0/1/1/1	-
8	EDO	A	526	-	-	1/1/1/1	-
2	NAG	B	501	1,2	-	2/6/23/26	0/1/1/1
2	NAG	B	502	3,2	-	0/6/23/26	0/1/1/1
3	BMA	B	503	9,2	-	0/2/19/22	0/1/1/1
9	MAN	B	504	3	-	2/2/19/22	0/1/1/1
9	MAN	B	505	9,3	-	1/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	MAN	B	506	9	-	1/2/19/22	0/1/1/1
2	NAG	B	507	1,2	-	2/6/23/26	0/1/1/1
2	NAG	B	508	2	-	1/6/23/26	0/1/1/1
2	NAG	B	509	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	510	3,2	-	3/6/23/26	0/1/1/1
3	BMA	B	511	2	-	0/2/19/22	0/1/1/1
7	HF8	B	516	1	-	1/7/27/31	0/1/1/1
8	EDO	B	517	-	-	1/1/1/1	-
8	EDO	B	518	-	-	0/1/1/1	-
8	EDO	B	519	-	-	1/1/1/1	-
8	EDO	B	520	-	-	0/1/1/1	-
8	EDO	B	521	-	-	1/1/1/1	-
8	EDO	B	522	-	-	1/1/1/1	-
8	EDO	B	523	-	-	0/1/1/1	-
8	EDO	B	524	-	-	1/1/1/1	-
8	EDO	B	525	-	-	1/1/1/1	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	513	ACT	CH3-C	5.10	1.55	1.48
5	A	511	ACT	CH3-C	4.36	1.54	1.48
7	A	517	HF8	O10-C4	-3.77	1.41	1.47
5	B	512	ACT	CH3-C	3.51	1.53	1.48
5	A	512	ACT	CH3-C	3.15	1.52	1.48
7	B	516	HF8	O10-C4	-2.21	1.43	1.47

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	510	NAG	C2-N2-C7	4.35	129.15	122.92
3	A	503	BMA	O5-C5-C6	3.64	112.91	107.15
2	B	508	NAG	O5-C5-C6	3.53	112.74	107.15
3	A	508	BMA	C1-O5-C5	3.53	116.99	112.20
9	B	505	MAN	O2-C2-C1	3.14	115.55	109.17
2	A	502	NAG	O4-C4-C5	3.06	116.95	109.29
7	A	517	HF8	C4-O10-S15	3.02	121.85	117.91
7	A	517	HF8	O10-C4-C5	-2.82	102.53	108.32
7	B	516	HF8	O10-C4-C5	-2.79	102.59	108.32
2	B	507	NAG	C1-C2-N2	2.75	115.19	110.49
2	A	506	NAG	C1-O5-C5	2.73	115.91	112.20
7	B	516	HF8	O11-C2-C3	2.65	115.59	110.04

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	502	NAG	C3-C4-C5	-2.60	105.55	110.23
2	A	505	NAG	O4-C4-C3	-2.55	104.43	110.34
2	B	509	NAG	O7-C7-N2	2.54	126.69	121.95
2	A	506	NAG	O5-C5-C6	2.53	111.16	107.15
2	A	505	NAG	C1-O5-C5	2.52	115.63	112.20
2	A	507	NAG	O4-C4-C3	-2.50	104.55	110.34
9	B	504	MAN	O5-C5-C6	2.48	111.07	107.15
3	B	511	BMA	O5-C5-C6	2.43	111.00	107.15
2	B	501	NAG	C6-C5-C4	-2.43	107.29	113.00
2	A	505	NAG	O3-C3-C4	-2.39	104.80	110.34
2	A	502	NAG	O4-C4-C3	-2.37	104.85	110.34
2	A	505	NAG	C3-C4-C5	2.36	114.47	110.23
2	A	504	NAG	O5-C5-C6	2.34	110.86	107.15
2	A	501	NAG	C4-C3-C2	-2.27	107.69	111.02
9	B	504	MAN	O2-C2-C1	2.27	113.78	109.17
7	B	516	HF8	O11-C2-C1	-2.24	105.16	110.34
2	B	509	NAG	C2-N2-C7	2.23	126.11	122.92
2	A	505	NAG	C2-N2-C7	-2.22	119.75	122.92
2	B	508	NAG	C1-O5-C5	2.15	115.12	112.20
2	A	505	NAG	C1-C2-N2	2.12	114.10	110.49
2	B	507	NAG	O5-C5-C6	2.06	110.41	107.15
9	B	506	MAN	O5-C1-C2	-2.06	107.62	110.79
2	A	502	NAG	C6-C5-C4	2.05	117.82	113.00
2	B	501	NAG	O5-C1-C2	-2.05	108.15	111.36
2	A	504	NAG	O4-C4-C3	-2.03	105.64	110.34

There are no chirality outliers.

All (33) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	510	NAG	C3-C2-N2-C7
9	B	504	MAN	O5-C5-C6-O6
2	A	507	NAG	C4-C5-C6-O6
2	B	510	NAG	C4-C5-C6-O6
2	A	501	NAG	C4-C5-C6-O6
2	A	507	NAG	O5-C5-C6-O6
8	A	526	EDO	O1-C1-C2-O2
2	B	501	NAG	O5-C5-C6-O6
2	B	510	NAG	O5-C5-C6-O6
2	B	507	NAG	C4-C5-C6-O6
9	B	504	MAN	C4-C5-C6-O6
4	A	510	PEG	C1-C2-O2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	501	NAG	O5-C5-C6-O6
8	A	524	EDO	O1-C1-C2-O2
8	A	520	EDO	O1-C1-C2-O2
8	B	517	EDO	O1-C1-C2-O2
2	B	507	NAG	O5-C5-C6-O6
4	A	510	PEG	O1-C1-C2-O2
9	B	506	MAN	O5-C5-C6-O6
8	A	518	EDO	O1-C1-C2-O2
8	A	522	EDO	O1-C1-C2-O2
8	B	524	EDO	O1-C1-C2-O2
4	A	509	PEG	O2-C3-C4-O4
2	B	501	NAG	C4-C5-C6-O6
8	B	519	EDO	O1-C1-C2-O2
4	A	509	PEG	C4-C3-O2-C2
2	B	508	NAG	C3-C2-N2-C7
8	B	522	EDO	O1-C1-C2-O2
8	B	525	EDO	O1-C1-C2-O2
8	A	521	EDO	O1-C1-C2-O2
9	B	505	MAN	C4-C5-C6-O6
7	B	516	HF8	C3-C4-O10-S15
8	B	521	EDO	O1-C1-C2-O2


There are no ring outliers.

3 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	506	NAG	1	0
4	A	509	PEG	4	0
4	A	510	PEG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand HF8 A 517




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	390/398 (97%)	-0.12	1 (0%) 93 94	40, 53, 70, 100	7 (1%)
1	B	392/398 (98%)	-0.09	1 (0%) 93 94	42, 59, 86, 116	4 (1%)
All	All	782/796 (98%)	-0.11	2 (0%) 93 94	40, 55, 81, 116	11 (1%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	209	TRP	2.2
1	A	102	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	B	511	11/12	0.33	0.26	122,136,146,150	0
4	PEG	A	510	7/7	0.58	0.23	65,87,91,93	0
3	BMA	A	508	11/12	0.61	0.18	114,124,137,137	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	ACT	A	511	4/4	0.63	0.16	82,91,92,96	0
3	BMA	A	503	11/12	0.64	0.24	94,106,119,128	0
8	EDO	A	518	4/4	0.70	0.24	64,67,70,70	0
8	EDO	B	525	4/4	0.70	0.17	85,98,102,106	0
9	MAN	B	504	11/12	0.75	0.46	95,111,121,121	0
2	NAG	B	509	14/15	0.75	0.16	78,102,112,120	0
8	EDO	B	518	4/4	0.77	0.15	91,94,95,95	0
9	MAN	B	506	11/12	0.79	0.30	125,134,139,150	0
8	EDO	B	524	4/4	0.80	0.14	74,75,77,83	0
8	EDO	A	526	4/4	0.81	0.25	74,83,84,84	0
2	NAG	B	510	14/15	0.81	0.23	112,124,137,140	0
8	EDO	A	524	4/4	0.83	0.31	64,80,81,84	0
4	PEG	A	509	7/7	0.84	0.12	63,71,75,86	0
5	ACT	A	512	4/4	0.84	0.23	57,68,72,76	0
2	NAG	B	508	14/15	0.84	0.20	130,139,143,155	0
2	NAG	A	505	14/15	0.84	0.18	91,100,106,109	0
8	EDO	A	523	4/4	0.84	0.18	74,74,75,81	0
8	EDO	B	517	4/4	0.85	0.13	78,82,83,83	0
8	EDO	B	521	4/4	0.85	0.18	90,94,97,104	0
5	ACT	A	513	4/4	0.85	0.21	69,78,78,87	0
6	SO4	B	515	5/5	0.85	0.26	122,123,129,142	0
8	EDO	A	522	4/4	0.87	0.29	63,75,77,80	0
8	EDO	B	523	4/4	0.87	0.18	69,74,75,76	0
2	NAG	A	506	14/15	0.88	0.09	71,81,87,91	0
8	EDO	A	519	4/4	0.88	0.24	56,69,70,73	0
8	EDO	A	521	4/4	0.88	0.30	43,45,49,53	4
2	NAG	A	507	14/15	0.89	0.15	91,104,119,130	0
3	BMA	B	503	11/12	0.89	0.13	75,95,105,106	0
8	EDO	A	525	4/4	0.89	0.13	80,81,82,87	0
6	SO4	A	516	5/5	0.90	0.15	94,102,118,120	0
9	MAN	B	505	11/12	0.90	0.12	82,100,112,117	0
2	NAG	B	501	14/15	0.92	0.11	64,74,83,83	0
8	EDO	B	522	4/4	0.92	0.27	65,70,72,74	0
2	NAG	B	507	14/15	0.92	0.09	83,107,117,126	0
8	EDO	B	520	4/4	0.92	0.11	84,87,89,99	0
2	NAG	B	502	14/15	0.92	0.11	79,85,89,89	0
6	SO4	B	514	5/5	0.93	0.16	58,59,63,66	5
2	NAG	A	504	14/15	0.93	0.09	72,76,94,96	0
5	ACT	B	512	4/4	0.94	0.11	77,81,84,86	0
6	SO4	A	515	5/5	0.94	0.10	74,78,81,92	0
8	EDO	A	520	4/4	0.94	0.18	61,62,64,72	0
2	NAG	A	502	14/15	0.95	0.10	62,79,91,96	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
8	EDO	B	519	4/4	0.96	0.24	70,73,73,77	0
6	SO4	A	514	5/5	0.97	0.11	63,74,82,83	0
2	NAG	A	501	14/15	0.98	0.11	57,61,66,68	0
7	HF8	A	517	16/17	0.98	0.09	45,53,62,68	0
7	HF8	B	516	16/17	0.98	0.09	48,54,64,74	0
6	SO4	B	513	5/5	0.99	0.09	69,72,79,88	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around HF8 A 517:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HF8 B 516:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.