



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 10:58 pm GMT

PDB ID : 1IM3
Title : Crystal Structure of the human cytomegalovirus protein US2 bound to the MHC class I molecule HLA-A2/tax
Authors : Gewurz, B.E.; Gaudet, R.; Tortorella, D.; Wang, E.W.; Ploegh, H.L.; Wiley, D.C.
Deposited on : 2001-05-09
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	trunk30967
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4343 (2.20-2.20)
Clashscore	122126	5027 (2.20-2.20)
Ramachandran outliers	120053	4952 (2.20-2.20)
Sidechain outliers	120020	4953 (2.20-2.20)
RSRZ outliers	108989	4245 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	<div> <div>%</div> <div> <div></div> <div>89%</div> <div>10%</div> <div>.</div> </div> </div>
1	E	275	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>9%</div> <div>.</div> </div> </div>
1	I	275	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>11%</div> </div> </div>
1	M	275	<div> <div>%</div> <div> <div></div> <div>85%</div> <div>13%</div> <div>.</div> </div> </div>
2	B	100	<div> <div>%</div> <div> <div></div> <div>89%</div> <div>10%</div> <div>.</div> </div> </div>
2	F	100	<div> <div>%</div> <div> <div></div> <div>85%</div> <div>14%</div> <div>.</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	J	100	 2% 88% 11% •
2	N	100	 4% 92% 7% •
3	C	9	 89% 11%
3	G	9	 100%
3	K	9	 100%
3	O	9	 78% 22%
4	D	95	 5% 74% 25% •
4	H	95	 3% 82% 17% •
4	L	95	 20% 75% 23% •
4	P	95	 11% 76% 22% •

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 17162 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			2247	1403	409	426	9			
1	E	275	Total	C	N	O	S	0	0	0
			2247	1403	409	426	9			
1	I	275	Total	C	N	O	S	0	0	0
			2247	1403	409	426	9			
1	M	275	Total	C	N	O	S	0	0	0
			2247	1403	409	426	9			

- Molecule 2 is a protein called beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	F	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	J	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	N	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	CLONING ARTIFACT	UNP P61769
F	0	MET	-	CLONING ARTIFACT	UNP P61769
J	0	MET	-	CLONING ARTIFACT	UNP P61769
N	0	MET	-	CLONING ARTIFACT	UNP P61769

- Molecule 3 is a protein called Human T-cell lymphotropic virus type 1 Tax peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	0	0	0
			77	56	9	12			
3	G	9	Total	C	N	O	0	0	0
			77	56	9	12			
3	K	9	Total	C	N	O	0	0	0
			77	56	9	12			
3	O	9	Total	C	N	O	0	0	0
			77	56	9	12			

- Molecule 4 is a protein called cytomegalovirus protein US2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	95	Total	C	N	O	S	0	0	0
			768	497	129	138	4			
4	H	95	Total	C	N	O	S	0	0	0
			768	497	129	138	4			
4	L	95	Total	C	N	O	S	0	0	0
			768	497	129	138	4			
4	P	95	Total	C	N	O	S	0	0	0
			768	497	129	138	4			

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	247	Total	O	0	0
			247	247		
5	B	63	Total	O	0	0
			63	63		
5	C	11	Total	O	0	0
			11	11		
5	D	79	Total	O	0	0
			79	79		
5	E	250	Total	O	0	0
			250	250		
5	F	67	Total	O	0	0
			67	67		
5	G	12	Total	O	0	0
			12	12		
5	H	58	Total	O	0	0
			58	58		
5	I	214	Total	O	0	0
			214	214		
5	J	60	Total	O	0	0
			60	60		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	K	9	Total 9	O 9	0	0
5	L	36	Total 36	O 36	0	0
5	M	220	Total 220	O 220	0	0
5	N	53	Total 53	O 53	0	0
5	O	11	Total 11	O 11	0	0
5	P	56	Total 56	O 56	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN




- Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN




- Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN



- Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN





- Molecule 2: beta-2-microglobulin







- Molecule 2: beta-2-microglobulin



- Molecule 2: beta-2-microglobulin



- Molecule 2: beta-2-microglobulin



- Molecule 3: Human T-cell lymphotropic virus type 1 Tax peptide



- Molecule 3: Human T-cell lymphotropic virus type 1 Tax peptide




There are no outlier residues recorded for this chain.

- Molecule 3: Human T-cell lymphotropic virus type 1 Tax peptide




There are no outlier residues recorded for this chain.


- Molecule 3: Human T-cell lymphotropic virus type 1 Tax peptide




- Molecule 4: cytomagalovirus protein US2




- Molecule 4: cytomegalovirus protein US2



- Molecule 4: cytomegalovirus protein US2



- Molecule 4: cytomegalovirus protein US2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	95.48Å 96.70Å 99.54Å 109.02° 109.46° 107.93°	Depositor
Resolution (Å)	38.85 – 2.20 45.66 – 2.20	Depositor EDS
% Data completeness (in resolution range)	94.2 (38.85-2.20) 94.3 (45.66-2.20)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.48 (at 2.20Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.206 , 0.240 0.203 , 0.238	Depositor DCC
R_{free} test set	2713 reflections (1.95%)	wwPDB-VP
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.121	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 50.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.013 for k,h,-h-k-l 0.010 for -k,-h,-l 0.011 for -h,-k,h+k+l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	17162	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/2312	0.70	0/3137
1	E	0.57	0/2312	0.70	1/3137 (0.0%)
1	I	0.54	0/2312	0.71	1/3137 (0.0%)
1	M	0.54	0/2312	0.70	1/3137 (0.0%)
2	B	0.48	0/860	0.67	0/1162
2	F	0.50	0/860	0.66	0/1162
2	J	0.46	0/860	0.66	0/1162
2	N	0.47	0/860	0.66	0/1162
3	C	0.59	0/80	0.86	0/108
3	G	0.61	0/80	0.88	0/108
3	K	0.69	0/80	0.84	0/108
3	O	0.71	0/80	0.84	0/108
4	D	0.54	0/788	0.82	1/1070 (0.1%)
4	H	0.50	0/788	0.72	0/1070
4	L	0.51	0/787	0.70	0/1067
4	P	0.52	0/788	0.74	0/1070
All	All	0.53	0/16159	0.71	4/21905 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	M	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	M	28	VAL	N-CA-C	-5.48	96.20	111.00
1	I	28	VAL	N-CA-C	-5.33	96.62	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	137	LEU	CA-CB-CG	5.13	127.10	115.30
1	E	28	VAL	N-CA-C	-5.03	97.43	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	M	209	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2247	0	2096	29	0
1	E	2247	0	2096	27	0
1	I	2247	0	2096	25	0
1	M	2247	0	2096	32	0
2	B	837	0	803	8	0
2	F	837	0	803	13	0
2	J	837	0	803	10	0
2	N	837	0	803	7	0
3	C	77	0	79	1	0
3	G	77	0	79	0	0
3	K	77	0	79	0	0
3	O	77	0	79	2	0
4	D	768	0	746	20	0
4	H	768	0	746	14	0
4	L	768	0	745	18	0
4	P	768	0	746	22	0
5	A	247	0	0	7	0
5	B	63	0	0	1	0
5	C	11	0	0	1	0
5	D	79	0	0	4	0
5	E	250	0	0	8	0
5	F	67	0	0	3	0
5	G	12	0	0	0	0
5	H	58	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	I	214	0	0	4	0
5	J	60	0	0	2	0
5	K	9	0	0	0	0
5	L	36	0	0	0	0
5	M	220	0	0	5	0
5	N	53	0	0	1	0
5	O	11	0	0	1	0
5	P	56	0	0	3	0
All	All	17162	0	14895	213	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 213 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:P:69:MET:HG2	4:P:124:LEU:HB3	1.46	0.98
4:L:69:MET:HG2	4:L:124:LEU:HB3	1.45	0.98
4:D:69:MET:HG2	4:D:124:LEU:HB3	1.48	0.95
4:H:69:MET:HG2	4:H:124:LEU:HB3	1.48	0.93
4:P:73:VAL:HB	4:P:123:VAL:HG13	1.50	0.92

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/275 (99%)	268 (98%)	5 (2%)	0	100	100
1	E	273/275 (99%)	268 (98%)	5 (2%)	0	100	100
1	I	273/275 (99%)	269 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	M	273/275 (99%)	268 (98%)	5 (2%)	0	100	100
2	B	98/100 (98%)	95 (97%)	3 (3%)	0	100	100
2	F	98/100 (98%)	95 (97%)	3 (3%)	0	100	100
2	J	98/100 (98%)	95 (97%)	3 (3%)	0	100	100
2	N	98/100 (98%)	95 (97%)	3 (3%)	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
3	G	7/9 (78%)	7 (100%)	0	0	100	100
3	K	7/9 (78%)	7 (100%)	0	0	100	100
3	O	7/9 (78%)	7 (100%)	0	0	100	100
4	D	93/95 (98%)	85 (91%)	7 (8%)	1 (1%)	16	13
4	H	93/95 (98%)	82 (88%)	11 (12%)	0	100	100
4	L	92/95 (97%)	84 (91%)	8 (9%)	0	100	100
4	P	93/95 (98%)	86 (92%)	7 (8%)	0	100	100
All	All	1883/1916 (98%)	1818 (96%)	64 (3%)	1 (0%)	53	62

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	50	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/231 (100%)	227 (98%)	4 (2%)	63	77
1	E	231/231 (100%)	227 (98%)	4 (2%)	63	77
1	I	231/231 (100%)	226 (98%)	5 (2%)	55	68
1	M	231/231 (100%)	226 (98%)	5 (2%)	55	68
2	B	95/95 (100%)	94 (99%)	1 (1%)	76	87

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	95/95 (100%)	94 (99%)	1 (1%)	76	87
2	J	95/95 (100%)	94 (99%)	1 (1%)	76	87
2	N	95/95 (100%)	94 (99%)	1 (1%)	76	87
3	C	8/8 (100%)	8 (100%)	0	100	100
3	G	8/8 (100%)	8 (100%)	0	100	100
3	K	8/8 (100%)	8 (100%)	0	100	100
3	O	8/8 (100%)	8 (100%)	0	100	100
4	D	83/83 (100%)	81 (98%)	2 (2%)	52	65
4	H	83/83 (100%)	81 (98%)	2 (2%)	52	65
4	L	83/83 (100%)	80 (96%)	3 (4%)	38	48
4	P	83/83 (100%)	80 (96%)	3 (4%)	38	48
All	All	1668/1668 (100%)	1636 (98%)	32 (2%)	60	74

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	121	LYS
1	I	215	LEU
4	P	51	ARG
1	I	183	ASP
1	I	224	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	86	ASN
1	I	141	GLN
1	M	151	HIS
1	E	224	GLN
2	F	2	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
4	L	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	L	136:GLU	C	137:LEU	N	2.75

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	-0.33	2 (0%) 87 86	19, 33, 52, 75	0
1	E	275/275 (100%)	-0.28	4 (1%) 73 72	17, 32, 53, 76	0
1	I	275/275 (100%)	-0.18	5 (1%) 68 66	22, 35, 54, 73	0
1	M	275/275 (100%)	-0.17	4 (1%) 73 72	21, 34, 54, 76	0
2	B	100/100 (100%)	-0.09	1 (1%) 82 81	24, 43, 72, 82	0
2	F	100/100 (100%)	-0.18	1 (1%) 82 81	20, 42, 69, 81	0
2	J	100/100 (100%)	-0.03	2 (2%) 65 63	23, 44, 72, 82	0
2	N	100/100 (100%)	0.10	4 (4%) 38 36	24, 44, 72, 82	0
3	C	9/9 (100%)	-0.60	0 100 100	24, 29, 37, 49	0
3	G	9/9 (100%)	-0.35	0 100 100	21, 28, 36, 49	0
3	K	9/9 (100%)	-0.07	0 100 100	25, 29, 37, 48	0
3	O	9/9 (100%)	-0.41	0 100 100	23, 28, 36, 47	0
4	D	95/95 (100%)	0.35	5 (5%) 26 25	26, 42, 67, 85	0
4	H	95/95 (100%)	0.17	3 (3%) 47 45	25, 46, 69, 86	0
4	L	95/95 (100%)	0.76	19 (20%) 1 1	29, 49, 71, 91	0
4	P	95/95 (100%)	0.44	10 (10%) 6 5	27, 45, 68, 86	0
All	All	1916/1916 (100%)	-0.07	60 (3%) 49 46	17, 37, 67, 91	0

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	137	LEU	6.7
4	H	137	LEU	6.6
4	P	43	PRO	6.3
4	L	43	PRO	6.0
4	L	137	LEU	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.