



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 09:45 pm GMT

PDB ID : 4IMP
Title : The missing linker: a dimerization motif located within polyketide synthase modules
Authors : Zheng, J.; Keatinge-Clay, A.T.
Deposited on : 2013-01-03
Resolution : 2.57 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	3182 (2.60-2.56)
Clashscore	122126	3541 (2.60-2.56)
Ramachandran outliers	120053	3489 (2.60-2.56)
Sidechain outliers	120020	3489 (2.60-2.56)
RSRZ outliers	108989	3120 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	580	<div> <div>7%</div> <div>68% 18% 10%</div> </div>
1	B	580	<div> <div>7%</div> <div>63% 21% 5% 11%</div> </div>
1	C	580	<div> <div>5%</div> <div>64% 21% 5% 11%</div> </div>
1	D	580	<div> <div>8%</div> <div>62% 22% 5% 11%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 15710 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyketide synthase extender modules 3-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	520	Total	C	N	O	S	0	0	0
			3885	2429	719	727	10			
1	B	518	Total	C	N	O	S	0	0	0
			3871	2422	717	722	10			
1	C	519	Total	C	N	O	S	0	0	0
			3875	2422	718	725	10			
1	D	515	Total	C	N	O	S	0	0	0
			3850	2407	714	719	10			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	EXPRESSION TAG	UNP Q9ALM4
A	-19	GLY	-	EXPRESSION TAG	UNP Q9ALM4
A	-18	SER	-	EXPRESSION TAG	UNP Q9ALM4
A	-17	SER	-	EXPRESSION TAG	UNP Q9ALM4
A	-16	HIS	-	EXPRESSION TAG	UNP Q9ALM4
A	-15	HIS	-	EXPRESSION TAG	UNP Q9ALM4
A	-14	HIS	-	EXPRESSION TAG	UNP Q9ALM4
A	-13	HIS	-	EXPRESSION TAG	UNP Q9ALM4
A	-12	HIS	-	EXPRESSION TAG	UNP Q9ALM4
A	-11	HIS	-	EXPRESSION TAG	UNP Q9ALM4
A	-10	SER	-	EXPRESSION TAG	UNP Q9ALM4
A	-9	SER	-	EXPRESSION TAG	UNP Q9ALM4
A	-8	GLY	-	EXPRESSION TAG	UNP Q9ALM4
A	-7	LEU	-	EXPRESSION TAG	UNP Q9ALM4
A	-6	VAL	-	EXPRESSION TAG	UNP Q9ALM4
A	-5	PRO	-	EXPRESSION TAG	UNP Q9ALM4
A	-4	ARG	-	EXPRESSION TAG	UNP Q9ALM4
A	-3	GLY	-	EXPRESSION TAG	UNP Q9ALM4
A	-2	SER	-	EXPRESSION TAG	UNP Q9ALM4
A	-1	HIS	-	EXPRESSION TAG	UNP Q9ALM4
A	0	MET	-	EXPRESSION TAG	UNP Q9ALM4

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-20	MET	-	EXPRESSION TAG	UNP Q9ALM4
B	-19	GLY	-	EXPRESSION TAG	UNP Q9ALM4
B	-18	SER	-	EXPRESSION TAG	UNP Q9ALM4
B	-17	SER	-	EXPRESSION TAG	UNP Q9ALM4
B	-16	HIS	-	EXPRESSION TAG	UNP Q9ALM4
B	-15	HIS	-	EXPRESSION TAG	UNP Q9ALM4
B	-14	HIS	-	EXPRESSION TAG	UNP Q9ALM4
B	-13	HIS	-	EXPRESSION TAG	UNP Q9ALM4
B	-12	HIS	-	EXPRESSION TAG	UNP Q9ALM4
B	-11	HIS	-	EXPRESSION TAG	UNP Q9ALM4
B	-10	SER	-	EXPRESSION TAG	UNP Q9ALM4
B	-9	SER	-	EXPRESSION TAG	UNP Q9ALM4
B	-8	GLY	-	EXPRESSION TAG	UNP Q9ALM4
B	-7	LEU	-	EXPRESSION TAG	UNP Q9ALM4
B	-6	VAL	-	EXPRESSION TAG	UNP Q9ALM4
B	-5	PRO	-	EXPRESSION TAG	UNP Q9ALM4
B	-4	ARG	-	EXPRESSION TAG	UNP Q9ALM4
B	-3	GLY	-	EXPRESSION TAG	UNP Q9ALM4
B	-2	SER	-	EXPRESSION TAG	UNP Q9ALM4
B	-1	HIS	-	EXPRESSION TAG	UNP Q9ALM4
B	0	MET	-	EXPRESSION TAG	UNP Q9ALM4
C	-20	MET	-	EXPRESSION TAG	UNP Q9ALM4
C	-19	GLY	-	EXPRESSION TAG	UNP Q9ALM4
C	-18	SER	-	EXPRESSION TAG	UNP Q9ALM4
C	-17	SER	-	EXPRESSION TAG	UNP Q9ALM4
C	-16	HIS	-	EXPRESSION TAG	UNP Q9ALM4
C	-15	HIS	-	EXPRESSION TAG	UNP Q9ALM4
C	-14	HIS	-	EXPRESSION TAG	UNP Q9ALM4
C	-13	HIS	-	EXPRESSION TAG	UNP Q9ALM4
C	-12	HIS	-	EXPRESSION TAG	UNP Q9ALM4
C	-11	HIS	-	EXPRESSION TAG	UNP Q9ALM4
C	-10	SER	-	EXPRESSION TAG	UNP Q9ALM4
C	-9	SER	-	EXPRESSION TAG	UNP Q9ALM4
C	-8	GLY	-	EXPRESSION TAG	UNP Q9ALM4
C	-7	LEU	-	EXPRESSION TAG	UNP Q9ALM4
C	-6	VAL	-	EXPRESSION TAG	UNP Q9ALM4
C	-5	PRO	-	EXPRESSION TAG	UNP Q9ALM4
C	-4	ARG	-	EXPRESSION TAG	UNP Q9ALM4
C	-3	GLY	-	EXPRESSION TAG	UNP Q9ALM4
C	-2	SER	-	EXPRESSION TAG	UNP Q9ALM4
C	-1	HIS	-	EXPRESSION TAG	UNP Q9ALM4
C	0	MET	-	EXPRESSION TAG	UNP Q9ALM4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-20	MET	-	EXPRESSION TAG	UNP Q9ALM4
D	-19	GLY	-	EXPRESSION TAG	UNP Q9ALM4
D	-18	SER	-	EXPRESSION TAG	UNP Q9ALM4
D	-17	SER	-	EXPRESSION TAG	UNP Q9ALM4
D	-16	HIS	-	EXPRESSION TAG	UNP Q9ALM4
D	-15	HIS	-	EXPRESSION TAG	UNP Q9ALM4
D	-14	HIS	-	EXPRESSION TAG	UNP Q9ALM4
D	-13	HIS	-	EXPRESSION TAG	UNP Q9ALM4
D	-12	HIS	-	EXPRESSION TAG	UNP Q9ALM4
D	-11	HIS	-	EXPRESSION TAG	UNP Q9ALM4
D	-10	SER	-	EXPRESSION TAG	UNP Q9ALM4
D	-9	SER	-	EXPRESSION TAG	UNP Q9ALM4
D	-8	GLY	-	EXPRESSION TAG	UNP Q9ALM4
D	-7	LEU	-	EXPRESSION TAG	UNP Q9ALM4
D	-6	VAL	-	EXPRESSION TAG	UNP Q9ALM4
D	-5	PRO	-	EXPRESSION TAG	UNP Q9ALM4
D	-4	ARG	-	EXPRESSION TAG	UNP Q9ALM4
D	-3	GLY	-	EXPRESSION TAG	UNP Q9ALM4
D	-2	SER	-	EXPRESSION TAG	UNP Q9ALM4
D	-1	HIS	-	EXPRESSION TAG	UNP Q9ALM4
D	0	MET	-	EXPRESSION TAG	UNP Q9ALM4


- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	26	Total	O	0	0
			26	26		
3	B	7	Total	O	0	0
			7	7		
3	C	1	Total	O	0	0
			1	1		
3	D	3	Total	O	0	0
			3	3		



• Molecule 1: Polyketide synthase extender modules 3-4



• Molecule 1: Polyketide synthase extender modules 3-4



ARG
HIS
GLU
LEU
VAL

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.81Å 211.69Å 101.74Å 90.00° 95.26° 90.00°	Depositor
Resolution (Å)	50.00 – 2.57 47.57 – 2.57	Depositor EDS
% Data completeness (in resolution range)	96.2 (50.00-2.57) 96.4 (47.57-2.57)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.17 (at 2.58Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.204 , 0.245 0.204 , 0.247	Depositor DCC
R_{free} test set	3670 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	55.0	Xtriage
Anisotropy	0.033	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15710	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	2/3958 (0.1%)	0.61	1/5395 (0.0%)
1	B	0.62	0/3944	0.56	1/5375 (0.0%)
1	C	0.60	0/3947	0.56	2/5379 (0.0%)
1	D	0.60	0/3920	0.55	2/5339 (0.0%)
All	All	0.63	2/15769 (0.0%)	0.57	6/21488 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	5
1	C	0	2
1	D	0	1
All	All	0	12

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	222	TRP	CD2-CE2	5.14	1.47	1.41
1	A	296	GLN	CD-OE1	5.00	1.34	1.24

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	77	PRO	N-CA-CB	7.01	111.72	103.30
1	D	199	LEU	CA-CB-CG	5.30	127.48	115.30
1	B	104	ASP	CB-CG-OD2	5.25	123.03	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	104	ASP	CB-CG-OD2	5.24	123.01	118.30
1	A	104	ASP	CB-CG-OD2	5.22	123.00	118.30

There are no chirality outliers.

5 of 12 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	400	ASP	Peptide
1	A	462	GLY	Peptide
1	A	528	ALA	Peptide
1	A	79	ARG	Peptide
1	B	23	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3885	0	3892	86	0
1	B	3871	0	3884	113	0
1	C	3875	0	3874	104	0
1	D	3850	0	3859	95	0
2	A	48	0	26	3	0
2	B	48	0	26	2	0
2	C	48	0	26	7	0
2	D	48	0	26	3	0
3	A	26	0	0	0	0
3	B	7	0	0	0	0
3	C	1	0	0	0	0
3	D	3	0	0	0	0
All	All	15710	0	15613	390	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 390 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:469:ARG:HG2	1:D:469:ARG:HH21	1.10	1.15
1:B:90:PRO:HG2	1:B:94:ILE:HD11	1.29	1.10
1:C:80:ARG:HG3	1:C:81:THR:H	1.08	1.09
1:C:470:ARG:HG2	1:C:470:ARG:HH21	1.10	1.07
1:A:239:ARG:HH21	1:A:239:ARG:HG3	1.14	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	518/580 (89%)	484 (93%)	29 (6%)	5 (1%)	17	34
1	B	516/580 (89%)	470 (91%)	35 (7%)	11 (2%)	8	14
1	C	517/580 (89%)	459 (89%)	43 (8%)	15 (3%)	5	8
1	D	511/580 (88%)	469 (92%)	29 (6%)	13 (2%)	6	10
All	All	2062/2320 (89%)	1882 (91%)	136 (7%)	44 (2%)	8	14

5 of 44 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	220	GLU
1	B	272	GLN
1	B	322	ALA
1	B	360	ALA
1	C	42	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	383/429 (89%)	342 (89%)	41 (11%)	7	12
1	B	381/429 (89%)	326 (86%)	55 (14%)	3	6
1	C	380/429 (89%)	340 (90%)	40 (10%)	7	13
1	D	378/429 (88%)	320 (85%)	58 (15%)	3	4
All	All	1522/1716 (89%)	1328 (87%)	194 (13%)	5	8

5 of 194 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	383	GLU
1	C	126	GLU
1	D	358	VAL
1	B	405	ASP
1	C	49	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NDP	A	1501	-	45,52,52	1.77	13 (28%)	54,80,80	1.56	9 (16%)
2	NDP	B	1501	-	45,52,52	1.46	8 (17%)	54,80,80	1.66	10 (18%)
2	NDP	C	1501	-	45,52,52	1.44	8 (17%)	54,80,80	1.53	11 (20%)
2	NDP	D	1501	-	45,52,52	1.42	6 (13%)	54,80,80	1.62	8 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	1501	-	-	0/30/77/77	0/5/5/5
2	NDP	B	1501	-	-	0/30/77/77	0/5/5/5
2	NDP	C	1501	-	-	0/30/77/77	0/5/5/5
2	NDP	D	1501	-	-	0/30/77/77	0/5/5/5

The worst 5 of 35 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1501	NDP	C8A-N9A	-5.06	1.30	1.36
2	D	1501	NDP	C8A-N9A	-4.09	1.32	1.36
2	C	1501	NDP	C8A-N9A	-3.66	1.32	1.36
2	B	1501	NDP	C8A-N9A	-3.45	1.32	1.36
2	A	1501	NDP	C4N-C5N	-3.25	1.42	1.49

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1501	NDP	N3A-C2A-N1A	-7.64	122.32	128.86
2	B	1501	NDP	N3A-C2A-N1A	-6.59	123.22	128.86
2	A	1501	NDP	N3A-C2A-N1A	-6.58	123.23	128.86
2	C	1501	NDP	N3A-C2A-N1A	-5.64	124.04	128.86
2	B	1501	NDP	O3B-C3B-C4B	-3.49	100.96	111.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1501	NDP	3	0
2	B	1501	NDP	2	0
2	C	1501	NDP	7	0
2	D	1501	NDP	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	520/580 (89%)	-0.02	8 (1%) 73 71	26, 46, 78, 137	0
1	B	518/580 (89%)	0.44	39 (7%) 14 11	34, 63, 98, 155	0
1	C	519/580 (89%)	0.30	28 (5%) 26 21	40, 67, 103, 139	0
1	D	515/580 (88%)	0.55	48 (9%) 8 6	40, 68, 103, 135	0
All	All	2072/2320 (89%)	0.32	123 (5%) 22 18	26, 61, 100, 155	0

The worst 5 of 123 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	529	MET	6.4
1	D	325	LEU	5.6
1	D	309	ILE	5.1
1	C	78	VAL	4.7
1	D	320	ALA	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NDP	B	1501	48/48	0.95	0.12	40,53,69,73	0
2	NDP	D	1501	48/48	0.95	0.12	39,61,85,97	0
2	NDP	C	1501	48/48	0.96	0.11	42,54,64,74	0
2	NDP	A	1501	48/48	0.98	0.13	22,34,39,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.