



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2018 – 01:50 pm GMT

PDB ID : 1J07
Title : Crystal structure of the mouse acetylcholinesterase-decidium complex
Authors : Bourne, Y.; Taylor, P.; Radic, Z.; Marchot, P.
Deposited on : 2002-11-07
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1015 (2.36-2.36)
Clashscore	122126	1081 (2.36-2.36)
Ramachandran outliers	120053	1066 (2.36-2.36)
Sidechain outliers	120020	1067 (2.36-2.36)
RSRZ outliers	108989	1002 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	543	<div> <div>6%</div> <div>83%</div> <div>15%</div> <div>..</div> </div>
1	B	543	<div> <div>7%</div> <div>80%</div> <div>17%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	544	-	-	-	X
2	NAG	B	601	-	-	-	X
3	FUC	A	545	-	-	-	X

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 8904 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called acetylcholinesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	536	Total	C	N	O	S	0	0	0
			4184	2684	726	760	14			
1	B	534	Total	C	N	O	S	0	0	0
			4171	2678	722	757	14			

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula: $C_6H_{12}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 4 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	1	3		
4	B	1	Total	C	O	0	0
			4	1	3		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total Cl 2 2	0	0

- Molecule 6 is 3,8-DIAMINO-5,10'-(TRIMETHYLAMMONIUM)DECYL-6-PHENYL PHENANTHRIDINIUM (three-letter code: DCU) (formula: $C_{32}H_{44}N_4$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C N 36 32 4	0	0
6	B	1	Total C N 36 32 4	0	0

- Molecule 7 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: $C_{12}H_{26}O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			19	12	7		

- Molecule 8 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			11	8	3		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	223	Total 223	O 223	0	0
9	B	162	Total 162	O 162	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: acetylcholinesterase



- Molecule 1: acetylcholinesterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.53Å 112.31Å 226.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.91 – 2.35 39.42 – 2.35	Depositor EDS
% Data completeness (in resolution range)	92.8 (29.91-2.35) 92.9 (39.42-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.88 (at 2.34Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.199 , 0.227 0.200 , 0.224	Depositor DCC
R_{free} test set	1589 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å ²)	43.5	Xtriage
Anisotropy	0.745	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 39.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8904	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CO3, NAG, CL, PG4, FUC, P6G, DCU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/4308	0.79	4/5887 (0.1%)
1	B	0.52	0/4295	0.74	1/5869 (0.0%)
All	All	0.54	0/8603	0.77	5/11756 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	257	CYS	C-N-CD	-20.10	76.37	120.60
1	A	257	CYS	C-N-CA	13.68	179.44	122.00
1	B	494	ASP	N-CA-C	-6.96	92.21	111.00
1	A	258	PRO	CA-N-CD	-5.76	103.44	111.50
1	A	161	LEU	CA-CB-CG	-5.53	102.59	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	70	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4184	0	4070	55	0
1	B	4171	0	4061	80	0
2	A	28	0	25	0	0
2	B	14	0	13	0	0
3	A	10	0	10	0	0
4	A	4	0	0	1	0
4	B	4	0	0	1	0
5	A	2	0	0	1	0
6	A	36	0	44	4	0
6	B	36	0	44	3	0
7	B	19	0	26	1	0
8	B	11	0	12	0	0
9	A	223	0	0	4	0
9	B	162	0	0	2	0
All	All	8904	0	8305	136	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 136 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:197:VAL:H	1:B:223:HIS:HD2	1.06	0.95
1:B:424:ARG:HH11	1:B:424:ARG:HG3	1.37	0.89
1:A:245:ARG:O	1:A:249:THR:HG23	1.79	0.83
1:B:197:VAL:H	1:B:223:HIS:CD2	1.98	0.76
1:B:497:SER:HB2	1:B:498:PRO:C	2.07	0.75

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	532/543 (98%)	506 (95%)	24 (4%)	2 (0%)	36	41
1	B	530/543 (98%)	509 (96%)	18 (3%)	3 (1%)	27	30
All	All	1062/1086 (98%)	1015 (96%)	42 (4%)	5 (0%)	31	34

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	494	ASP
1	B	542	ALA
1	A	494	ASP
1	B	497	SER
1	A	523	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	440/443 (99%)	425 (97%)	15 (3%)	40	49
1	B	439/443 (99%)	435 (99%)	4 (1%)	81	88
All	All	879/886 (99%)	860 (98%)	19 (2%)	55	66

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	291	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	301	PRO
1	B	23	LYS
1	A	258	PRO
1	B	295	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	16	GLN
1	B	499	GLN
1	B	223	HIS
1	A	291	GLN
1	B	322	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	544	1,3	14,14,15	0.86	1 (7%)	17,19,21	0.75	0
3	FUC	A	545	2	9,10,11	0.74	0	13,14,16	0.68	0
2	NAG	A	701	1	14,14,15	0.70	0	17,19,21	0.76	1 (5%)
4	CO3	A	903	-	0,3,3	0.00	-	0,3,3	0.00	-
6	DCU	A	951	-	38,39,39	1.84	11 (28%)	48,54,54	1.27	5 (10%)
2	NAG	B	601	1	14,14,15	0.86	0	17,19,21	0.72	0
7	P6G	B	901	-	18,18,18	2.12	6 (33%)	17,17,17	1.17	1 (5%)
8	PG4	B	902	-	10,10,12	1.86	2 (20%)	9,9,11	0.79	0
4	CO3	B	904	-	0,3,3	0.00	-	0,3,3	0.00	-
6	DCU	B	952	-	38,39,39	1.87	11 (28%)	48,54,54	1.35	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	544	1,3	-	0/6/23/26	0/1/1/1
3	FUC	A	545	2	-	0/0/17/20	0/1/1/1
2	NAG	A	701	1	-	0/6/23/26	0/1/1/1
4	CO3	A	903	-	-	0/0/0/0	0/0/0/0
6	DCU	A	951	-	-	0/18/18/18	0/4/4/4
2	NAG	B	601	1	-	0/6/23/26	0/1/1/1
7	P6G	B	901	-	-	0/16/16/16	0/0/0/0
8	PG4	B	902	-	-	0/8/8/10	0/0/0/0
4	CO3	B	904	-	-	0/0/0/0	0/0/0/0
6	DCU	B	952	-	-	0/18/18/18	0/4/4/4

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	951	DCU	C32-C31	2.04	1.42	1.38
6	A	951	DCU	C34-C33	2.08	1.43	1.38
6	B	952	DCU	C32-C31	2.09	1.43	1.38
6	B	952	DCU	C26-C27	2.14	1.43	1.39
6	A	951	DCU	C26-C24	2.19	1.45	1.40

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	952	DCU	C26-C24-C25	-3.99	114.90	120.65

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	951	DCU	C26-C24-C25	-3.97	114.93	120.65
6	B	952	DCU	C2-N23-C24	-3.80	115.41	118.97
6	A	951	DCU	C2-N23-C24	-3.55	115.64	118.97
2	A	701	NAG	C2-N2-C7	-2.21	119.71	122.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	903	CO3	1	0
6	A	951	DCU	4	0
7	B	901	P6G	1	0
4	B	904	CO3	1	0
6	B	952	DCU	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	536/543 (98%)	0.07	35 (6%)	19 28	32, 47, 69, 98	0
1	B	534/543 (98%)	0.18	37 (6%)	17 25	36, 52, 75, 99	0
All	All	1070/1086 (98%)	0.12	72 (6%)	18 27	32, 49, 72, 99	0

The worst 5 of 72 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	543	THR	6.5
1	B	542	ALA	5.0
1	A	495	SER	4.5
1	B	118	ILE	4.5
1	A	497	SER	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	DCU	B	952	36/36	0.62	0.30	88,90,94,94	36
8	PG4	B	902	11/13	0.68	0.22	95,97,98,98	0
2	NAG	B	601	14/15	0.73	0.51	94,97,99,99	0
6	DCU	A	951	36/36	0.74	0.24	87,91,99,99	0
5	CL	A	999	1/1	0.75	0.34	85,85,85,85	0
3	FUC	A	545	10/11	0.75	0.69	97,99,99,99	0
2	NAG	A	544	14/15	0.77	0.51	84,88,92,95	0
2	NAG	A	701	14/15	0.87	0.42	90,94,96,97	0
4	CO3	B	904	4/4	0.88	0.28	74,75,76,76	0
4	CO3	A	903	4/4	0.89	0.31	61,62,64,66	0
5	CL	A	1867	1/1	0.90	0.24	90,90,90,90	1
7	P6G	B	901	19/19	0.92	0.16	59,67,75,77	0

6.5 Other polymers [i](#)

There are no such residues in this entry.