



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 12:35 am GMT

PDB ID : 1J2B
Title : Crystal Structure Of Archaeosine tRNA-Guanine Transglycosylase Complexed With lambda-form tRNA(Val)
Authors : Ishitani, R.; Nureki, O.; Nameki, N.; Okada, N.; Nishimura, S.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2002-12-29
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1168 (3.36-3.24)
Clashscore	122126	1022 (3.34-3.26)
Ramachandran outliers	120053	1004 (3.34-3.26)
Sidechain outliers	120020	1003 (3.34-3.26)
RSRZ outliers	108989	1133 (3.36-3.24)
RNA backbone	2636	1009 (3.74-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	77	<div> <div>22%</div> <div>48%</div> <div>23%</div> <div>6%</div> </div>
1	D	77	<div> <div>17%</div> <div>22%</div> <div>43%</div> <div>27%</div> <div>8%</div> </div>
2	A	582	<div> <div>%</div> <div>31%</div> <div>56%</div> <div>12%</div> <div>.</div> </div>
2	B	582	<div> <div>%</div> <div>29%</div> <div>56%</div> <div>14%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	B	1601	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12495 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called tRNA(Val).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	77	Total	C	N	O	P	0	0	0
			1645	731	294	543	77			
1	D	71	Total	C	N	O	P	0	0	0
			1517	674	272	500	71			

- Molecule 2 is a protein called Archaeosine tRNA-guanine transglycosylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	576	Total	C	N	O	S	0	0	0
			4643	2964	813	847	19			
2	B	576	Total	C	N	O	S	0	0	0
			4643	2964	813	847	19			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Mg	0	0
			2	2		
3	A	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		
4	A	1	Total	Zn	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	9	Total 9	O 9	0	0
5	D	8	Total 8	O 8	0	0
5	A	11	Total 11	O 11	0	0
5	B	13	Total 13	O 13	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: tRNA(Val)




• Molecule 1: tRNA(Val)




• Molecule 2: Archaeosine tRNA-guanine transglycosylase





• Molecule 2: Archaeosine tRNA-guanine transglycosylase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	230.83Å 230.83Å 269.25Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.54 – 3.30 48.54 – 3.16	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.54-3.30) 99.0 (48.54-3.16)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 3.19Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.225 , 0.288 0.221 , 0.274	Depositor DCC
R_{free} test set	4151 reflections (8.85%)	wwPDB-VP
Wilson B-factor (Å ²)	80.5	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 73.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.038 for -2/3*h-1/3*k+2/3*l,-1/3*h-2/3*k-2/3*l,2/3*h-2/3*k+1/3*l 0.035 for -h,1/3*h-1/3*k+2/3*l,2/3*h+4/3*k+1/3*l 0.038 for -1/3*h+1/3*k-2/3*l,-k,-4/3*h-2/3*k+1/3*l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12495	wwPDB-VP
Average B, all atoms (Å ²)	88.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.63	1/1836 (0.1%)	0.84	4/2860 (0.1%)
1	D	0.55	1/1692 (0.1%)	0.80	1/2633 (0.0%)
2	A	0.63	0/4736	0.80	2/6380 (0.0%)
2	B	0.64	0/4736	0.81	0/6380
All	All	0.62	2/13000 (0.0%)	0.81	7/18253 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	5
2	A	0	1
All	All	0	6

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	901	G	OP3-P	-7.11	1.52	1.61
1	D	901	G	OP3-P	-6.99	1.52	1.61

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	961	U	N1-C1'-C2'	7.38	123.60	114.00
1	C	916	U	N1-C1'-C2'	7.24	123.41	114.00
2	A	274	LEU	CA-CB-CG	5.85	128.76	115.30
2	A	236	VAL	CB-CA-C	-5.79	100.39	111.40
1	C	914	A	C1'-O4'-C4'	-5.59	105.43	109.90

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	911	U	Sidechain
1	C	915	G	Sidechain
1	C	916	U	Sidechain
1	C	923	U	Sidechain
1	C	961	U	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1645	0	835	91	0
1	D	1517	0	772	94	0
2	A	4643	0	4735	522	0
2	B	4643	0	4734	568	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
3	C	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	11	0	0	0	0
5	B	13	0	0	0	0
5	C	9	0	0	1	0
5	D	8	0	0	0	0
All	All	12495	0	11076	1202	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 51.

The worst 5 of 1202 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:525:ASP:HA	2:B:577:VAL:HG23	1.28	1.15
2:A:166:ALA:HB1	2:A:184:LEU:HD12	1.24	1.07
1:D:937:C:H3'	1:D:938:G:H5''	1.06	1.05

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:439:TYR:O	2:B:443:ILE:HG12	1.58	1.04
2:A:362:ILE:HD12	2:A:383:ARG:NH2	1.71	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	574/582 (99%)	467 (81%)	86 (15%)	21 (4%)	4	23
2	B	574/582 (99%)	472 (82%)	78 (14%)	24 (4%)	3	19
All	All	1148/1164 (99%)	939 (82%)	164 (14%)	45 (4%)	3	21

5 of 45 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	412	SER
2	A	433	LYS
2	A	452	ALA
2	A	493	ILE
2	A	528	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	498/503 (99%)	432 (87%)	66 (13%)	4	19
2	B	498/503 (99%)	422 (85%)	76 (15%)	3	14
All	All	996/1006 (99%)	854 (86%)	142 (14%)	3	17

5 of 142 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	542	ASP
2	B	172	THR
2	B	488	LEU
2	B	7	MET
2	B	80	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
2	A	471	GLN
2	A	570	GLN
2	B	471	GLN
2	A	339	HIS
2	B	475	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	C	76/77 (98%)	19 (25%)	9 (11%)
1	D	69/77 (89%)	18 (26%)	6 (8%)
All	All	145/154 (94%)	37 (25%)	15 (10%)

5 of 37 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	C	908	U
1	C	909	G
1	C	914	A
1	C	915	G
1	C	916	U

5 of 15 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	C	959	A
1	C	961	U
1	D	916	U
1	C	949	C
1	D	914	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	77/77 (100%)	-0.42	0 100 100	56, 109, 141, 168	0
1	D	71/77 (92%)	0.59	13 (18%) 1 1	76, 150, 198, 200	0
2	A	576/582 (98%)	-0.17	5 (0%) 84 84	32, 74, 121, 162	0
2	B	576/582 (98%)	-0.16	5 (0%) 84 84	27, 70, 115, 142	0
All	All	1300/1318 (98%)	-0.14	23 (1%) 68 67	27, 76, 137, 200	0

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	937	C	4.1
1	D	935	U	3.7
1	D	957	C	3.5
2	B	84	ARG	3.2
1	D	958	A	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MG	B	1601	1/1	0.67	0.41	83,83,83,83	0
3	MG	C	1602	1/1	0.85	0.29	52,52,52,52	0
3	MG	B	602	1/1	0.95	0.29	38,38,38,38	0
3	MG	A	601	1/1	0.96	0.15	85,85,85,85	0
4	ZN	A	600	1/1	0.99	0.16	76,76,76,76	0
4	ZN	B	600	1/1	1.00	0.13	70,70,70,70	0

6.5 Other polymers [i](#)

There are no such residues in this entry.