



wwPDB/EMDataBank EM Map/Model Validation Summary Report ⓘ

Feb 16, 2018 – 01:47 pm GMT

PDB ID : 3J2X
EMDB ID: : EMD-5576
Title : Electron Cryo-microscopy of Chikungunya VLP in complex with neutralizing antibody Fab m242
Authors : Sun, S.; Xiang, Y.; Rossmann, M.G.
Deposited on : 2013-01-28
Resolution : 15.60 Å(reported)
Based on PDB ID : 4GQ8

This is a wwPDB/EMDataBank EM Map/Model Validation Summary Report
for a publicly released PDB/EMDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

MolProbity : 4.02b-467
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30686


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 15.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	136279	1886
Ramachandran outliers	132675	1663
Sidechain outliers	132484	1531

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	217	
1	C	217	
1	E	217	
1	G	217	
2	B	217	
2	D	217	
2	F	217	
2	H	217	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 13228 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called m242 light chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	217	Total	C	N	O	S	0	0
			1676	1041	285	344	6		
1	C	217	Total	C	N	O	S	0	0
			1676	1041	285	344	6		
1	E	217	Total	C	N	O	S	0	0
			1676	1041	285	344	6		
1	G	217	Total	C	N	O	S	0	0
			1676	1041	285	344	6		


- Molecule 2 is a protein called m242 heavy chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	217	Total	C	N	O	S	0	0
			1631	1035	263	326	7		
2	D	217	Total	C	N	O	S	0	0
			1631	1035	263	326	7		
2	F	217	Total	C	N	O	S	0	0
			1631	1035	263	326	7		
2	H	217	Total	C	N	O	S	0	0
			1631	1035	263	326	7		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: m242 light chain




- Molecule 1: m242 light chain




- Molecule 1: m242 light chain




- Molecule 1: m242 light chain




- Molecule 2: m242 heavy chain





- Molecule 2: m242 heavy chain

Chain D: 83% 15%




- Molecule 2: m242 heavy chain

Chain F: 84% 14%



- Molecule 2: m242 heavy chain

Chain H: 84% 14%



4 Experimental information

Property	Value	Source
Reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	1728	Depositor
Resolution determination method	FSC 0.5 CUT-OFF	Depositor
CTF correction method	Each Micrograph	Depositor
Microscope	FEI/PHILIPS CM300FEG/HE	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	20	Depositor
Minimum defocus (nm)	Not provided	Depositor
Maximum defocus (nm)	Not provided	Depositor
Magnification	39190	Depositor
Image detector	Not provided	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >2	RMSZ	# Z >2
1	A	0.31	0/1713	0.45	0/2327
1	C	0.31	0/1713	0.45	0/2327
1	E	0.31	0/1713	0.45	0/2327
1	G	0.31	0/1713	0.45	0/2327
2	B	0.31	0/1674	0.48	0/2287
2	D	0.31	0/1674	0.48	0/2287
2	F	0.31	0/1674	0.48	0/2287
2	H	0.31	0/1674	0.48	0/2287
All	All	0.31	0/13548	0.46	0/18456

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1676	0	1605	12	0
1	C	1676	0	1602	15	0
1	E	1676	0	1602	16	0
1	G	1676	0	1602	16	0
2	B	1631	0	1591	17	0
2	D	1631	0	1591	17	0
2	F	1631	0	1591	16	0
2	H	1631	0	1591	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	13228	0	12775	122	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 122 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:2656:LEU:HD13	2:F:2683:SER:HB2	1.80	0.64
2:B:656:LEU:HD13	2:B:683:SER:HB2	1.80	0.64
1:A:30:ASP:HA	1:A:35:SER:HA	1.81	0.63
1:C:1040:TYR:HE1	1:C:1093:GLN:HG2	1.64	0.63
1:G:3030:ASP:HA	1:G:3035:SER:HA	1.81	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/217 (99%)	199 (93%)	15 (7%)	1 (0%)	31	74
1	C	215/217 (99%)	199 (93%)	15 (7%)	1 (0%)	31	74
1	E	215/217 (99%)	199 (93%)	15 (7%)	1 (0%)	31	74
1	G	215/217 (99%)	199 (93%)	15 (7%)	1 (0%)	31	74
2	B	215/217 (99%)	194 (90%)	16 (7%)	5 (2%)	7	41
2	D	215/217 (99%)	194 (90%)	16 (7%)	5 (2%)	7	41
2	F	215/217 (99%)	194 (90%)	16 (7%)	5 (2%)	7	41
2	H	215/217 (99%)	194 (90%)	16 (7%)	5 (2%)	7	41
All	All	1720/1736 (99%)	1572 (91%)	124 (7%)	24 (1%)	17	52

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	601	GLY
2	D	1601	GLY
2	F	2601	GLY
2	H	3601	GLY
2	B	556	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/191 (100%)	185 (97%)	6 (3%)	43	69
1	C	191/191 (100%)	185 (97%)	6 (3%)	43	69
1	E	191/191 (100%)	185 (97%)	6 (3%)	43	69
1	G	191/191 (100%)	185 (97%)	6 (3%)	43	69
2	B	187/187 (100%)	179 (96%)	8 (4%)	32	61
2	D	187/187 (100%)	179 (96%)	8 (4%)	32	61
2	F	187/187 (100%)	179 (96%)	8 (4%)	32	61
2	H	187/187 (100%)	179 (96%)	8 (4%)	32	61
All	All	1512/1512 (100%)	1456 (96%)	56 (4%)	41	65

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	1657	THR
1	E	2106	THR
2	H	3551	ILE
2	D	1681	LEU
1	E	2032	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	1539	GLN
1	E	2042	GLN
1	G	3042	GLN
1	C	1096	ASN
1	G	3094	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.