



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 9, 2019 – 06:55 AM EDT

PDB ID : 4J23
Title : Low resolution crystal structure of the FGFR2D2D3/FGF1/SR128545 complex
Authors : Kudlinzki, D.; Saxena, K.; Sreeramulu, S.; Schieborr, U.; Dreyer, M.; Schreuder, H.; Schwalbe, H.
Deposited on : 2013-02-04
Resolution : 3.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.3.2
Percentile statistics	:	20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac	:	5.8.0158
CCP4	:	7.0 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.3.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1084 (4.16-3.60)
Clashscore	122126	1155 (4.16-3.60)
Ramachandran outliers	120053	1118 (4.16-3.60)
Sidechain outliers	120020	1111 (4.16-3.60)
RSRZ outliers	108989	1021 (4.18-3.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	
2	B	138	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2642 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fibroblast growth factor receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	198	Total	C	N	O	S	0	0	0
			1571	1001	276	286	8			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	146	MET	-	EXPRESSION TAG	UNP P21802
A	367	LEU	-	EXPRESSION TAG	UNP P21802
A	368	GLU	-	EXPRESSION TAG	UNP P21802

- Molecule 2 is a protein called Fibroblast growth factor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	134	Total	C	N	O	S	0	0	0
			1071	675	187	205	4			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	18	GLY	-	EXPRESSION TAG	UNP P05230
B	19	HIS	-	EXPRESSION TAG	UNP P05230
B	20	MET	-	EXPRESSION TAG	UNP P05230


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Fibroblast growth factor receptor 2



• Molecule 2: Fibroblast growth factor 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	210.67Å 210.67Å 72.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.81 – 3.88 182.45 – 3.88	Depositor EDS
% Data completeness (in resolution range)	99.8 (39.81-3.88) 100.0 (182.45-3.88)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 3.89Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496, REFMAC 5.5.0102	Depositor
R, R_{free}	0.321 , 0.384 0.330 , 0.391	Depositor DCC
R_{free} test set	438 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	175.0	Xtriage
Anisotropy	0.238	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 363.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	2642	wwPDB-VP
Average B, all atoms (Å ²)	284.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	4/1613 (0.2%)	1.38	17/2191 (0.8%)
2	B	0.87	1/1095 (0.1%)	1.19	5/1477 (0.3%)
All	All	0.94	5/2708 (0.2%)	1.31	22/3668 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6
2	B	0	4
All	All	0	10

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	278	CYS	CB-SG	7.16	1.94	1.82
1	A	231	CYS	CB-SG	-6.13	1.71	1.82
1	A	344	ALA	CA-CB	-5.34	1.41	1.52
1	A	178	ARG	CG-CD	5.21	1.65	1.51
2	B	75	GLU	CG-CD	-5.03	1.44	1.51

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	342	CYS	CA-CB-SG	16.34	143.42	114.00
1	A	312	LEU	CA-CB-CG	8.02	133.75	115.30
2	B	121	ASN	N-CA-CB	-6.84	98.28	110.60
1	A	207	TYR	CA-CB-CG	6.59	125.92	113.40
1	A	251	ARG	NE-CZ-NH1	6.54	123.57	120.30

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	173	ASN	Peptide
1	A	197	GLU	Peptide
1	A	198	PHE	Peptide
1	A	199	LYS	Peptide
1	A	218	MET	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1571	0	1541	266	0
2	B	1071	0	1049	176	0
All	All	2642	0	2590	425	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 81.

The worst 5 of 425 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:TRP:CH2	1:A:233:VAL:HG21	1.20	1.62
1:A:156:TRP:CD1	1:A:240:ILE:CG2	1.77	1.58
1:A:156:TRP:NE1	1:A:240:ILE:CG2	1.71	1.49
1:A:156:TRP:CD1	1:A:240:ILE:HG21	0.94	1.46
1:A:156:TRP:CZ2	1:A:233:VAL:HG21	1.50	1.46

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/223 (87%)	134 (69%)	43 (22%)	17 (9%)	1	14
2	B	132/138 (96%)	102 (77%)	20 (15%)	10 (8%)	1	17
All	All	326/361 (90%)	236 (72%)	63 (19%)	27 (8%)	1	15

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	159	THR
1	A	175	VAL
1	A	214	TRP
1	A	280	VAL
1	A	313	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/190 (90%)	138 (81%)	32 (19%)	1	13
2	B	118/120 (98%)	105 (89%)	13 (11%)	7	33
All	All	288/310 (93%)	243 (84%)	45 (16%)	3	20

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	293	HIS
1	A	322	LYS
2	B	103	ARG
1	A	312	LEU
1	A	325	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	242	HIS
2	B	142	GLN
2	B	36	HIS
1	A	213	HIS
1	A	265	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	198/223 (88%)	1.57	64 (32%) 0 0	134, 278, 416, 527	0
2	B	134/138 (97%)	1.63	47 (35%) 0 0	162, 281, 425, 455	0
All	All	332/361 (91%)	1.60	111 (33%) 0 0	134, 279, 424, 527	0

The worst 5 of 111 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	274	VAL	9.4
1	A	352	PHE	7.5
2	B	22	ASN	7.2
2	B	38	LEU	7.1
2	B	68	GLU	6.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.