



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2018 – 12:56 pm GMT

PDB ID : 4JBE
Title : 1.95 Angstrom Crystal Structure of Gamma-glutamyl phosphate Reductase from *Saccharomonospora viridis*.
Authors : Minasov, G.; Filippova, E.V.; Halavaty, A.; Shuvalova, L.; Kiryukhina, O.; Endres, M.; Joachimiak, A.; Anderson, W.F.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2013-02-19
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2220 (1.96-1.96)
Clashscore	122126	2333 (1.96-1.96)
Ramachandran outliers	120053	2314 (1.96-1.96)
Sidechain outliers	120020	2314 (1.96-1.96)
RSRZ outliers	108989	2174 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	445	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red 0%, orange 1%, yellow 87%, green 95%, grey 100%);"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red 0%, orange 1%, yellow 87%, green 95%, grey 100%);"></div> </div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> % 87% 5% 8% </div> </div>
1	B	445	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red 0%, orange 3%, yellow 87%, green 96%, grey 100%);"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red 0%, orange 3%, yellow 87%, green 96%, grey 100%);"></div> </div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> 3% 87% 6% 7% </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7053 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-glutamyl phosphate reductase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	411	Total	C	N	O	S	Se	0	15	0
			3214	1991	596	621	2	4			
1	B	412	Total	C	N	O	S	Se	0	8	0
			3145	1956	569	615	2	3			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-31	MSE	-	EXPRESSION TAG	UNP C7MW73
A	-30	HIS	-	EXPRESSION TAG	UNP C7MW73
A	-29	HIS	-	EXPRESSION TAG	UNP C7MW73
A	-28	HIS	-	EXPRESSION TAG	UNP C7MW73
A	-27	HIS	-	EXPRESSION TAG	UNP C7MW73
A	-26	HIS	-	EXPRESSION TAG	UNP C7MW73
A	-25	HIS	-	EXPRESSION TAG	UNP C7MW73
A	-24	SER	-	EXPRESSION TAG	UNP C7MW73
A	-23	SER	-	EXPRESSION TAG	UNP C7MW73
A	-22	GLY	-	EXPRESSION TAG	UNP C7MW73
A	-21	VAL	-	EXPRESSION TAG	UNP C7MW73
A	-20	ASP	-	EXPRESSION TAG	UNP C7MW73
A	-19	LEU	-	EXPRESSION TAG	UNP C7MW73
A	-18	TRP	-	EXPRESSION TAG	UNP C7MW73
A	-17	SER	-	EXPRESSION TAG	UNP C7MW73
A	-16	HIS	-	EXPRESSION TAG	UNP C7MW73
A	-15	PRO	-	EXPRESSION TAG	UNP C7MW73
A	-14	GLN	-	EXPRESSION TAG	UNP C7MW73
A	-13	PHE	-	EXPRESSION TAG	UNP C7MW73
A	-12	GLU	-	EXPRESSION TAG	UNP C7MW73
A	-11	LYS	-	EXPRESSION TAG	UNP C7MW73
A	-10	GLY	-	EXPRESSION TAG	UNP C7MW73
A	-9	THR	-	EXPRESSION TAG	UNP C7MW73
A	-8	GLU	-	EXPRESSION TAG	UNP C7MW73
A	-7	ASN	-	EXPRESSION TAG	UNP C7MW73

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	LEU	-	EXPRESSION TAG	UNP C7MW73
A	-5	TYR	-	EXPRESSION TAG	UNP C7MW73
A	-4	PHE	-	EXPRESSION TAG	UNP C7MW73
A	-3	GLN	-	EXPRESSION TAG	UNP C7MW73
A	-2	SER	-	EXPRESSION TAG	UNP C7MW73
A	-1	ASN	-	EXPRESSION TAG	UNP C7MW73
A	0	ALA	-	EXPRESSION TAG	UNP C7MW73
B	-31	MSE	-	EXPRESSION TAG	UNP C7MW73
B	-30	HIS	-	EXPRESSION TAG	UNP C7MW73
B	-29	HIS	-	EXPRESSION TAG	UNP C7MW73
B	-28	HIS	-	EXPRESSION TAG	UNP C7MW73
B	-27	HIS	-	EXPRESSION TAG	UNP C7MW73
B	-26	HIS	-	EXPRESSION TAG	UNP C7MW73
B	-25	HIS	-	EXPRESSION TAG	UNP C7MW73
B	-24	SER	-	EXPRESSION TAG	UNP C7MW73
B	-23	SER	-	EXPRESSION TAG	UNP C7MW73
B	-22	GLY	-	EXPRESSION TAG	UNP C7MW73
B	-21	VAL	-	EXPRESSION TAG	UNP C7MW73
B	-20	ASP	-	EXPRESSION TAG	UNP C7MW73
B	-19	LEU	-	EXPRESSION TAG	UNP C7MW73
B	-18	TRP	-	EXPRESSION TAG	UNP C7MW73
B	-17	SER	-	EXPRESSION TAG	UNP C7MW73
B	-16	HIS	-	EXPRESSION TAG	UNP C7MW73
B	-15	PRO	-	EXPRESSION TAG	UNP C7MW73
B	-14	GLN	-	EXPRESSION TAG	UNP C7MW73
B	-13	PHE	-	EXPRESSION TAG	UNP C7MW73
B	-12	GLU	-	EXPRESSION TAG	UNP C7MW73
B	-11	LYS	-	EXPRESSION TAG	UNP C7MW73
B	-10	GLY	-	EXPRESSION TAG	UNP C7MW73
B	-9	THR	-	EXPRESSION TAG	UNP C7MW73
B	-8	GLU	-	EXPRESSION TAG	UNP C7MW73
B	-7	ASN	-	EXPRESSION TAG	UNP C7MW73
B	-6	LEU	-	EXPRESSION TAG	UNP C7MW73
B	-5	TYR	-	EXPRESSION TAG	UNP C7MW73
B	-4	PHE	-	EXPRESSION TAG	UNP C7MW73
B	-3	GLN	-	EXPRESSION TAG	UNP C7MW73
B	-2	SER	-	EXPRESSION TAG	UNP C7MW73
B	-1	ASN	-	EXPRESSION TAG	UNP C7MW73
B	0	ALA	-	EXPRESSION TAG	UNP C7MW73


- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	4	Total	Ca	0	0
			4	4		
2	A	6	Total	Ca	0	0
			6	6		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	4	Total	Cl	0	1
			5	5		
3	A	8	Total	Cl	0	2
			8	8		

- Molecule 4 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C₂H₆OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	385	Total 398	O 398	0	17
7	B	231	Total 237	O 237	0	7


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Gamma-glutamyl phosphate reductase



- Molecule 1: Gamma-glutamyl phosphate reductase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	265.51Å 55.12Å 64.25Å 90.00° 91.18° 90.00°	Depositor
Resolution (Å)	29.73 – 1.95 29.73 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.73-1.95) 99.6 (29.73-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.91 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.168 , 0.208 0.174 , 0.211	Depositor DCC
R_{free} test set	3417 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	28.0	Xtriage
Anisotropy	0.741	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7053	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CL, EDO, MES, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	0/3259	0.71	2/4429 (0.0%)
1	B	0.50	0/3190	0.65	0/4343
All	All	0.56	0/6449	0.68	2/8772 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	72	ARG	NE-CZ-NH2	7.82	124.21	120.30
1	A	160	ARG	NE-CZ-NH1	5.39	123.00	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3214	0	3228	26	0
1	B	3145	0	3152	19	0
2	A	6	0	0	0	0
2	B	4	0	0	0	0
3	A	8	0	0	0	0
3	B	5	0	0	0	0
4	A	4	0	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	4	0	5	0	0
5	A	4	0	6	1	0
6	A	12	0	13	0	0
6	B	12	0	13	0	0
7	A	398	0	0	5	0
7	B	237	0	0	2	0
All	All	7053	0	6422	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 41 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:65:MSE:HE2	1:B:69:LEU:HB3	1.48	0.96
1:A:99[B]:THR:HG23	1:A:112:VAL:HG11	1.50	0.94
1:A:99[B]:THR:HG23	1:A:112:VAL:CG1	2.15	0.76
1:B:160:ARG:O	1:B:164[B]:VAL:HG22	1.88	0.73
1:A:156[B]:LYS:CE	7:A:680:HOH:O	2.37	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	424/445 (95%)	420 (99%)	4 (1%)	0	100	100
1	B	418/445 (94%)	413 (99%)	5 (1%)	0	100	100
All	All	842/890 (95%)	833 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/344 (97%)	332 (100%)	1 (0%)	93	92
1	B	327/344 (95%)	325 (99%)	2 (1%)	87	86
All	All	660/688 (96%)	657 (100%)	3 (0%)	90	89

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	ARG
1	B	98	ARG
1	B	149	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	254	ASN
1	A	358	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 28 ligands modelled in this entry, 23 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BME	A	515	1	3,3,3	0.55	0	1,2,2	0.68	0
5	EDO	A	516	-	3,3,3	0.36	0	2,2,2	0.10	0
6	MES	A	517	-	12,12,12	2.00	1 (8%)	14,16,16	1.13	2 (14%)
4	BME	B	509	1	3,3,3	0.45	0	1,2,2	0.58	0
6	MES	B	510	-	12,12,12	2.11	1 (8%)	14,16,16	0.98	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BME	A	515	1	-	0/1/1/1	0/0/0/0
5	EDO	A	516	-	-	0/1/1/1	0/0/0/0
6	MES	A	517	-	-	0/6/14/14	0/1/1/1
4	BME	B	509	1	-	0/1/1/1	0/0/0/0
6	MES	B	510	-	-	0/6/14/14	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	510	MES	C8-S	-6.94	1.67	1.77
6	A	517	MES	C8-S	-6.49	1.68	1.77

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	517	MES	O1S-S-C8	-2.10	104.39	106.92
6	B	510	MES	O3S-S-C8	2.15	109.25	105.77
6	A	517	MES	O2S-S-C8	2.96	110.48	106.92

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	516	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	408/445 (91%)	-0.10	5 (1%) 79 84	19, 28, 47, 59	0
1	B	409/445 (91%)	0.06	15 (3%) 41 51	22, 37, 60, 80	0
All	All	817/890 (91%)	-0.02	20 (2%) 59 68	19, 33, 52, 80	0

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	67	ALA	4.7
1	B	62	ALA	3.9
1	A	3	ASN	3.3
1	B	66	SER	3.1
1	B	188	ARG	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CA	B	504	1/1	0.71	0.09	42,42,42,42	1
2	CA	A	505	1/1	0.85	0.11	53,53,53,53	1
3	CL	A	511	1/1	0.88	0.10	60,60,60,60	0
3	CL	A	507	1/1	0.91	0.06	46,46,46,46	0
2	CA	B	503	1/1	0.91	0.06	71,71,71,71	0
3	CL	B	506[B]	1/1	0.91	0.33	44,44,44,44	1
3	CL	B	506[A]	1/1	0.91	0.33	36,36,36,36	1
2	CA	B	502	1/1	0.92	0.14	56,56,56,56	1
3	CL	B	507	1/1	0.94	0.14	56,56,56,56	0
5	EDO	A	516	4/4	0.94	0.12	29,30,36,38	0
4	BME	B	509	4/4	0.95	0.10	43,48,49,50	0
6	MES	B	510	12/12	0.96	0.17	53,58,62,62	0
3	CL	A	514[B]	1/1	0.96	0.22	32,32,32,32	1
3	CL	A	513[A]	1/1	0.96	0.21	33,33,33,33	1
2	CA	A	506	1/1	0.96	0.12	56,56,56,56	0
4	BME	A	515	4/4	0.96	0.10	32,37,37,39	0
2	CA	B	501	1/1	0.97	0.04	42,42,42,42	1
3	CL	A	508	1/1	0.97	0.15	50,50,50,50	0
2	CA	A	502	1/1	0.97	0.11	47,47,47,47	0
6	MES	A	517	12/12	0.97	0.09	28,37,44,46	0
2	CA	A	501	1/1	0.98	0.19	54,54,54,54	0
2	CA	A	503	1/1	0.98	0.05	60,60,60,60	0
3	CL	A	509	1/1	0.98	0.05	46,46,46,46	0
3	CL	B	508	1/1	0.98	0.09	47,47,47,47	0
2	CA	A	504	1/1	0.99	0.11	52,52,52,52	0
3	CL	A	512	1/1	0.99	0.05	36,36,36,36	0
3	CL	B	505	1/1	0.99	0.06	35,35,35,35	0
3	CL	A	510	1/1	1.00	0.05	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.