



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2018 – 05:54 am GMT

PDB ID : 5JCA
Title : NADP(H) bound NADH-dependent Ferredoxin:NADP Oxidoreductase (NfnI) from *Pyrococcus furiosus*
Authors : Zadvornyy, O.A.; Schut, G.J.; Nguyen, D.M.; Artz, J.H.; Tokmina-Lukaszewska, M.; Lipscomb, G.; Adams, M.W.; Peters, J.W.
Deposited on : 2016-04-14
Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2534 (1.50-1.50)
Clashscore	122126	2727 (1.50-1.50)
Ramachandran outliers	120053	2661 (1.50-1.50)
Sidechain outliers	120020	2659 (1.50-1.50)
RSRZ outliers	108989	2481 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	474	<div> <div>8%</div> <div> <div></div> <div>93%</div> <div>7%</div> <div>.</div> </div> </div>
2	S	284	<div> <div>%</div> <div> <div></div> <div>95%</div> <div>.</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NDP	L	504	-	-	X	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 13118 atoms, of which 6046 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH-dependent Ferredoxin:NADP Oxidoreductase (NfnI) subunit alpha.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	L	471	Total	C	H	N	O	S	0	0	0
			7432	2337	3752	650	676	17			


- Molecule 2 is a protein called NADH-dependent Ferredoxin:NADP Oxidoreductase (NfnI) subunit beta.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	S	279	Total	C	H	N	O	S	0	0	0
			4370	1389	2200	370	396	15			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
S	1	MET	-	expression tag	UNP Q8U194
S	2	VAL	-	expression tag	UNP Q8U194
S	3	VAL	-	expression tag	UNP Q8U194
S	4	MET	-	expression tag	UNP Q8U194
S	5	VAL	-	expression tag	UNP Q8U194
S	6	MET	-	expression tag	UNP Q8U194

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	L	1	Total	Fe	S	0	0
			8	4	4		
3	L	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	L	1	Total	C	H	N	O	P	0	0
			83	27	30	9	15	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	S	1	Total	C	H	N	O	P	0	0
			83	27	30	9	15	2		

- Molecule 5 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	L	1	Total	C	H	N	O	P	0	0
			74	21	26	7	17	3		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	L	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 7 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe_2S_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	S	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	S	1	Total	Mg	0	0
			1	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	L	594	Total	O	0	0
			594	594		
9	S	447	Total	O	0	0
			447	447		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NADH-dependent Ferredoxin:NADP Oxidoreductase (NfnI) subunit alpha



- Molecule 2: NADH-dependent Ferredoxin:NADP Oxidoreductase (NfnI) subunit beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	179.68Å 179.68Å 80.67Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.52 – 1.50 47.52 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.8 (47.52-1.50) 98.8 (47.52-1.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.20 (at 1.50Å)	Xtriage
Refinement program	PHENIX (dev_2313: ???)	Depositor
R, R_{free}	0.139 , 0.153 0.162 , 0.176	Depositor DCC
R_{free} test set	11765 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	12.7	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 47.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13118	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, SF4, FES, NDP, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.49	1/3749 (0.0%)	0.70	3/5062 (0.1%)
2	S	0.62	2/2210 (0.1%)	0.89	4/2979 (0.1%)
All	All	0.54	3/5959 (0.1%)	0.77	7/8041 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	S	102	ASP	CB-CG	12.98	1.79	1.51
2	S	102	ASP	CG-OD2	8.93	1.45	1.25
1	L	254	ILE	C-N	-5.49	1.21	1.34

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	S	102	ASP	CB-CG-OD2	24.26	140.13	118.30
2	S	102	ASP	OD1-CG-OD2	-7.82	108.44	123.30
2	S	102	ASP	CB-CG-OD1	-7.73	111.34	118.30
1	L	14	ARG	NE-CZ-NH1	6.77	123.69	120.30
2	S	102	ASP	CB-CA-C	6.28	122.96	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	3680	3752	3750	33	0
2	S	2170	2200	2226	13	2
3	L	16	0	0	0	0
4	L	53	30	31	10	0
4	S	53	30	31	1	0
5	L	48	26	26	23	0
6	L	6	8	8	1	0
7	S	4	0	0	0	0
8	S	1	0	0	0	0
9	L	594	0	0	6	0
9	S	447	0	0	4	0
All	All	7072	6046	6072	52	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 52 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:S:102:ASP:CG	2:S:102:ASP:CB	1.79	1.47
1:L:449:GLU:OE1	5:L:504:NDP:C5D	1.89	1.19
4:L:503:FAD:N10	5:L:504:NDP:O7N	1.81	1.14
1:L:449:GLU:OE1	5:L:504:NDP:H51N	1.47	1.13
4:L:503:FAD:C1'	5:L:504:NDP:O7N	1.97	1.12

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:S:102:ASP:OD2	2:S:102:ASP:OD2[6_555]	1.13	1.07
2:S:102:ASP:OD1	2:S:102:ASP:OD1[6_555]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	469/474 (99%)	455 (97%)	14 (3%)	0	100	100
2	S	277/284 (98%)	271 (98%)	6 (2%)	0	100	100
All	All	746/758 (98%)	726 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	385/388 (99%)	384 (100%)	1 (0%)	93	86
2	S	232/237 (98%)	232 (100%)	0	100	100
All	All	617/625 (99%)	616 (100%)	1 (0%)	94	87

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	65	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	303	ASN
1	L	363	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SF4	L	501	1	0,12,12	0.00	-	0,24,24	0.00	-
3	SF4	L	502	1	0,12,12	0.00	-	0,24,24	0.00	-
4	FAD	L	503	-	50,58,58	1.47	6 (12%)	56,89,89	1.81	9 (16%)
5	NDP	L	504	-	45,52,52	1.04	2 (4%)	54,80,80	1.52	5 (9%)
6	GOL	L	505	-	5,5,5	0.34	0	5,5,5	0.62	0
4	FAD	S	301	8	50,58,58	1.48	6 (12%)	56,89,89	1.81	10 (17%)
7	FES	S	302	2	0,4,4	0.00	-	0,4,4	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SF4	L	501	1	-	0/0/48/48	0/6/5/5
3	SF4	L	502	1	-	0/0/48/48	0/6/5/5
4	FAD	L	503	-	-	0/28/50/50	0/6/6/6
5	NDP	L	504	-	-	0/30/77/77	0/5/5/5
6	GOL	L	505	-	-	0/4/4/4	0/0/0/0
4	FAD	S	301	8	-	0/28/50/50	0/6/6/6
7	FES	S	302	2	-	0/0/4/4	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	S	301	FAD	C9A-N10	2.08	1.41	1.38
4	L	503	FAD	C9A-N10	2.11	1.41	1.38
4	L	503	FAD	C5A-C4A	3.04	1.47	1.40
4	S	301	FAD	C5A-C4A	3.05	1.47	1.40
5	L	504	NDP	C5A-C4A	3.06	1.47	1.40

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	L	504	NDP	N3A-C2A-N1A	-6.93	122.93	128.86
4	L	503	FAD	N3A-C2A-N1A	-6.90	122.96	128.86
4	S	301	FAD	N3A-C2A-N1A	-6.90	122.96	128.86
5	L	504	NDP	PN-O3-PA	-3.61	120.49	132.63
4	S	301	FAD	P-O3P-PA	-3.60	120.52	132.63

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	L	503	FAD	10	0
5	L	504	NDP	23	0
6	L	505	GOL	1	0
4	S	301	FAD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	471/474 (99%)	0.25	38 (8%) 12 13	8, 18, 43, 74	0
2	S	279/284 (98%)	-0.12	4 (1%) 75 80	9, 15, 29, 68	0
All	All	750/758 (98%)	0.11	42 (5%) 24 27	8, 17, 39, 74	0

The worst 5 of 42 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	386	ILE	7.9
1	L	374	LEU	6.0
1	L	376	GLU	6.0
2	S	6	MET	5.7
1	L	49	TYR	5.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NDP	L	504	48/48	0.69	0.42	22,45,81,92	0
4	FAD	L	503	53/53	0.77	0.22	7,11,16,17	0
6	GOL	L	505	6/6	0.87	0.21	23,37,48,58	0
3	SF4	L	502	8/8	0.89	0.17	13,13,14,14	0
3	SF4	L	501	8/8	0.90	0.18	10,10,11,11	0
4	FAD	S	301	53/53	0.93	0.14	7,10,13,19	0
8	MG	S	303	1/1	0.98	0.19	9,9,9,9	0
7	FES	S	302	4/4	1.00	0.10	9,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.