



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2018 – 02:42 PM EDT

PDB ID : 1JRB  
Title : The P56A mutant of Lactococcus lactis dihydroorotate dehydrogenase A  
Authors : Norager, S.; Arent, S.; Bjornberg, O.; Ottosen, M.; Lo Leggio, L.; Jensen, K.F.; Larsen, S.  
Deposited on : 2001-08-13  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031172  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031172


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	5502 (1.90-1.90)
Clashscore	122126	6115 (1.90-1.90)
Ramachandran outliers	120053	6048 (1.90-1.90)
Sidechain outliers	120020	6048 (1.90-1.90)
RSRZ outliers	108989	5379 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	311	<div> <div>5%</div> <div> <div></div> <div>78%</div> <div>19%</div> <div>..</div> </div> </div>
1	B	311	<div> <div>5%</div> <div> <div></div> <div>84%</div> <div>14%</div> <div>.</div> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5226 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called dihydroorotate dehydrogenase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	311	Total	C	N	O	S	0	1	0
			2397	1541	385	459	12			
1	B	311	Total	C	N	O	S	0	0	0
			2387	1534	385	457	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	56	ALA	PRO	ENGINEERED	UNP P54321
B	56	ALA	PRO	ENGINEERED	UNP P54321


- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C<sub>17</sub>H<sub>21</sub>N<sub>4</sub>O<sub>9</sub>P).



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 3 is OROTIC ACID (three-letter code: ORO) (formula:  $C_5H_4N_2O_4$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			11	5	2	4		
3	B	1	Total	C	N	O	0	0
			11	5	2	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	180	Total	O	0	0
			180	180		
4	B	178	Total	O	0	0
			178	178		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: dihydroorotate dehydrogenase A



- Molecule 1: dihydroorotate dehydrogenase A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.07Å 108.82Å 66.45Å 90.00° 103.92° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 19.43 – 1.93	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-1.90) 98.8 (19.43-1.93)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.28 (at 1.93Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.196 , 0.236 0.174 , 0.210	Depositor DCC
$R_{free}$ test set	5431 reflections (10.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.2	Xtriage
Anisotropy	0.527	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 54.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5226	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, ORO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.09	4/2451 (0.2%)	1.68	50/3309 (1.5%)
1	B	1.02	2/2436 (0.1%)	1.69	44/3287 (1.3%)
All	All	1.05	6/4887 (0.1%)	1.69	94/6596 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	8
1	B	0	6
All	All	0	14

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	194	SER	CB-OG	-7.79	1.32	1.42
1	B	194	SER	CB-OG	-7.29	1.32	1.42
1	A	40	TYR	CE2-CZ	-6.18	1.30	1.38
1	B	194	SER	CA-CB	5.46	1.61	1.52
1	A	78	TYR	CG-CD1	5.06	1.45	1.39
1	A	194	SER	CA-CB	5.02	1.60	1.52

All (94) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	238	ARG	NE-CZ-NH1	22.52	131.56	120.30
1	A	238	ARG	NE-CZ-NH1	16.66	128.63	120.30
1	A	40	TYR	CB-CG-CD1	15.89	130.54	121.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	40	TYR	CB-CG-CD2	12.68	128.61	121.00
1	B	40	TYR	CD1-CG-CD2	-12.01	104.68	117.90
1	A	40	TYR	CD1-CG-CD2	-12.01	104.69	117.90
1	A	233	ARG	NE-CZ-NH1	10.83	125.71	120.30
1	B	238	ARG	NE-CZ-NH2	-10.82	114.89	120.30
1	A	191	SER	O-C-N	-9.81	107.00	122.70
1	B	55	LEU	O-C-N	-9.75	107.09	122.70
1	B	40	TYR	CB-CG-CD1	9.51	126.70	121.00
1	B	191	SER	O-C-N	-9.49	107.51	122.70
1	B	191	SER	CA-C-N	9.05	137.11	117.20
1	A	40	TYR	CG-CD1-CE1	9.04	128.53	121.30
1	B	284	ARG	NE-CZ-NH1	8.77	124.68	120.30
1	B	127	ASN	OD1-CG-ND2	-8.66	101.97	121.90
1	B	40	TYR	CG-CD1-CE1	8.65	128.22	121.30
1	A	17	ASN	CB-CG-OD1	8.59	138.78	121.60
1	B	17	ASN	OD1-CG-ND2	-8.31	102.79	121.90
1	A	58	TYR	CB-CG-CD1	8.20	125.92	121.00
1	B	40	TYR	CG-CD2-CE2	8.12	127.79	121.30
1	B	40	TYR	CZ-CE2-CD2	7.93	126.94	119.80
1	B	40	TYR	CE1-CZ-CE2	-7.90	107.16	119.80
1	A	97	PHE	CB-CG-CD2	-7.72	115.39	120.80
1	A	127	ASN	OD1-CG-ND2	-7.71	104.17	121.90
1	B	55	LEU	CB-CA-C	-7.70	95.58	110.20
1	A	40	TYR	CG-CD2-CE2	7.67	127.43	121.30
1	A	17	ASN	OD1-CG-ND2	-7.65	104.30	121.90
1	A	40	TYR	CE1-CZ-CE2	-7.60	107.64	119.80
1	A	40	TYR	CZ-CE2-CD2	7.49	126.54	119.80
1	B	55	LEU	CA-C-N	7.36	133.38	117.20
1	B	28	ASP	CB-CG-OD1	7.28	124.85	118.30
1	B	191	SER	N-CA-CB	7.12	121.17	110.50
1	A	169	PHE	CB-CG-CD1	-7.01	115.89	120.80
1	B	229	LEU	CA-C-O	6.86	134.51	120.10
1	B	229	LEU	O-C-N	-6.71	111.96	122.70
1	A	175	ASP	CB-CG-OD1	6.60	124.24	118.30
1	A	55	LEU	O-C-N	-6.54	112.23	122.70
1	A	57	ARG	NE-CZ-NH1	6.52	123.56	120.30
1	B	24	MET	CG-SD-CE	-6.37	90.00	100.20
1	B	79	TYR	CB-CG-CD2	-6.37	117.18	121.00
1	A	23	CYS	O-C-N	-6.33	112.57	122.70
1	A	223	TYR	CB-CG-CD2	-6.30	117.22	121.00
1	A	40	TYR	CB-CG-CD2	6.29	124.78	121.00
1	A	26	ILE	O-C-N	-6.28	112.66	122.70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	303	ASP	CB-CG-OD2	6.24	123.92	118.30
1	B	58	TYR	CB-CG-CD2	-6.23	117.26	121.00
1	A	58	TYR	CB-CG-CD2	-6.23	117.27	121.00
1	A	40	TYR	CD1-CE1-CZ	5.93	125.14	119.80
1	A	49	LYS	CA-CB-CG	5.93	126.45	113.40
1	A	259	GLU	OE1-CD-OE2	5.91	130.39	123.30
1	B	40	TYR	CD1-CE1-CZ	5.86	125.08	119.80
1	B	252	GLU	OE1-CD-OE2	-5.69	116.47	123.30
1	A	192	VAL	N-CA-CB	5.69	124.02	111.50
1	B	127	ASN	CB-CG-OD1	5.68	132.96	121.60
1	A	23	CYS	CA-C-O	5.67	132.01	120.10
1	A	43	LYS	O-C-N	-5.67	113.62	122.70
1	A	194	SER	CB-CA-C	-5.66	99.35	110.10
1	A	127	ASN	CB-CG-ND2	5.65	130.27	116.70
1	B	152	GLU	OE1-CD-OE2	-5.64	116.53	123.30
1	B	78	TYR	CZ-CE2-CD2	5.60	124.84	119.80
1	B	39	ALA	N-CA-CB	5.56	117.88	110.10
1	A	191	SER	CA-C-N	5.53	129.37	117.20
1	A	200	PHE	O-C-N	5.48	131.47	122.70
1	A	55	LEU	CA-C-N	5.46	129.22	117.20
1	A	269	GLN	O-C-N	5.45	131.42	122.70
1	B	191	SER	CA-C-O	-5.45	108.65	120.10
1	B	194	SER	CB-CA-C	-5.44	99.76	110.10
1	A	272	THR	O-C-N	-5.43	114.00	122.70
1	A	238	ARG	NE-CZ-NH2	-5.38	117.61	120.30
1	A	189	VAL	CA-CB-CG1	5.38	118.97	110.90
1	A	192	VAL	O-C-N	5.37	131.29	122.70
1	B	258	PHE	CA-C-O	5.36	131.35	120.10
1	B	238	ARG	NH1-CZ-NH2	-5.34	113.53	119.40
1	B	78	TYR	CG-CD2-CE2	-5.31	117.05	121.30
1	B	262	LEU	CA-CB-CG	5.27	127.43	115.30
1	B	81	ASP	CB-CG-OD1	5.26	123.03	118.30
1	A	168	TYR	CB-CG-CD2	-5.24	117.86	121.00
1	A	97	PHE	CB-CG-CD1	5.23	124.46	120.80
1	A	67	ASN	O-C-N	-5.23	114.33	122.70
1	A	202	ASP	CB-CG-OD1	5.23	123.01	118.30
1	A	214	ASP	CB-CG-OD2	5.22	123.00	118.30
1	A	241	PRO	N-CA-CB	5.22	109.56	103.30
1	A	79	TYR	CD1-CG-CD2	5.18	123.59	117.90
1	B	17	ASN	CB-CG-ND2	5.17	129.11	116.70
1	B	125	GLU	OE1-CD-OE2	-5.16	117.11	123.30
1	A	238	ARG	NH1-CZ-NH2	-5.13	113.75	119.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	141	TYR	CD1-CE1-CZ	5.13	124.42	119.80
1	A	256	ASP	CB-CG-OD2	-5.12	113.69	118.30
1	B	107	GLU	OE1-CD-OE2	-5.07	117.21	123.30
1	B	215	GLY	CA-C-O	-5.07	111.48	120.60
1	A	79	TYR	CB-CG-CD1	-5.05	117.97	121.00
1	B	236	TYR	CB-CG-CD2	-5.03	117.98	121.00
1	B	11	PHE	CB-CG-CD1	-5.02	117.28	120.80

There are no chirality outliers.

All (14) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	122	GLY	Mainchain
1	A	130	CYS	Peptide
1	A	191	SER	Mainchain,Peptide
1	A	37	ALA	Mainchain
1	A	55	LEU	Mainchain,Peptide
1	A	93	GLU	Peptide
1	B	130	CYS	Peptide
1	B	191	SER	Mainchain,Peptide
1	B	55	LEU	Mainchain,Peptide
1	B	93	GLU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2397	0	2375	24	0
1	B	2387	0	2362	13	0
2	A	31	0	19	0	0
2	B	31	0	19	0	0
3	A	11	0	3	2	0
3	B	11	0	3	0	0
4	A	180	0	0	2	0
4	B	178	0	0	2	0
All	All	5226	0	4781	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:17:ASN:HD21	1:A:41:ILE:H	1.26	0.81
1:B:7:ALA:H	1:B:244:GLN:HE22	1.36	0.73
1:A:131:PRO:HD2	3:A:1313:ORO:C4	2.20	0.70
1:A:7:ALA:H	1:A:244:GLN:HE22	1.37	0.69
1:A:129:SER:HB3	1:A:166:PRO:HD3	1.81	0.63
1:A:129:SER:O	1:A:131:PRO:HD3	2.01	0.61
1:B:53:ASN:HB3	1:B:212:PRO:HG3	1.83	0.60
1:A:210:ILE:HG13	1:A:215:GLY:HA2	1.85	0.57
1:B:180:ILE:O	1:B:183:GLN:HG2	2.03	0.57
1:B:210:ILE:HG13	1:B:215:GLY:HA2	1.86	0.56
1:A:195:ILE:HD12	1:A:224:ILE:HG22	1.86	0.56
1:A:53:ASN:HB3	1:A:212:PRO:HG3	1.94	0.50
1:B:51:GLU:HG3	4:B:2370:HOH:O	2.11	0.50
1:A:180:ILE:O	1:A:183:GLN:HG2	2.12	0.49
1:B:195:ILE:HD12	1:B:224:ILE:HG22	1.95	0.49
1:A:101:ALA:HB2	1:A:127:ASN:HB3	1.95	0.49
1:B:7:ALA:H	1:B:244:GLN:NE2	2.09	0.48
1:A:190:ASN:HD21	1:A:269:GLN:NE2	2.12	0.47
1:A:93:GLU:HG2	4:A:1482:HOH:O	2.13	0.47
1:A:129:SER:HB3	1:A:166:PRO:CD	2.45	0.47
1:A:94:GLY:HA2	1:A:95:PRO:HD2	1.74	0.47
1:B:94:GLY:HA2	1:B:95:PRO:HD2	1.66	0.46
1:A:137:PRO:O	1:A:138:GLN:C	2.54	0.46
1:B:1:MET:HG2	4:B:2465:HOH:O	2.15	0.46
1:A:131:PRO:HD2	3:A:1313:ORO:N3	2.32	0.45
1:A:17:ASN:H	1:A:17:ASN:HD22	1.65	0.45
1:A:16:MET:HB2	1:A:269:GLN:HG2	1.99	0.45
1:A:51:GLU:HG3	4:A:1401:HOH:O	2.18	0.44
1:B:137:PRO:O	1:B:138:GLN:C	2.56	0.43
1:A:164:LYS:HA	1:A:190:ASN:HB3	2.01	0.42
1:B:103:MET:N	1:B:107:GLU:OE1	2.45	0.42
1:A:1:MET:HG3	1:A:290:GLU:HG2	2.01	0.42
1:A:211:LYS:HB3	1:A:212:PRO:HD3	2.02	0.41
1:B:190:ASN:HD21	1:B:269:GLN:NE2	2.18	0.41
1:A:84:LEU:O	1:A:88:LYS:HG3	2.20	0.41
1:A:93:GLU:HG3	1:A:93:GLU:O	2.20	0.41
1:B:13:ASN:HB2	1:B:14:PRO:HD2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/311 (100%)	290 (94%)	17 (6%)	3 (1%)	17	7
1	B	309/311 (99%)	290 (94%)	16 (5%)	3 (1%)	17	7
All	All	619/622 (100%)	580 (94%)	33 (5%)	6 (1%)	17	7

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	92	GLN
1	B	92	GLN
1	A	93	GLU
1	B	93	GLU
1	B	136	LYS
1	A	136	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/258 (99%)	252 (99%)	3 (1%)	74	73
1	B	252/258 (98%)	250 (99%)	2 (1%)	83	83
All	All	507/516 (98%)	502 (99%)	5 (1%)	78	78

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	130	CYS
1	A	189	VAL
1	B	1	MET
1	B	189	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	86	ASN
1	A	244	GLN
1	A	269	GLN
1	B	244	GLN
1	B	269	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FMN	A	1312	-	31,33,33	3.07	14 (45%)	39,50,50	3.88	17 (43%)
3	ORO	A	1313	-	6,11,11	1.92	2 (33%)	4,15,15	8.91	3 (75%)
2	FMN	B	2312	-	31,33,33	2.57	14 (45%)	39,50,50	3.32	13 (33%)
3	ORO	B	2313	-	6,11,11	2.10	3 (50%)	4,15,15	10.54	3 (75%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	A	1312	-	-	0/16/18/18	0/3/3/3
3	ORO	A	1313	-	-	0/0/4/4	0/1/1/1
2	FMN	B	2312	-	-	0/16/18/18	0/3/3/3
3	ORO	B	2313	-	-	0/0/4/4	0/1/1/1

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1312	FMN	C1'-N10	-8.64	1.39	1.48
2	B	2312	FMN	C1'-N10	-6.96	1.41	1.48
2	B	2312	FMN	C6-C5A	-4.46	1.34	1.41
2	A	1312	FMN	P-O3P	-2.52	1.44	1.54
2	B	2312	FMN	P-O5'	-2.29	1.52	1.60
2	A	1312	FMN	C9-C9A	-2.23	1.36	1.40
2	A	1312	FMN	C6-C5A	-2.18	1.38	1.41
2	B	2312	FMN	C2-N1	-2.06	1.34	1.38
2	A	1312	FMN	C8M-C8	2.02	1.55	1.51
2	A	1312	FMN	C5'-C4'	2.06	1.54	1.51
3	B	2313	ORO	C2-N1	2.13	1.42	1.38
2	B	2312	FMN	C4A-C10	2.13	1.44	1.41
2	A	1312	FMN	O4'-C4'	2.16	1.48	1.43
2	B	2312	FMN	C4'-C3'	2.24	1.57	1.53
3	A	1313	ORO	C2-N1	2.27	1.42	1.38
2	B	2312	FMN	O4-C4	2.37	1.30	1.24
2	B	2312	FMN	C4-C4A	2.37	1.45	1.41
2	B	2312	FMN	C5A-N5	2.46	1.39	1.35
2	B	2312	FMN	C2-N3	2.75	1.43	1.38
2	A	1312	FMN	C4-C4A	2.84	1.46	1.41
2	B	2312	FMN	C4-N3	2.91	1.38	1.33
3	B	2313	ORO	C6-N1	2.93	1.39	1.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2312	FMN	C7M-C7	3.03	1.57	1.51
3	B	2313	ORO	C4-N3	3.14	1.38	1.33
2	A	1312	FMN	C2-N3	3.32	1.44	1.38
3	A	1313	ORO	C4-N3	3.58	1.39	1.33
2	B	2312	FMN	C4A-N5	4.12	1.39	1.33
2	A	1312	FMN	C9A-N10	4.27	1.43	1.38
2	A	1312	FMN	C4-N3	4.59	1.41	1.33
2	A	1312	FMN	C7M-C7	5.44	1.61	1.51
2	A	1312	FMN	C4A-N5	5.86	1.41	1.33
2	B	2312	FMN	C5'-C4'	6.06	1.60	1.51
2	A	1312	FMN	C4'-C3'	6.66	1.66	1.53

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2313	ORO	C5-C4-N3	-9.43	112.97	124.06
2	A	1312	FMN	C4A-C4-N3	-9.31	110.23	123.47
2	B	2312	FMN	C4A-C4-N3	-7.82	112.35	123.47
3	A	1313	ORO	C5-C4-N3	-7.64	115.08	124.06
2	A	1312	FMN	C6-C5A-N5	-6.69	111.28	118.95
2	A	1312	FMN	C7-C6-C5A	-5.89	112.38	121.17
2	A	1312	FMN	C4'-C3'-C2'	-5.43	101.96	113.40
2	A	1312	FMN	C4-C4A-N5	-5.36	112.68	118.70
2	A	1312	FMN	C9A-N10-C10	-5.22	114.81	121.77
2	B	2312	FMN	C4-C4A-N5	-4.20	113.99	118.70
2	B	2312	FMN	C7M-C7-C8	-3.58	113.24	120.72
2	B	2312	FMN	C1'-N10-C10	-3.28	115.18	118.46
2	A	1312	FMN	O2P-P-O5'	-3.20	98.21	106.73
2	A	1312	FMN	C7M-C7-C8	-3.17	114.08	120.72
2	B	2312	FMN	C9A-N10-C10	-2.90	117.90	121.77
2	A	1312	FMN	O3'-C3'-C2'	-2.73	102.10	108.82
2	B	2312	FMN	C4'-C3'-C2'	-2.32	108.50	113.40
2	B	2312	FMN	C8M-C8-C7	-2.23	116.06	120.72
2	A	1312	FMN	C1'-C2'-C3'	2.12	115.89	109.82
2	B	2312	FMN	C7M-C7-C6	2.12	125.58	120.36
3	A	1313	ORO	C5-C6-N1	2.29	125.11	122.38
2	A	1312	FMN	O3P-P-O5'	2.65	113.79	106.73
2	A	1312	FMN	O3P-P-O2P	2.78	118.57	107.59
2	B	2312	FMN	C5A-C9A-N10	2.81	119.86	117.71
2	A	1312	FMN	C4A-C10-N10	3.03	123.22	120.40
3	B	2313	ORO	C4-C5-C6	3.10	118.73	116.73
2	B	2312	FMN	P-O5'-C5'	4.82	131.58	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	B	2312	FMN	C4-C4A-C10	4.83	123.56	119.95
2	A	1312	FMN	C4-C4A-C10	5.44	124.02	119.95
2	A	1312	FMN	P-O5'-C5'	6.44	136.05	118.30
2	A	1312	FMN	C1'-N10-C9A	6.68	124.25	118.31
2	B	2312	FMN	C1'-N10-C9A	8.70	126.04	118.31
2	A	1312	FMN	C4-N3-C2	11.79	125.18	115.14
2	B	2312	FMN	C4-N3-C2	12.30	125.61	115.14
3	A	1313	ORO	C4-N3-C2	15.89	127.81	114.14
3	B	2313	ORO	C4-N3-C2	18.55	130.10	114.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1313	ORO	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	311/311 (100%)	-0.08	14 (4%) 33 36	10, 19, 47, 82	10 (3%)
1	B	311/311 (100%)	-0.08	14 (4%) 33 36	10, 19, 47, 82	13 (4%)
All	All	622/622 (100%)	-0.08	28 (4%) 33 36	10, 19, 47, 82	23 (3%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	133	VAL	10.8
1	A	132	ASN	10.4
1	B	93	GLU	10.3
1	A	91	ALA	9.7
1	A	90	ASN	9.6
1	A	133	VAL	9.0
1	B	132	ASN	7.9
1	B	91	ALA	7.3
1	B	90	ASN	6.5
1	A	134	PRO	6.0
1	B	130	CYS	5.9
1	A	93	GLU	5.9
1	B	89	GLU	5.0
1	A	89	GLU	5.0
1	B	136	LYS	4.7
1	B	92	GLN	4.7
1	A	88	LYS	4.5
1	A	136	LYS	4.5
1	A	135	GLY	4.4
1	A	131	PRO	4.4
1	B	134	PRO	4.1
1	B	135	GLY	4.1
1	A	130	CYS	3.8
1	A	85	LYS	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	88	LYS	3.0
1	B	131	PRO	2.9
1	A	92	GLN	2.8
1	B	137	PRO	2.6

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ORO	A	1313	11/11	0.95	0.10	13,17,19,20	0
3	ORO	B	2313	11/11	0.97	0.07	13,17,19,19	0
2	FMN	B	2312	31/31	0.97	0.08	8,12,14,15	0
2	FMN	A	1312	31/31	0.98	0.07	8,12,14,15	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.