



Full wwPDB NMR Structure Validation Report ⓘ

Feb 13, 2020 – 05:08 PM EST

PDB ID : 2JR3
Title : Antibacterial Peptide from Eggshell Matrix: Structure and Self-assembly of beta-defensin Like Peptide from the Chinese Soft-shelled Turtle Eggshell
Authors : Vivekanandan, S.; Lakshminarayanan, R.; Jois, S.D.S.; Perumal Samy, R.; Banerjee, Y.; Chi-Jin, E.O.; Teo, K.W.; Kini, R.M.; Valiyaveetil, S.
Deposited on : 2007-06-20

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.4
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 46%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	136327	12091
Ramachandran outliers	132723	10835
Sidechain outliers	132532	10811

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	42	<div> <div>50%</div> <div>29%</div> <div>• •</div> <div>17%</div> </div>

2 Ensemble composition and analysis

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:8-A:42 (35)	0.11	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 4, 9, 11, 12, 14, 15
2	3, 5, 6, 8, 10, 18, 19
3	7, 13, 16
Single-model clusters	17; 20

3 Entry composition [i](#)


There is only 1 type of molecule in this entry. The entry contains 560 atoms, of which 272 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Pelovaterin.

Mol	Chain	Residues	Atoms						Trace
1	A	42	Total	C	H	N	O	S	0
			560	177	272	51	54	6	


4.2.3 Score per residue for model 3

- Molecule 1: Pelovaterin




4.2.4 Score per residue for model 4

- Molecule 1: Pelovaterin




4.2.5 Score per residue for model 5

- Molecule 1: Pelovaterin




4.2.6 Score per residue for model 6

- Molecule 1: Pelovaterin




4.2.7 Score per residue for model 7

- Molecule 1: Pelovaterin




4.2.8 Score per residue for model 8

- Molecule 1: Pelovaterin




4.2.9 Score per residue for model 9

- Molecule 1: Pelovaterin




4.2.10 Score per residue for model 10

- Molecule 1: Pelovaterin




4.2.11 Score per residue for model 11

- Molecule 1: Pelovaterin




4.2.12 Score per residue for model 12

- Molecule 1: Pelovaterin




4.2.13 Score per residue for model 13

- Molecule 1: Pelovaterin




4.2.14 Score per residue for model 14

- Molecule 1: Pelovaterin




4.2.15 Score per residue for model 15

- Molecule 1: Pelovaterin




4.2.16 Score per residue for model 16

- Molecule 1: Pelovaterin




4.2.17 Score per residue for model 17

- Molecule 1: Pelovaterin




4.2.18 Score per residue for model 18

- Molecule 1: Pelovaterin




4.2.19 Score per residue for model 19

- Molecule 1: Pelovaterin



4.2.20 Score per residue for model 20

- Molecule 1: Pelovaterin



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.1
CYANA	refinement	2.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2jr3_cs.cif
Number of chemical shift lists	1
Total number of shifts	250
Number of shifts mapped to atoms	250
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	46%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	235	227	227	6±1
All	All	4700	4540	4540	112

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:8:CYS:SG	1:A:32:CYS:SG	0.80	2.78	10	19
1:A:18:PRO:HB2	1:A:20:VAL:HG12	0.58	1.75	14	20
1:A:22:LEU:HD21	1:A:27:HIS:CE1	0.56	2.36	17	20
1:A:29:THR:HG22	1:A:37:GLY:CA	0.55	2.32	3	17
1:A:27:HIS:CE1	1:A:29:THR:HG23	0.55	2.37	8	20
1:A:19:ILE:HG23	1:A:22:LEU:HD22	0.48	1.84	19	6
1:A:14:GLY:N	1:A:41:ILE:HD12	0.46	2.25	4	3
1:A:32:CYS:SG	1:A:38:CYS:SG	0.44	3.16	16	1
1:A:26:VAL:O	1:A:26:VAL:HG12	0.42	2.15	8	1
1:A:26:VAL:HG12	1:A:26:VAL:O	0.41	2.16	20	5

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	34/42 (81%)	23±0 (68±1%)	9±1 (27±2%)	2±1 (5±2%)	4	25
All	All	680/840 (81%)	463 (68%)	182 (27%)	35 (5%)	4	25

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	27	HIS	20
1	A	11	GLY	11
1	A	34	GLY	4

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	26/33 (79%)	20±1 (76±4%)	6±1 (24±4%)	3	27
All	All	520/660 (79%)	396 (76%)	124 (24%)	3	27

All 9 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	16	CYS	20
1	A	27	HIS	20
1	A	24	CYS	20
1	A	22	LEU	14
1	A	39	CYS	14
1	A	23	LEU	14
1	A	33	SER	11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	21	ASP	7
1	A	10	SER	4

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 46% for the well-defined parts and 46% for the entire structure.

7.1 Chemical shift list 1

File name: 2jr3_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	250
Number of shifts mapped to atoms	250
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 46%, i.e. 163 atoms were assigned a chemical shift out of a possible 354. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	68/171 (40%)	68/68 (100%)	0/70 (0%)	0/33 (0%)
Sidechain	85/155 (55%)	85/91 (93%)	0/61 (0%)	0/3 (0%)
Aromatic	10/28 (36%)	10/15 (67%)	0/11 (0%)	0/2 (0%)
Overall	163/354 (46%)	163/174 (94%)	0/142 (0%)	0/38 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 46%, i.e. 200 atoms were assigned a chemical shift out of a possible 432. 0 out of 7 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	80/204 (39%)	80/81 (99%)	0/84 (0%)	0/39 (0%)
Sidechain	110/200 (55%)	110/118 (93%)	0/76 (0%)	0/6 (0%)
Aromatic	10/28 (36%)	10/15 (67%)	0/11 (0%)	0/2 (0%)
Overall	200/432 (46%)	200/214 (93%)	0/171 (0%)	0/47 (0%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

